GENERAL INDEX—VOLUME 34

(A1 and A2 = Abstracts Supplements 1 and 2; E = Editorial; L = Letter to the Editor)

AA amyloidosis, experimental reactive proteoglycans associated with, (A1) 118 therapeutic effect of brefeldin, (A1) 118

AARONS L See Herrick AL

ABAD MA, MONTEAGUDO I, LÓPEZ-LONGO FJ, GONZÁLEZ CM, ALONSO JM, MORENO AC, ARIZA A, GONZÁLEZ M, MÍNGUEZ MD, CARREÑO L. Hepatitis C viral infection and rheumatic diseases in HIV infected patients, (A1) 111 ABAD MA See also González CM; Monteagudo I; Moreno FC

ABBOT SE, HARLEY SL STEVENS CR, BLAKE DR. Synovial microvascular endothelial cells exhibit enhanced shear rate dependent leucocyte adhesion in vitro, (A1) 51

ABBOT SE, STEVENS CR, BLAKE DR. Hypoxia induces prolonged neutrophil adhesion to endothelial cells in vitro at physiological shear rates, (A1) 50 ABBOT SE See also Sahinoglu T

Abdominal scintigraphy pattern, different in patients with ulcerative colitis, Crohn's disease and seronegative spondylarthropathies, 946 Abdominal sepsis, chronic and migratory polyarthritis, (L) 478

Abducent and trochlear palsies in a patient with Sjögren's syndrome, (L) 484 Abnormal anal function in patients with systemic sclerosis, (A1) 26

Abnormal collagen cross-linking in the cartilage of a diastrophic dysplasia patient, 512

Abnormal tear evaporation rate and osmolality in rheumatoid arthritis, (A1) 91

Absence

of correlation between IL-1a intron 6 polymorphism and rheumatoid arthritis, 1123

of exogenous retroviruses in Felty's syndrome, (L) 185

Abstracts, they turned down my abstract again, (E) 303 ACASUSO M, CISNAL A, COLLANTES E. Joint hyperlaxity: quantification

according to age, (A1) 39

ACASUSO-DIAZ M. Quantification of joint laxity. The non-dominant (Spanish) modification), (L) 795

ACCARDO S See Cutolo M

Acetabular dysplasia does not cause hip osteoarthritis in a Chinese population, (A1) 34

Acne, lupus-like syndrome in patients treated for, (L) 584

ACQUAVIVA P See Combe B

Acquired Brown's syndrome in systemic lupus erythematosus, 1092 Acupuncture

influence of knowledge, attitudes and personality characteristics on the outcome of treatment, (A2) 51 place of in medicine today, (E) 404 Acute calcific discitis. A report of two cases with scintigraphy and SPECT,

(A2) 55 Acute necrotizing vasculitis (polyarteritis nodosa?) confined to the nerve with

sponatneous recovery, (L) 694 Acute phase proteins, effect of minocycline in rheumatoid arthritis patients,

(A1) 96 Acute sarcoid arthritis: occurrence, seasonal onset, clinical features and outcome, 45

ADEBAJO A See Ali-Gombe A; Cawston TE

ADEBAJO AO, CAWSTON TE, HAZLEMAN BL. False positive IgA and IgG rheumatoid factor isotypes in infectious diseases, (A1) 109

ADEBAJO AO, MAGEED RA, HAZLEMAN BL. Rheumatoid factor—associated cross reactive idiotypes in patients with tropical infections, (A1) 109

ADEBAJO AO, WILLIAMS DG, HAZLEMAN BL, MAINI RN. Antibodies to the 65 kDa mycobacterial stress protein in West Africans with rheumatoid arthritis, tuberculosis and malaria, 352

Adhesion molecule-1 and -3, circulating intercellular, but not circulating endothelial leucocyte adhesion molecule are increased in patients with rheumatoid arthritis, 311

Adhesion molecules

LFA-1 and CD2 gene expression, differential regulation by corticosteroids in vitro, (A1) 146

soluble, altered in rheumatoid arthritis, vasculitis and systemic sclerosis, 814

special circulating levels reflect disease severity in systemic sclerosis, 1048 Adjuvant arthritis, tolerance to induced by anti-CD4 monoclonal antibodies does not confer tolerance to type II collagen-induced arthritis, (A1) 65 Adjuvant-induced arthritis, protective role for testosterone, 1117

ADU D See Luqmani RA

Adult onset Still's disease in Northern India: comparison with juvenile onset Still's disease, 429

Adult respiratory syndrome, toxic epidermal necrolysis and pancytopenia, (L)

Adult- and childhood-onset systemic lupus erythematosus: a comparison of onset, clinical features, serology and outcome, 866

Adverse events in methotrexate-treated rheumatoid arthritis patients, (S2) 49 Advice to car drivers with rheumatoid arthritis, (L) 796

ADYA CM See Uppal SS

Aerobic training, effectiveness of minimally supervised home training in patients with systemic rheumatic disease, 1064

Aetiology and presenting symptoms in male osteoporosis, 936 Aetiopathogenesis of giant cell arteritis, 960

Africa, gout in patients attending the rheumatology unit of Lomé Hospital, 843

AGGARWAL A See Agrawal S
AGRAWAL NM. Epidemiology and prevention of non-steroidal antiinflammatory drug effects in the gastrointestinal tract, (S1) 5

AGRAWAL S, AGGARWAL A, DABADGHAO S, NAIK S, MISRA R. Compound heterozygosity of HLA-DR4 and DR1 antigens in Asian Indians increases the risk of extra-articular features in rheumatoid arthritis, 41

AHMED S, DAWES PT. An audit of thyroid function tests in a rheumatic disease clinic-results and impression that thyroxine excerbates RA, (A2) 21

Ано K See Heliövaara M; Paimela L

AJMANI AK See Satoh M

ÅKESSON A See Scheja A AKINTUNDE C See Humphries J

AKIZUKI M See Satoh M

AL BALLA SR See Al Rikabi AC

AL NAHHAS A See Jawad ASM

RIKABI AC, NADDAF HO, AL BALLA SR, AL SOHAIBANI MO. Proteinaceous lymphadenopathy in a patient with known rheumatoid arthritis-case report and review of literature, 1087

AL-MUGHALES J, BLYTH TH, WILKINSON PC, HUNTER JA. The chemoattractant effect of rheumatoid synovial fluid on lymphocytes, (A1) 57 ALARCÓN GS See Sandoval DM

ALAVI A, BOND A, BROWN ST, ARDEN N, SPECTOR TD, HAY FC, AXFORD JS. Pregnancy induced glycosylation changes in rheumatoid arthritis, (A1) 62

ALAVI A See also Dalziel M; Soltys AJ

ALBANI S See Massa M

Alcoholic beverage effects on urate metabolism in gout sufferers, (A1) 120,

ALI-GOMBE A, ADEBAJO A, SILMAN A. Methodological problems in comparing the severity of rheumatoid arthritis between populations, 781 Alkaline phosphatase, bone-specific as a marker of bone turnover in osteoarthritis and rheumatoid arthritis, (A1) 56

ALLDERSEA J See Hassell A; Ollier WER

ALMOGUERA I See Alonso JC

ALONSO JC, LÓPEZ-LONGO FJ, LAMPREAVE JL, GONZÁLEZ CM, GONZÁLEZ M, ALMOGUERA I, CARREÑO L. Different abdominal scintigraphy pattern in patients with ulcerative colitis, Crohn's disease and seronegative spondylarthropathies, 946

ALONSO JM See Abad MA; González CM; Gónzalez M; López-Longo FJ Altered levels of soluble adhesion molecules in rheumatoid arthritis, vasculitis and systemic sclerosis, 814

ALVAREZ L See Peris P

Ambulatory recording, diurnal variation in 24 hour recordings of patient activity is different in rheumatoid and osteoarthritis, (A2) 16

AMES PRJ See Rosman A

5-Aminosalicylic acid, sulphasalazine or sulphapyridine-which is the active moiety in rheumatoid arthritis?, (S2) 16

AMOR B See Claudepierre P

Amos M See Srinivasan A

AMOS N See Williams AS

Amos RS. The history of the use of sulphasalazine in rheumatology, (S2) 2 Anal function, abnormal in patients with systemic sclerosis, (A1) 26 Analysis

of cell types and mediator production from tissues around loosening joint implants, 1127

of childhood relationships in adults with juvenile chronic arthritis, (A1) 90

of Chlamydia trachomatis specific synovial fluid derived T cell clones from a patient with recurent reactive arthritis, (A1) 152 of the discrepancy between the clinical and radiological course in

rheumatoid arthritis, (A1) 105 thrombotic events in antiphospholipid antibody positive lupus patients, (A1) 15

ANCA, antigen specificity in systemic vasculitis, (L) 394

Anchor drug role, how and when should combination therapy be used?, (S2)

ANDERSON FH See Francis RM

ANDERSON JD See Dobbie JW

ANDERSON ME, MOORE TL, JAYSON MIV, HERRICK AL. Neurovascular control mechanisms in primary Raynaud's phenomenon and systemic sclerosis as assessed by iontophoresis and laser Doppler techniques, (A2) 24

ANDERSONE D See Stichtenoth DO

ANDERSSON R See Schaufelberger C ANDREOTTI F See Mackworth-Young CG

ANDREWS D See Gough A

ANDRUP O See Glennås A

Angiotensin converting enzyme

and von Willebrand factor level value in vasculitis, (A1) 22

and von Willebrand factor release kinetics by endothelial cells, (A1) 23 Ankle

and foot in rheumatoid arthritis, 443

surgery to, 854

Ankylosing spondylitis

antibodies to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of patients with axial and peripheral form, 413 antibodies to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of patients with/without iritis and enthesitis, 418

atypical antineutrophil cytoplasmic antibodies, (L) 695

axial skeletal changes in paralysed patients may mimic, 174 Bath Ankylosing Spondylitis Functional Index, (L) 793, (L) 794

Bath Ankylosing Spondylitis Radiology Index (BASRI): a new validated approach to disease assessment, (A2) 37

Bath patient global score, (A1) 114

the body chart: a meaningful assessment of pain compared to visual analogue scales, (A2) 36

the correlation between a new metrology score and radiology, 767

diagnostic features, (E) 301

early, bone density and bone turnover, (A2) 34

effect of smoking on outcome, (A2) 35

effect of syndesmophyte formation on lumbar spine bone density in patients with ankylosing spondylitis, (A1) 117

high prevalence of apical pulmonary fibrosis by high resolution computerized tomography, (A1) 117

HLA class II associations, (A1) 142

HLA component of the genetic contribution, (A1) 76

the individual with: defining disease status and the impact of the illness, 663

juvenile onset—more girls than we thought?, (A2) 34 nature of surgical intervention, (A1) 115

outcome of total hip replacement: a thirteen year follow-up study, (A1)

relationship of fatigue to disease activity, sleep and other factors, (A2) 36 risk factors for a patient with spondylarthropathy to develop, (A1) 116 single photon emission computerised tomography identifies pain sites,

tarsal tunnel syndrome, (L) 488

and tonsillectomy, (A2) 35

use of focus group interview in the evaluation of patients, 150

ANÓN J See Tinahones FJ

Anterior cruciate ligament

quadriceps activation and rehabilitation following rupture, (A1) 135 shortly after transection dogs develop characteristics similar to human moderate knee osteoarthritis, (A1) 135

Anterior knee pain in thalassaemia major treated with L1, a new iron chelating agent, (A1) 119

Anti ganglioside antibodies in patients with rheumatoid arthritis and peripheral neuropathy, (A2) 3

Anti-Clq antibodies are associated with hypo-complementaemia, defective opsonisation of gram-positive organisms, and death due to infection in patients with SLE, (A1) 10

Anti-cardiolipin antibodies

and lupus anticoagulant, systemic lupus erythematosus patients with both are at risk for clinical manifestations related to the antiphospholipid syndrome, 873

thrombosis is not associated with activated protein-C resistance due to factor VLeiden mutation, (A1) 15

Anti-CD4 chimaeric monoclonal antibody cM-T412, human anti-chimaeric antibody response in rheumatoid arthritis patients after treatment with,

Anti-CD4 monoclonal antibodies, tolerance to adjuvant arthritis induced by

does not confer tolerance to type II collagen-induced arthritis, (A1) 65
Anti-CD44 antibody induces receptor shedding from synovial leukocytes and fibroblasts from rheumatoid arthritis patients, (A2) 30

Anti-endothelial cell antibodies

and anti-neutrophil cytoplasmic antibodies are not increased in Kawasaki disease, 882

in inflammatory myopathies: association with interstitial lung disease.

Anti-endothelial cell membrane antibodies, characterization in systemic lupus erythematosus, (A1) 11

Anti-rheumatic drug

audit of the outpatient use of disease modifying and immunosuppressive, (A1) 103

disease-modifying, use of a contrast enhanced MRI in the assessment of therapeutic response, 956

Anti-ribonucleoprotein antibodies

cross-sectional and longitudinal study in lupus nephropathy, (A1) 8 precipitating and non-precipitating, immunoblot analysis, (A1) 11

Anti-RNA polymerase antibodies, patients with have diffuse scleroderma and exhibit a speckled pattern anti-nuclear antibody distinct from anti-Scl-70, (A1) 66

Anti-Ro antibodies, homozygosity for glutathione-transferase GSTM1 and GSTT1 null alleles may confer risk for in SLE, (A1) 5

Anti-Ro/SS-A antibody association with progressive multifocal leukencephalopathy in HIV infected patients, (A1) 112

Anti-Sa antibodies in patients with systemic lupus erythematosus, (A1) 9 Anti-Sm antibodies: a potential risk factor for lupus nephritis, (A1) 9

Anti-thyroid drug treatment, antineutrophil cytoplasmic antibody-positive crescentic glomerulonephritis associated with, 1090

Anti-TNFα antibody CDP571

engineered, repeated dose in patients with rheumatoid arthritis, (A1) 81 engineered human, serological effects following treatment in patients with rheumatoid arthritis, (A1) 101

Antibiotics, role of prophylactic in the prevention of late infection of prosthetic joints. Results of a questionnaire and review of the literature, 380

Antibodies

to the 65 kDa mycobacterial stress protein in West Africans with rheumatoid arthritis, tuberculosis and malaria, 352

to heat shock proteins, increased levels with increasing age in MRL/MP-LPR/LPR mice, 610

to a human endogenous retrovirus envelope glycoprotein in rheumatoid arthritis, (A1) 154

to intermediate filaments and human immunodeficiency virus, (A1) 112 to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of ankylosing spondylitis patients with/without iritis and enthesitis, 418

to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of patients with axial and peripheral form of ankylosing spondylitis, 413

Antibody dependent cellular cytotoxicity (ADCC) inhibition in rheumatoid arthritis by the non-immune complex of IgA-alpha-1-antitrypsin, (A2) 30 Antigen specificity of ANCA in systemic vasculitis, (L) 394

Antimalarials, incidence of myopathy in patients treated with. A report of three cases and a review of the literature, 166

Antineutrophil cytoplasmic antibodies

and anti-endothelial cell antibodies are not increased in Kawasaki disease, 882

atypical in patients with ankylosing spondylitis, (L) 695 in inflammatory arthritis-potential for misdiagnosis, 820

Antineutrophil cytoplasmic antibody-positive crescentic glomerulonephritis associated with anti-thyroid drug treatment, 1090

Antinuclear antibodies in early onset pauciarticular juvenile chronic arthritis (JCA) are associated with HLA-DQB1*0603: a possible JVA-associated human leucocyte antigen haplotype, 461 Antiperinuclear factor

specificity for rheumatoid arthritis in rheumatoid factor-positive sera,

in spondylarthritis, 912

Antiperinuclear factor-IgG, high diagnostic value: prospective analysis of 1004 sera diluted 1:100 and more, (L) 87

Antiphospholipid antibodies in Hughes syndrome: move closer to thrombosis in 1994, (E) 493

in paediatric systemic lupus erythematosus, juvenile chronic arthritis and overlap syndromes: SLE patients with both lupus anticoagulant and high-titre anticardiolipin antibodies are at risk for clinical manifestations related to the antiphospholipid syndrome, 873

Antiphospholipid syndrome

and endothelium-derived haemostatic factors, 201

prevalence and clinical significance of subungual splinter haemorrhages, (A1) 16

primary

aortitis as a new manifestation, (L) 686

development of systemic lupus erythematosus features, (A1) 16 increased levels of β 2 glycoprotein-1 antigen and β 2 glycoprotein-1 binding antibodies are associated with a history of thromboembolic complications, 1031

Antirheumatic agents sulphasalazine and methotrexate share an antiinflammatory mechanism, (S2) 30

Antirheumatic therapy, effect of 2nd line on pyridinium crosslink excretion in patients with established RA, (A1) 99

Antiribosomal P protein antibody detection in patients with systemic sclerosis, 908

Antithrombin III inactivation in rheumatoid arthritis, (A1) 84

Aortitis: a new manifestation of primary antiphospholipid syndrome, (L) 686 Apoptosis

from worm to wonder drug, (E) 95

pathological synovial fluids promote apoptosis of peripheral blood and synovial neutrophilis: a role in resolution of synovitis?, (A1) 58 APPELBOOM T. See Wile G

ARC Low Dose Corticosteroid Study Group See Kirwan JR

ARCHER JR. Seronegative arthritis mailbase discussion group, (L) 1185

ARDEN N See Alavi A; Dolan AL

ARDEN NK, BAKER J, BAAN K, SPECTOR TD. The heritability of hip axis length and ultrasound of the calcaneus in postmenopausal females: a twin study, (A1) 128

ARDEN NK See also Keen RW; Lloyd ME

Are slow-acting anti-rheumatic drugs monitored too often? An audit of current clinical practice, 966

ARINGER M See Koeller M

ARIZA A See Abad MA; González CM; González M; Monteagudo I

ARMSTRONG S See Munro R

ARNAIZ-VILLENA A See Pablos JL

ARNOLD MM, MCKENNA F. A double blind comparison of the endoscopic and clinical effects of tenoxicam and diclofenae in rheumatoid arthritis, (A1) 95

AROMAA A See Heliövaara M

Arthritic disease identification, fluorophore assisted carbohydrate electrophoresis: a potential tool, (A1) 61

Arthritis

associated with HIV infection in Zambia, (A2) 32

brucellar, pseudogout masking, (L) 294 chronic nephrotoxicity in the treatment, 162

education programme, five-year follow-up of a controlled trial, 647

hepatitis C virus (HCV)-associated. Quantitative analysis of HCV RNA of the synovial fluid and the serum, (L) 691

lymphocyte reactivity to Clostridium perfringens antigens and toxoid, (L) 1095

panniculitis and pancreatitis, 680

receiver operating characteristic analysis of urinary pyridinium crosslinks, (L) 792

recent-onset inflammatory, muscle strength in healthy people and in patients suffering from, 355

in scleroderma, 831

seronegative mailbase discussion group, (L) 1185

sexually acquired in nineteenth century London, (A2) 31

staphylococcal in human immunodeficiency virus infection, (L) 397 Arthrogenic inhibition determinants in knee osteoarthritis, (A1) 134

Arthropathy, leucopenia and recurrent infection associated with a TcR $\gamma\delta$ population, 978

Arthroplasty, lower limb, second-line drug experience of patients with rheumatoid arthritis, (A2) 20

Arthroscopy, needle, (E) 1102

Articular cartilage

compressive modulus measurement under physiological loading conditions, (A2) 48

human metalloproteinase release and cyclical pressure from, (A2) 28
Articular damage severity, prognostic factors in rheumatoid arthritis.

Comparative study of two subsets of patients according to severity of

damage, 529

Articular indices
comparison of the validity and reliability of self-reported, 760
joints count: a review of old and new of joint inflammation, (E) 1003

Aseptic necrosis, cutaneous, autoimmune phenomena associated with during interferon-alpha treatment for chronic myelogenous leukaemia, (L) 582 ASHERSON RA See Khamashta MA

ASHERSON KA See Khamashta MA
Assessing clinical competence: recognition of case descriptions of rheumatic

diseases by general practitioners, 375

Assessing systemic lupus erythematosus activity using plasma factor VIII von Willebrand factor antigen, (A1) 8

Assessment

of bone in Ehlers Danlos syndrome by ultrasound and densitometry, (A1) 130

of bone mineral density in women with Marfan syndrome, 516 of primary Sjögren's syndrome in patients with dry eyes, (A2) 25

of referral guidelines to a bone densitometry service, (A1) 37
Association

of anti-Ro/SS-A antibodies with progressive multifocal leukencephalopathy in HIV infected patients, (A1) 112

between physical function and psychological problems in children with juvenile chronic arthritis, 470

of HLA-DR4, protease inhibitor phenotypes and keratoconjunctivitis sicca with pulmonary abnormalities in rheumatoid arthritis, 37

Asymptomatic mothers of babies with congenital heart block with features of Sjögren's syndrome, (A1) 19

ATHANASSIOU P, IOAKIMIDIS D, WESTON J, GEORGANAS K, KONTOMERKOS T, DAVIES KA. Type I cryoglobulinaemia associated with a thymic tumour: successful treatment with plasma exchange, 285

Athletes, female postmenopausal, vitamin D receptor gene polymorphisms, (A1) 79

Athritis, active inflammatory, lymphoedema associated with in a patient with congenital lymphatic hypoplasia, (L) 689

ATKINS RM See Perry MJ

Atlanto-occipital subluxation in rheumatoid arthritis demonstrated by magnetic resonance imaging, 787

ATRA E See Bosi Ferraz M; Cury SE

ATSUMI T See Rosman A

Atypical antineutrophil cytoplasmic antibodies in patients with ankylosing spondylitis, (L) 695

Audit

are slow-acting anti-rheumatic drugs monitored too often? An audit of current clinical practice, 966

of methotrexate and folic acid for rheumatoid arthritis. Experience from a teaching centre, 971

of the outpatient use of disease modifying and immunosuppressive antirheumatic drugs, (A1) 103

of patients taking sulphasalazine and penicillamine individually, sequentially and concurrently in inflammatory arthritis, (A2) 17

preliminary of the management of acute low back pain in the Kettering district, (L) 693

regional of adult onset Still's disease, (A2) 22

shared care between hospital and general practice: disease-modifying drug monitoring in rheumatoid arthritis, 673

of thyroid function tests in a rheumatic disease clinic—results and impression that thyroxine excerbates RA, (A2) 21

AUDRAIN M See Berthelot JM

AUNG TS, PLANT MJ, RAHMATALLA AA, WYNN-JONES CH, WARD AB, DAWES PT. Measurement of shoulder movement using FASTRAK 3-D tracking system, (A2) 47

Australia, letter from, 57

Autoantibodies

changing profiles with variable clinical manifestations in a patient with relapsing systemic systemic lupus erythematosus and polymyositis, 915

to a major epitope on the carboxyl terminal fragment of CENP-B in patients with autoimmune disease, 407

to myeloperoxidase in systemic sclerosis, (A2) 24

Autoimmune abnormalities, sulphasalazine-induced in patients with rheumatic disease, 435

Autoimmune antibodies, sulphasalazine-induced in patients with rheumatic disease, (L) 1097

Autoimmune conditions, decreased DNA repair proficiency: a factor in susceptibility or a feature of disease?, (A1) 4

Autoimmune phenomena associated with cutaneous aseptic necrosis during

interferon-alpha treatment for chronic myelogenous leukaemia, (L) 582

Autoimmune rheumatic disease, new therapeutic approaches in, with special

emphasis on rheumatoid arthritis, (E) 193 Autoimmunity, mercuric chloride-induced, oxypentifylline enhances the Th2 environment and inhibits TNFα mRNA production and inflammatory

arthritis, (A2) 29
Automimmune disease, frequency of autoantibodies to a major epitope on the carboxyl terminal fragment of CENP-B, 407

Avascular necrosis, multiple joint: beware of tuberculosis and human immunodeficiency virus—a rare but important cause, 387

AVERNS HL, CHEUNG NT, DAWES PT. Painful knees associated with a septic focus 86?

AVERNS HL, KERRY R. Role of prophylactic antibiotics in the prevention of late infection of prosthetic joints. Results of a questionnaire and review of the literature, 380

AVERNS HL, OXTOBY J, TAYLOR HG, DZIEDZIC K, JONES PW, DAWES PT. Effect of smoking on outcome of ankylosing spondylitis, (A2) 35

AVINA-ZUBIETA JA, JOHNSON ES, SUAREZ-ALMAZOR ME, RUSSELL AS.
Incidence of myopathy in patients treated with antimalarials. A report of three cases and a review of the literature, 166

Awad M, Corrigall VM, Dayer J-M, Panayi GS. Effect of gold salts on monocytes and their pattern of cytokine secretion, (A1) 150

AXFORD JS See Alavi A; Dalziel M; Martin K; Rees DHE; Soltys AJ

Axial skeletal changes in paralysed patients may mimic ankylosing spondylitis, 174

AYRAL X. Radiological medial joint space narrowing and meniscal lesions, (L) 991 β 2 glycoprotein-1 antigen and β 2 glycoprotein-1 binding antibodies, increased levels are associated with a history of thromboembolic complications in patients with SLE and primary antiphospholipid syndrome, 1031

BAAN K See Arden NK; Etherington J

BAARN K See Dolan AL

Back pain

inflammatory in primary care, 1074

influence of physical, psychological and behavioural factors on consultations, 156

more than pain in the back II: tender points in graded back pain, (A2) 38 relationship with psychological status and quality of life in women in the general population: the Chingford study, (A1) 40

BACON PA, FARR M, BIRD H, HICKLING P, SWANNELL A, TEALE R. A double blind placebo controlled trial of minocycline in patients with active rheumatoid arthritis, (A2) 16

BACON PA See also Farr M; Gordon C; Gough A; Kitas GD; Luqmani RA;

McRorie ER; Waterhouse L; Zwede M

Bacterial flora-derived antigen presence in synovial tissue macrophages and dendritic cells, 1112

Bacterial joint infection in England & Wales 1990-1993, (A2) 31

Bacterial overgrowth, small bowel in systemic sclerosis: detection using direct and indirect methods and treatment outcome, 265

BADCOCK LJ, CLARKE S, FISHER J, JONES P, DAWES PT. Rheumatoid arthritis and primary selective IgA deficiency, (A2) 3

BAGGIOLINI M See Seitz M
BAILDAM EM, HOLT PJL, CONWAY SC, MORTON MJS. The association between physical function and psychological problems in children with juvenile chronic arthritis, 470

BAILEY AJ, SIMS TJ, STANESCU V, MAROTEAUX P, STANESCU R. Abnormal collagen cross-linking in the cartilage of a diastrophic dysplasia patient, 512

BAILEY I See Vaghmaria A

BAILEY R See Jones HW

BAILLY S, HAYEM G, FAY M, KAHN MF, GOUGEROT-POCIDALO MA Absence of correlation between IL-1a intron 6 polymorphism and rheumatoid arthritis, 1123

BAJOCCHI GL, PITZALIS C, HALL M, LEE A, PIPITONE N, LANCHBURY J, PANAYI G. Differential regulation by corticosteroids of LFA-1 and CD2 adhesion molecule gene expression in vitro, (A1) 146

BAJOCCHI GL See also Pitzalis C BAKER J See Arden NK BAKER JR See Keen RW

BAKER LRI See D'Cruz D

BALINT G. Sixth European Regional Conference of Rehabilitation International, 283

BAMBER S See Brennan P; Macgregor AJ; Newman J BANKHEAD C See Brennan P; Harrison BJ; Jones PBB

BANKHEAD CR, SYMMONS DPM, BARRETT EM, SCOTT DGI, SILMAN AJ. The incidence of rheumatoid arthritis is not related to socioeconomi factors, (A2) 7

BANKHEAD CR, SYMMONS DPM, JAYSON MIV. Continuing medical education for rheumatologists: the current picture, 172

BARBANO GC See Gattorno M

BARILE L See Medina F

BARLOW JD See Herrick AL

BARON D See SARAUX A BARONE A See Cutolo M.

BARRERA P, HAAGSMA CJ, BOERBOOMS AMTh, VAN RIEL PLCM, BORM GF, VAN DE PUTTE LBA, VAN DER MEER JWM. Effect of methotrexate alone or in combination with sulphasalazine on the production and ciculating concentrations of cyokines and their antagonists, 747

BARRETT B See Jones M BARRETT E See Brennan P

BARRETT EM See Bankhead CR; Harrison BJ

BARROW S See Wright D

BARRY M See Barry R

BARRY R, COLLINS M, SMITH M, HAZLEMAN B, BARRY M. Feasibility study of homeopathic remedy in rheumatoid arthritis, (A2) 17
BARTÓK I, VYSE TJ, SPÄTH PJ, MORLEY BJ, WALPORT MJ. The molecular

basis of hereditary complement factor I deficiency, (A1) 155 BARTRAM SA, DENTON CP, DU BOIS RM, BLACK CM. Use of intravenous

prostacyclin to treat pulmonary hypertension associated with systemic sclerosis, (A1) 30

Basic calcium phosphate salts and supraspinatus tendon degeneration, (A1)

BASSIOUNI H See Troughton PR

BASSIOUNI M See Troughton PR Bath Ankylosing Spondylitis

Functional Index, (L) 793, (L) 794 patient global score, (A1) 114

Radiology Index (BASRI): a new validated approach to disease assessment, (A2) 37

BAUMAN A See Lindroth Y

BAVINGTON C See Jobanputra P; Wright M

BAXENDALE RH See Hall MG

BAYLISS M See Lewis S; Yaqub R

BCG immunotherapy for bladder carcinoma, psoriatic arthropathy after, (L) 1097

BCL-2 expression is unaltered in unfractionated peripheral blood mononuclear cells in patients with systemic lupus erythematosus, 316

BECKER BE. Home exercises and out-patient hydrotherapy, (L) 991

BEEVERS DJ, SEEDHOM BB, KAY SPJ. Design of a non constrained, non cemented, modular metacarpophalangeal prosthesis, (A2) 47

BEHAN WMH See Neilly JB

Behçet's disease

in India: a clinical, immunological, immunogentic and outcome study, 825

Lhermitte's sign, (L) 184

and pregnancy relationship study, (A1) 24

sensitivity and specificity of different diagnostic criteria according to the latent class approach, 932

skin hyper-reactivity in patients with benign intracranial hypertension as

an early manifestation, (L) 184
BELCH JJF See Maple C; Mulherin D; Veale DJ

BELCHER C, FAWTHROP F, DOHERTY M. Levels of urokinase and plasminogen activator inhibitor-1 in synovial fluids in normal, osteoarthritis and rheumatoid arthritis knees, (A1) 59

BELL A See Doherty E

BELL AL See Bruce IN; Magill MK

BELZUNEGUI J, PLAZAOLA I, URIARTE E, PEGO JM. Primary tuberculous muscle abscess in a patient with systemic lupus erythematosus, 1177

Bends, polyarthritis masquerading as, (A2) 53 BENGTSSON B-Å See Schaufelberger C

BENKE-STUDNICKA A See Erlacher L BENLAHRACHE S See Kennedy LG

BERKSON L See Spitzer WO BERNARD J-C See Combe B

BERNARDS ATM See Hogeweg JA

BERNSTEIN RM See Brammah TB; Kay RA

BERRETTINI S, FERRI C, LA CIVITA L, SEGNINI G, LOMBARDINI F, BRUSCHINI P, LONGOMBARDO G, SELLARI-FRANCESCHINI S. Inner ear involvement in mixed cryoglobulinaemia patients, 370

BERRY H See Houssein DA; Houssien DA; Scott DL; Smith EC

BERTHELOT JM. High diagnostic value of antiperinuclear factor-IgG: prospective analysis of 1004 sera diluted 1:100 and more, (L) 87

BERTHELOT JM, MAUGARS Y, AUDRAIN M, YOUINOU P, PROST A. Specificity of antiperinuclear factor for rheumatoid arthritis in rheumatoid factorpositive sera, 716

BEYELER C, IM HOF V, JORDI B, GERBER NJ. Pulmonary function in rheumatoid arthritis treated with low dose methotrexate: a longitudinal study, (A1) 100

BEYNON HLC See Davies KA

BHAKTA BB, PEASE CT. Rheumatoid arthritis in the elderly: is peripheral oedema at onset a good prognostic feature?, (A2) 10

BHALLA A See Patel S

BHALLA AK See Elvins DM; McFarlane XA

BHASKARAN V See Thompson PW

BHATT B See Sahinoglu T

BICKERSTAFF MCM See Denton CP

BIENVENU J See Tebib JG

BIGG HF, HARRALL RL, RILEY GP, CAWSTON TE. The effect of growth factors on collagenase and TIMP production from human tendon fibroblasts in culture, (A1) 138

BULSMA JWJ See Geenen R; Lems WF; van Valburg AA

BILLINGHAM MEJ, CARNEY SL, HICKS CA, KINGSTON AE. Tolerance to adjuvant arthritis induced by anti-CD4 monoclonal antibodies does not confer tolerance to type II collagen-induced arthritis, (A1) 65

BINKS MH, KATZ DR, CHAIN BM. In vitro maturation of human peripheral blood dendritic cells and the extracellular matrix environment, (A1) 145

BIRCH A See James IT BIRD H See Bacon PA

BIRD HA See Troughton PR

BIRD HA. Sulphasalazine, sulphapyridine or 5-aminosalicylic acid-which is the active moiety in rheumatoid arthritis?, (S2) 16

BIRK MA, ZEUTHEN EL. Phrenicus palsy in progressive systemic sclerosis, 684

BIRRELL FN, SHEERAN TP. A case of CREST syndrome with hypermobility: an unusual association, (A2) 54

Bits and pieces in a puzzle-rheumatoid arthritis and pregnancy, (E) 1

BJARNASON I See Mahmud T; Rafi S BJERREGARD K See Drewes AM

BJÖRK J See Smedegård G

BLAAUW AAM, SCHUWIRTH LWT, VAN DER VLEUTEN CPM, SMITS F, VAN DER LINDEN Sj. Assessing clinical competence: recognition of case descriptions of rheumatic diseases by general practitioners, 375

BLACK A See Oien KA

BLACK C See Marguerie C

BLACK CM, DENTON CP. The management of systemic sclerosis, (E) 3
BLACK CM See also Bartram SA; Bunn CC; Carulli MT; Denton CP; Kaye SA; Misra R; Shiwen X; Silman AJ; Vancheeswaran R; Worrall JG

BLADES S See Gaffney K BLAKE D See Gaffney K

BLAKE DR See Abbot SE; Dai L; Jawed S; Jones HW; Naughton D; Nazhat NB; Sahinoglu T; Winrow VR; Zhang Z

BLANCO FJ See Gonzalez-Crespo MR

BLANN AD, HERRICK A, JAYSON MIV. Altered levels of soluble adhesion molecules in rheumatoid arthritis, vasculitis and systemic sclerosis, 814 BLYTH TH See Al-Mughales J

BODART C See Dobbie JW

BODEN T See Ward PS

BODMAN KB See Rankin ECC

Body chart: a meaningful assessment of pain compared to visual analogue scales in ankylosing spondylitis, (A2) 36

Body image, impact of rheumatoid arthritis on, (A1) 89

BOERBOOMS AMTh See Barrera P

BOERSMA JW See Griep EN

BOLOGNA C See Combe B; Jorgensen C

BOLTEN W See Wylie G

BOND A See Alavi A: Dalziel M: Martin K

Bone

assessment in Ehlers Danlos syndrome by ultrasound and densitometry, (A1) 130

early postmenopausal loss and polymorphisms of the vitamin D receptor, (A1) 79

effect of high-dose corticosteroid pulse therapy on metabolism in rheumatoid arthritis, (A1) 128

structure and function, near infra-red laser in the assessment of, (A2) 42 three-phase scan use in the diagnosis of repetitive strain injury, (A2) 39 Bone densitometry

assessment of referral guidelines to service, (A1) 37

hand in early rheumatoid arthritis: a measure of progression despite clinical control, (A1) 108

Bone density

and bone turnover in early AS, (A2) 34 effect of thiazide use: the Chingford study, (A1) 129

lumbar spine, syndesmophyte formation effect on in patients with ankylosing spondylitis, (A1) 117

and serum electrolytes: the Chingford study, (A1) 129

utility of lateral measurements in osteoarthritis of the lumbar spine, (A1)

and vitamin D receptor gene alleles in postmenopausal twins, (A1) 82 Bone marrow transplant, rheumatological complications, (A1) 120

peak in women, life-style, environmental and medical factors influencing, 620

pre-menopausal women, effect of systemic lupus erythematosus and long-term steroid therapy, 742

Bone mineral density

assessment in women with Marfan syndrome, 516

axial and intestinal strontium absorption, (A1) 75

can changes in appendicular bone predict it during treatment with hormone replacement therapy and cyclical etidronate?, (A2) 41

changes with bisphosphonate treatment of Paget's disease of bone, (A1) 130

and exercise: a comparison of female ex-elite athletes and aged matched controls, (A1) 72

serial changes in polymyalgia rheumatica, (A1) 74

Bone-specific alkaline phosphatase as a marker of bone turnover in osteoarthritis and rheumatoid arthritis, (A1) 56

BONIFACE F. How does neurophysiological assessment influence the management and outcome of patients with carpal tunnel syndrome?, (L) 581

BOONEN S, VAN DISTEL G, WESTHOVENS R, DEQUEKER J. Steroid myopathy induced by epidural triamcinolone injection, 385

BOOT-HANDFORD RP See Hillarby MC BORG AA, DAVIS MJ, JONES PW, DAWES PT. Measurement of pain in rheumatoid arthritis, (A1) 91

BORG AA, PLANT MJ, SAKLATVALA J, CLARKE D, MATTEY D, DAWES PT. Wrist predominance in seronegative rheumatoid arthritis: fact, fiction or natural history, (A2) 9
BORG AA. Sulphasalazine-induced autoimmune abnormalities in patients

with rheumatic disease, (L) 1097

BORG AA See also Kay LJ

BORM GF See Barrera P BOSCH A. Skin hyper-reactivity in patients with benign intracranial hypertension as an early manifestation of Behçet's disease, (L) 184

BOSCH JA See Marsal S BOSCH-GIL JA. Churg-Strauss syndrome with inflammatory orbital pseudotumour, (L) 485

BOSI FERRAZ M, WALTER SD, HEYMANN R, ATRA E. Sensitivity and specificity of different diagnostic criteria for Behçet's disease according to the latent class approach, 932

BOSSINGHAM D. Bush rheumatology or rheumatologist gone bush?, 785

Вотто M See Naughton M

BOUMBA D, SKOPOULI FN, MOUTSOPOULOS HM. Cytokine mRNA expression in the labial salivary gland from patients with primary Sjögren's syndrome, 326

BOURGEOIS P See Kaye O BOURKE BE See Laversuch CJ

BOUVIER M See Tebib JG

BOWDEN A See Herrick AL BOWDEN AR See Seed MP

BOWLES J See Dasgupta B BOYD MT See Venables PJW

BOYLSTON A See Ross DA

BRABYN J See Deodhar AA

Brachial plexitis in a patient with acute parvovirus B19 infection, (L) 480

BRADBURY S See Vaghmaria A
BRADLEY SM, LE GALLEZ P, GOOI HC, TROUGHTON PR. The effect of sulphasalazine on neutrophil superoxide protection in RA, (A2) 14

BRADLOW A. Continuing medical education in rheumatology, (E) 98

BRADLOW A See also Murphy K

BRADSHAW C, WATLING B, BRYCE C, STEEN IN. Manipulative physiotherapy for spinal problems in primary care: outcomes of care, 1070 BRADY G See Hillarby MC

Bradykinin-induced intra-articular plasma extravasation, neurogenic contribution in the rat knee joint, (A1) 52

Bramley R See Etherington J
Brammah TB, Guy M, Cooper RG, Holt L, Bernstein RM, Jayson MIV. Urinary retinol binding protein excretion in SLE, (A1) 17

BRANDER ML See Buckley MG

BRAZIER DJ See Forbes JE; Whitefield L

Breast feeding, other reproductive factors and rheumatoid arthritis. A prospective study, 542

BREATHNACH D, COSTELLO P, MULHERIN D, BRESNIHAN B, FITZGERALD O. Cutaneous lymphocyte antigen expression by peripheral blood and synovial fluid lymphocytes in psoriasis and psoriatic arthritis, (A1) 147

BREATHNACH D, COSTELLO P, MULHERIN D, BRESNIHAN B, FITZGERALD O. HML-1 expression by peripheral blood and synovial fluid lymphocytes in rheumatoid and psoriatic arthritis, (A1) 148

BREATNACH ES See Doran M

Breedveld FC See Kloppenburg M; ten Wolde S; van den Borne BEEM; van den Ende CHM; Vliet Vlieland TPM; Voskuyl AE; Wylie G

Brefeldin, therapeutic effect on experimental reactive AA amyloidosis, (A1)

BRENNAN FR, MIKECZ K, JOBANPUTRA P, GLANT TT, NUKI G. An anti-CD44 antibody induces receptor shedding from synovial leukocytes and fibroblasts from rheumatoid arthritis patients. (A2) 30

BRENNAN P, BANKHEAD C, SILMAN A, BARRETT E, SCOTT DGI, SYMMONS DPM. Predictors of disability one year after presentation with inflammatory polyarthritis, (A1) 86

BRENNAN P, SILMAN AJ, PAYTON A, BAMBER S, OLLIER WER. Foetally inherited paternal HLA does not explain post partum susceptibility to rheumatoid arthritis in the mother, (A1) 142

BRENNAN P See also Silman AJ

Bresnihan B See Breathnach D; Doherty E; Eustace J; Mulherin D BRESSOLLE F See Combe B

British Paediatric Association report on paediatric rheumatology, 60

British Paediatric Rheumatology Group news, 59, 453

BROAD I See Etherington J

Broadband ultrasound characteristics in rheumatoid arthritis and diabetes, (A1) 123

BRONS MR See Geenen R

BROOKES SM, PRICE EJ, VENABLES PJW, MAINI RN. Interferon-gamma and epithelial cell activation in Sjögren's syndrome, 226

BROOKES SM See also Venables PJW
BROOKS P, KIRWAN JR. Evidence-based medical practice: the Cochrane collaboration and osteoarthritis, (E) 403

BROOKS PM. Letter from Australia, 57

BROOKS PM See also Lindroth Y

BROPHY D See Eustace J Brown GMM See Kalla AA

BROWN K See De Lord D

Brown M. Transient regional osteoporosis of the hip, (L) 296

BROWN MA, PILE KD, GIBSON K, KENNEDY LG, CALIN A, FARRALL M, WORDSWORTH BP. The HLA component of the genetic contribution to ankylosing spondylitis, (A1) 76

BROWN MA, PILE KD, KLINKE J, GIBSON K, BUNCE M, CALIN A, DARKE C, DAVIS J. HLA class II associations of ankylosing spondylitis, (A1) 142 Brown MM See Laversuch CJ; Rees DHE

BROWN ST See Alavi A

Brown's syndrome, acquired in systemic lupus erythematosus, 1092

BRUCE IN, JOHNSTON GD, HARRIS CM, McNALLY JA, BELL AL. Increased endothelium derived relaxing factor/nitric oxide production in patients with systemic vasculitis, (A1) 23

Brucellar arthritis, pseudogout masking, (L) 294

BRUCKNER FE See Johnson SM BRÜHLMANN P See Stucki G

BRUN JG, NILSSEN S, KVALE G. Breast feeding, other reproductive factors and rheumatoid arthritis. A prospective study, 542

BRUSCHINI P See Berrettini S

BRYAN C See Silman AJ

BRYCE C See Bradshaw C

BUCHANAN N See Etherington J; Lima F

BUCHANAN NMM, TOUBLE, KHAMASHTA MA, LIMA F, KERSLAKE S, HUGHES GRV. The safety of hydroxychloroquine in lupus pregnancy: experience in 27 pregnancies, (A1) 14

BUCKLAND-WRIGHT JC, MACFARLANE D, WILLIAMS S, WARD R. Joint space width measured more accurately and precisely in semi-flexed than extended view of OA knees, (A1) 121

BUCKLAND-WRIGHT JC See also Lynch JA

BUCKLEY MG, WALTERS C, BRANDER ML, WONG R, CAWLEY MID, WALLS AF. Mast cell tryptase in the synovial fluid of patients with arthritis, (A1) 56

BUKHARI M, HERRICK A, MANNING J, MOORE T, JAYSON MIV. Capillary measurements by video capillary microscopy in patients with systemic sclerosis and primary Raynaud's phenomenon, (A1) 25

BULGEN DY. Minocycline-related lupus, (L) 398

BûLow PM See Nørregaard J

BUNCE M See Brown MA

BUNN CC, McMorrow M, BLACK CM. Patients with anti-RNA polymerase antibodies have diffuse scleroderma and exhibit a speckled pattern antinuclear antibody distinct from anti-Scl-70. (A1) 66

BLONCOMPAGNI A See Gattorno M

BURKE FD. Surgery to the wrist joint, 274

BURT J See Croft P

BUSH CH See Fiske DN

Bush rheumatology or rheumatologist gone bush?, 785

BUTCHER S See Hardwick LJ; Loughlin J

BXSB lupus prone mice, effect of dominant susceptibility genes on phenotype, (A1) 3

BYNG P See Campbell L

BYRNE J, JAWED S, JAMES MF, DACRE JE. Radiographic within-subject variability is low when measuring knee joint space with digital image analysis, (A1) 122

BYRNE P, RAJAN KT, EVANS WD. Does vasectomy predispose to osteoporosis?, (A2) 42

BYRON MA See Pimm TJ

C3 nephritic factor, hypocomplementaemia and type III mesangiocapillary glomerulonephritis progressing to systemic lupus erythematosus, (L) 90 C7 produced, where is it produced?, (A1) 155

C-SORIGUER FC See Tinahones FJ

Calcaneus, ultrasound and hip axis length in postmenopausal females: a twin study, (A1) 128

Calcification, subcutaneous and periarticular-a variant of limited scleroderma?, (A2) 53

Calcinosis, diltiazem induces remission in scleroderma, 576

Calcium phosphate basic salts and supraspinatus tendon degeneration, (A1) 41

Calin A. Bath Ankylosing Spondylitis Functional Index, (L) 793
Calin A, Jones SD, Kennedy LG. Juvenile onset ankylosing spondylitis more girls than we thought?, (A2) 34

CALIN A. The individual with ankylosing spondylitis: defining disease status and the impact of the illness, 663 CALIN A See also Brown MA; Garrett SL; Jones SD; Kabasakal Y; Kennedy

LG; Koh WH; Pande I CALLAHAN LF See Pincus T

CAMBRIDGE G See Worrall JG CAMP AV, PHILIPS DI, COLLIER SM, KIRK AP. The influence of knowledge, attitudes and personality characteristics on the outcome of acupuncture treatment, (A2) 51

CAMP AV See also Collier SM

CAMP V. The place of acupuncture in medicine today, (E) 404

CAMPBELL DA See Field M

CAMPBELL F, WEISSENSTEINER T, WARD ME, LANCHBURY J, PANAYI GS, KINGSLEY GH. Analysis of Chlamydia trachomatis specific synovial fluid derived T cell clones from a patient with recurent reactive arthritis, (A1)

CAMPBELL L, BYNG P, EGGER P, McLAREN M, CROFT P, COGGON D, COOPER C. Gender bias in patient selection for total hip arthroplasty, (A1) 34

CAMPBELL MK See Torgerson DJ

CAMPBELL RDC See Thomson W

Can changes in appendicular bone predict axial bone mineral density during treatment with hormone replacement therapy and cyclical etidronate?, (A2) 41

Can we differentiate between radiological progression over one year in patients with rheumatoid arthritis randomly allocated to im gold or hydroxychloroquine?, (A1) 107

Can we isolate the EEG activity related to on-going pain in the brain?, (A2) 13

CANDAL COUTO J See Veale DJ

CANTER RJ See Jones SD

CANVIN JMG, MACPHERSON BCM, STURROCK RD. High prevalence of apical pulmonary fibrosis in ankylosing spondylitis by high resolution computerized tomography, (A1) 117 CAPELL HA See McInnes IB; Morrison E

CAPELL HA. Clinical efficacy of sulphasalazine—a review, (S2) 35 Capillary measurements by video capillary microscopy in patients with systemic sclerosis and primary Raynaud's phenomenon, (A1) 25 CAPPER R See Jones SD

Capsaicin, topical, effects of upper motor neurons on the cutaneous response to, (A1) 53

Car drivers with rheumatoid arthritis, advice to, (L) 796

CARELESS DJ. Musculoskeletal complaints in patients with hyperlipoproteinaemia, (L) 393

CARKE S See Russell G

CARNEY SL See Billingham MEJ CARNEY T See Ross DA

Carpal tunnel syndrome

how does neurophysiological assessment influence the management and outcome of patients?, (L) 581

non-surgical management, (L) 890 use of a hand-pain diagram as a screening tool, (A1) 68

validation of a new method of assessing median nerve conduction at the wrist, (A1) 43

CARPENTER EH, PLANT MJ, FITZGERALD DA, SMITH DA, HASSELL AB, RYAN S, SHADFORTH MF, HOTHERSALL TE, MONTGOMERY B, FISHER J, CLARK S, DAWES PT. Management of oral complications of disease modifying drugs in rheumatoid arthritis, (A2) 20

CARPENTER EH See also Mayer B

CARR AJ, NANDRA D, HART DJ, DOYLE DV, THOMPSON PW, SPECTOR TD. The relationship between back pain, psychological status and quality of life in women in the general population: the Chingford study, (A1) 40 CARR AJ, THOMPSON PW. The impact of rheumatoid arthritis on body image,

(A1) 89

CARR AJ See also Hardwick LJ CARREIRA PE See Pablos JL

CARREÑO L See Abad MA; Alonso JC; González CM; González M; López-Longo FJ; Monteagudo I; Moreno FC CARRUTHERS DM, WATTS RA, SCOTT DGI. Pulse cyclophosphamide therapy

in systemic vasculitis, (A1) 21

CARRUTHERS DM See also Watts RA

Carson Dick, Dr W., obituary, 998

CARTER ME, CASHMAN PMM, KITNEY RI. Knee joint magnetic resonance images: enhancement by advanced 3D computer visualisation, (A1) 123 CARTHY D See Donn RP; Macgregor AJ

Cartilage

matrix, IL-6 and IL-4 role in IL-1-dependent degradation, 101

oligomeric matrix protein serum level relationship to disease progression in osteoarthritis of the knee joint, 306

quantitative imaging, (S1) 29

turnover markers, insulin-like growth factor-1: serum and synovial fluid levels and associations in osteoarthritis and rheumatoid arthritis, (A1) 55

CARTRON JP See Kerr JR

CARULLI MT, EPSTEIN O, BLACK CM. Small bowel bacterial overgrowth and omeprazole in patients with systemic sclerosis, (A1) 67

CARULLI MT See also Denton CP

of CREST syndrome with hypermobility: an unusual association, (A2)

of hepatitis C virus (HCV)-associated arthritis. Quantitative analysis of HCV RNA of the synovial fluid and the serum, (L) 691 report: dermatomyositis-scleroderma overlap in childhood, (A2) 54

CASEY ATH, CROCKARD A. In the rheumatoid patient: surgery to the cervical spine, 1078

CASEY EB See Donnelly S; McGonagle D

CASEY MC, WOOD C, HERBERT KE, FEEHALLY J, ISENBERG D, LUNEC J. Measurement of antibodies to reactive oxygen species damaged DNA as a simple marker of disease activity in SLE, (A1) 10

CASHMAN PMM See Carter ME CASSIMAN J-J See Vandevyer C

CATTENACH M See Cook A

CAULI A, YANNI G, PANAYI GS. IL-1 and IL-1ra production by monocyte/ macrophage in rheumatoid arthritis synovial membranes, (A1) 150

CAULI C See Pitzalis C

CAWLEY MID See Buckley MG; Peacock DJ; Wong WM

CAWSTON TE, CURRY V, RAMSEY S, CLARK IM, KYLE VA, ADEBAJO A, SILVERMAN B, DAYMOND T, HAZLEMAN BL. Levels of collagenase, tissue inhibitor metalloproteinases (TIMP), collagenase-TIMP compelxes and glycosaminoglycan in sequential samples of synovial fluid aspirated from patients with osteoarthritis, (A1) 138

CAWSTON TE. High free and latent collagenase activity in psoriatic arthritis synovial fluid, (L) 83

CAWSTON TE See also Adebajo AO; Bigg HF; Riley GP

CD4+ T-cell, synovial antigen-specific, prostanoid modulation of cytotoxic function in rheumatoid arthritis, 113

CD19 + peripheral blood lymphocytes, α2,6 sialyltransferase and β1, 4 galactosyltransferase activity are simultaneously reduced in rheumatoid arthritis, (A1) 62
CDP571

engineered anti-TNF α antibody, repeated dose in rheumatoid arthritis, (A1) 81

therapeutic effects of an engineered human anti-tumour necrosis factor alpha antibody in rheumatoid arthritis, 334

Cell adhesion molecules

an overview, (E) 1101

level reduction following sulphasalazine in patients with rheumatoid arthritis, (A1) 98

Cell type and mediator production analysis from tissues around loosening joint implants, 1127

Cellular sources of nitric oxide production in the joint, (A1) 44

CENP-B carboxyl terminal fragment, frequency of autoantibodies to a major epitope on in patients with autoimmune disease, 407

Centile curves of radiological damage: rheumatoid arthritis population, (A1) 106

Cerebral venous thrombosis and acquired protein S deficiency: an uncommon cause of headache in systemic lupus erythematosus, 572

Cervical spine

outcome of surgery to stabilise in rheumatoid arthritis—pain, neurological status and patient satisfaction, (A1) 92 surgery: in the rheumatoid patient, 1078

CHAIN BM See Binks MH

CHAKRAVARTY K See Jones PBB; Thomson W

CHALLACOMBE SJ See Kassimos DG

CHALTON D See Kalla AA

CHAMBERLAIN MA. Home exercises and out-patient hydrotherapy, (L) 992 CHAMBERLAIN MA See also Hillman M; Tennant A

CHAMBERS H, POPE D, HAY EM, SILMAN AJ. Weak association between selfreported dry eyes and dry mouth and objective tests of salivary and lacrimal function: a population study, (A1) 33

CHAMPOUX WC See Whyte J

CHAN EYT See Lau CS CHAN SY See Lau YL

CHANDLER C See Walker DJ: Wright D

CHANDLER P See Hogarth MB

Changes

in bone mineral density with bisphosphonate treatment of Paget's disease of bone, (A1) 130

in heel ultrasound bone measurements following strenuous exercise in army recruit training, (A1) 131

Changing autoantibody profiles with variable clinical manifestations in a patient with relapsing systemic systemic lupus erythematosus and polymyositis, 915

CHAPMAN PT, JAMAR F, KEELAN ETM, PETERS AM, HASKARD DO. Targeting endothelial activation in inflammation using 111In-labelled anti-E-selectin monoclonal antibody, (A1) 81

Characteristics of ulcers of the small bowel induced by non-steroidal antiinflammatory drugs in the rat: implications for clinical practice, 727

Characterization

of anti-endothelial cell membrane antibodies in systemic lupus erythematosus, (A1) 11

by capillary electrophoresis of synovial fluid chondroitin sulphates in arthritis, (A1) 54

CHARLES PJ See Laversuch CJ

CHATFIELD C See Pande I

Chemo-attractant effect of rheumatoid synovial fluid on lymphocytes, (A1) 57
CHEN ACN, JAYSON MIV, JONES AKP. Can we isolate the EEG activity related to on-going pain in the brain?, (A2) 13

CHESSER AMS See D'Cruz D

CHEUNG NT, SAKLATVALA J, DAWES PT. Multiple joint avascular necrosis: beware of tuberculosis and human immunodeficiency virus—a rare but important cause, 387

CHEUNG NT, TAYLOR D. DAWES PT, MATTEY D. In vitro effect of leflunomide on RA synovial fibroblasts proliferation and production of the matrix metallo-proteinase pro-MMP3, (A1) 60 CHEUNG NT, TAYLOR DJ, MATTEY DL, DAWES PT. Serological joint markers—a comparison of serum proMMP-3, proMMP-1 and disease activity in RA, (A1) 60

CHEUNG NT, WISEMAN M, ROLFE P, JONES PW, TAYLOR DJ, MATTEY DL, DAWES PT. Cyclical pressure and metalloproteinase release from human articular cartilage, (A2) 28

CHEUNG NT See also Averns HL; Russell G

CHICHEPORTICHE V See Lioté F

CHIKANZA I See Kassimos D

CHIKANZA IC, RYAN PJ, FOGELMAN I, GIBSON T. Single photon emission computerised tomography identifies pain sites in ankylosing spondylitis, (A1) 126

CHILD AH See Tobias JH

Childhood relationship analysis in adults with juvenile chronic arthritis, (A1)

CHIN GL See Segasothy M

CHINN R See Clunie G

Chlamydia trachomatis, analysis of specific synovial fluid derived T cell clones from a patient with recurent reactive arthritis, (A1) 152

Chloroquine retinopathy screening, (A1) 18

Chondrocalcinosis, familial, localisation of a gene in a UK kindred to chromosome 5P, (A2) 41

Chondrocyte integrins: increase in $\alpha 5$ and αV on stimulation with inflammatory synovial fluid, (A1) 57

Chondrocytes, human, signal transduction pathways activated by intermittent pressurisation, (A1) 58

Chondrodestructive and chondrosparing properties of non-steroidal antiinflammatory drugs in a model of destructive mono-articular arthritis, (A2) 15

Chondroitin sulphate

characterization by capillary electrophoresis of synovial fluid in arthritis, (A1) 54

and keratan sulphate epitopes, glycosaminoglycans and hyaluronan in normal knee synovial fluid, (A1) 55

CHOY E See Houssien DA

CHOY EHS, PANAYI GS, KINGSLEY GH. Therapeutic monoclonal antibodies, 707

CHOY EHS, SCHANTZ A, WOODY J, KINGSLEY GH, PANAYI GS. Human antichimaeric antibody response in rheumatoid arthritis patients after treatment with a chimaeric anti-CD4 monoclonal antibody, cM-T412, (A1) 101

CHOY EHS See also Houssein DA; Kassimos D; Kassimos DG; Rankin ECC CHRISTMAS S See Hodges E

Chronic abdominal sepsis and migratory polyarthritis, (L) 478

Chronic fatigue syndrome, hypothalamic-pituitary-adrenal function in patients with fibromyalgia, (A1) 69

Chronic nephrotoxicity in the treatment of arthritis, 162

Churg-Strauss syndrome with inflammatory orbital pseudotumour, (L) 485 CICUTTINI F See Spector TD

CIRUELO E See Gonzalez-Crespo MR

CISNAL A See Acasuso M

CLAAS FHJ See ten Wolde S CLARK D See Jobanputra P

CLARK IM See Cawston TE

CLARK S See Carpenter EH

CLARKE D See Borg AA

CLARKE S See Badcock LJ; Hassell AB; Russell G

CLAUDEPIERRE P, GUEGUEN A, LADJOUZE A, HAJJAJ-HASSOUNI N, SELLAMI S, AMOR B, DOUGADOS M. Predictive factors of severity of spondylarthropathy in North Africa, 1139

CLEGG DO See Tilley BC

CLIFTON A See Laversuch CJ

Clindamycin, combined intravenous with oral tetracycline in rheumatoid arthritis: an open evaluation, (A1) 96

Clinical competence, assessing: recognition of case descriptions of rheumatic diseases by general practitioners, 375

Clinical effects of a nitric oxide inhibitor on developing and established collagen induced arthritis in DBA/1 mice, (A1) 64

Clinical efficacy of sulphasalazine—a review, (S2) 35

Clinical patterns at presentation in 722 patients with early rheumatoid arthritis, examining the influences of age at onset and gender, (A2) 8 Clinical trial design for evaluating combination therapies, (S2) 92

Clonality of immunoglobulin production in rheumatoid synovium, (A1) 154
CLOSE DR, KIRBY JT, DAWES PT, STANWORTH DR, MATTEY DL. Inhibition
of antibody dependent cellular cytotoxicity (ADCC) in rheumatoid
arthritis by the non-immune complex of IgA-alpha-1-antitrypsin, (A2) 30

Clostridium perfringens antigens and toxoid, lymphocyte reactivity in patients with arthritis. (L) 1095

CLOT J See Combe B

CLUNIE G, GARBER S, PALEY M, WILKINSON I, CHINN R, EDWARDS J, ELL P, HALL-CRAGGS M. Synovial tissue volume quantification with gadolinium-DTPA enhanced magnetic resonance, (A1) 125 Clustering of sleep electroencephalographic patterns in patients with the fibromyalgia syndrome, 1151

Co-trimoxazole, fatal vasculitis following treatment, (L) 84

COAKLEY G. Toxic epidermal necrolysis, pancytopenia and adult respiratory syndrome, (L) 798

Cochrane collaboration and osteoarthritis, evidence-based medical practice, (E) 403

Coeliac disease, malabsorption caused by in patients who have scleroderma, 858

COGGON D See Campbell L; Dunne CA; Egger P

COLE J See Dunne CA

COLES CJ See Kirwan JR

Collagen

abnormal cross-linking in the cartilage of a diastrophic dysplasia patient, 512

type V (COL5A1), linkage of the gene that encodes the αl chain to type II Ehlers-Danlos syndrome, (A2) 40

Collagen induced arthritis, clinical effects of a nitric oxide inhibitor on developing and established disease in DBA/1 mice, (A1) 64

Collagenase

effect of growth factors on production from human tendon fibroblasts in culture, (A1) 138

high free and latent activity in psoriatic arthritis synovial fluid, (L) 83, (L) 84

tissue inhibitor of metalloproteinase (TIMP), collagenase-TIMP complexes and glycosaminoglycan levels in sequential samples of synovial fluid aspirated from patients with osteoarthritis, (A1) 138

COLLANTES E See Acasuso M; Tinahones FJ
COLLIER SM, PHILIPS DI, KIRK AP, CAMP AV. The effects of rheumatoid arthritis on the patient's spouse, (A2) 10

COLLIER SM See also Camp AV

COLLINS A See Elvins DM

COLLINS AJ, REID J, SOPER CJ, NOTARIANNI LJ. Characteristics of ulcers of the small bowel induced by non-steroidal anti-inflammatory drugs in the rat: implications for clinical practice, 727

Collins DA. Antigen specificity of ANCA in systemic vasculitis, (L) 394
Collins DA, Duke O. Combined intravenous clindamycin and oral tetracycline in rheumatoid arthritis: an open evaluation, (A1) 96

COLLINS DA See also Caulfield F; Johnson SM; Laversuch CJ COLLINS M See Barry R

COMBE B, EDNO L, LAFFORGUE P, BOLOGNA C, BERNARD J-C, ACQUAVIVA P, SANY J, BRESSOLLE F. Total and free methotrexate pharmacokinetics, with and without piroxicam, in rheumatoid arthritis patients, 421

COMBE B, ELIAOU J-F, DAURÉS J-P, MEYER O, CLOT J, SANY J. Prognostic factors in rheumatoid arthritis. Comparative study of two subsets of patients according to severity of articular damage, 529

Combination second-line or disease-modifying antirheumatic drug therapy in rheumatoid arthritis, (S2) 96

Combination of sulphasalazine and methotrexate in the management of rheumatoid arthritis—view based on personal clinical experience, (S2) 109

Combination therapy

clinical trial design for evaluating, (S2) 92

how and when should it be used? The role of an anchor drug, (S2) 100 in rheumatoid arthritis, (L) 580

Combined intravenous clindamycin and oral tetracycline in rheumatoid arthritis: an open evaluation, (A1) 96

Combining sulphasalazine and methotrexate in rheumatoid arthritis: early clinical impressions, (S2) 104

COMER M, SCOTT DL, DOYLE DV, HUSKISSON EC, HOPKINS A. Are slowacting anti-rheumatic drugs monitored too often? An audit of current clinical practice, 966

COMER M See also D'Cruz D

Community-based rheumatology clinic, shoulder pain, 440

Comparable co-expression of VCAM-1* and DAF* on synovial intimal cells and follicular dendritic cells explains synovial targeting in rheumatoid arthritis, (A2) 4

Comparative study of tenidap, a cytokine-modulating anti-rheumatic drug, and diclofenac in rheumatoid arthritis: a 24-week analysis of a 1-year clinical trial, 554

Comparison

of the accuracy of steroid placement with clinical outcome in patients with shoulder symptoms, (A1) 41

of anti-neutrophil cytoplasmic antibody positive with ANCA negative microscopic polyarteritis, (A1) 67

of the effectiveness and cost of treatment for vertebral fractures in women, 1167

of intravenous, intramuscular and oral pulse steroid therapies in the management of active rheumatoid arthritis, (A2) 19

of S-oxidation status in Asian and Caucasian rheumatoid arthritis patients, (A2) 19

of the validity and reliability of self-reported articular indices, 760 Complement, immune complexes and systemic lupus erythematosus, (A1) 82 Complement C4 polymorphism, and mannose-binding protein are associated with SLE in a Spanish population, (A1) 78

Complement C4A loci and HLA-DQA, relative contributions of in determining susceptibility to systemic lupus erythematosus, 221

determining susceptibility to systemic lupus erythematosus, 221 Complement component C4 gene expression regulation, (A1) 156 Complementary therapies, the baby and the bath water, (L) 479

Compound heterozygosity of HLA-DR4 and DR1 antigens in Asian Indians increases the risk of extra-articular features in rheumatoid arthritis, 41 Conceptualizing and defining outcome, (E) 899

Congenital heart block, asymptomatic mothers of babies with features of Sjögren's syndrome, (A1) 19

CONROY S See Faulds G

CONROY SE See Ehrenstein MR

Consequences of sustained activation of small sensory fibres on pain sensation and neurogenic inflammation, (A1) 53

Consultation, influence of physical, psychological and behavioural factors on for back pain, 156

Continuing medical education

for rheumatologists: the current picture, 172

in rheumatology, (E) 98

Controlled trial of intramuscular methylprednisolone against oral prednisolone in polymyalgia rheumatica: a 48 week follow up, (A1) 69

CONWAY SC See Baildam EM

COOK A, CATTENACH M, GALLAGHER G, FIELD M. Macrophage derived granulocyte macrophage colony stimulating factor may augment HLA-DR expression on inflammatory cells in temporal arteritis, (A1) 151 COOK NJ. Further possible cause of diarrhoea caused by oral gold, (L) 395

COOK NJ. Further possible cause of Cookson J See Gaffney K

COOPER AM See Williams AS

COOPER C See Campbell L; Dunne CA; Egger P; Peacock DJ

COOPER RG See Brammah TB; Davies EJ

CORKILL M See Rajapakse C

Correlation of scleroderma skin score to change in soluble ICAM-1 levels reflects endothelial activation, (A1) 28

CORRIGALL V See Robinson C CORRIGALL VM See Awad M

Corticosteroid osteoporosis, 8

Corticosteroid osteoporosis, 8

Corticosteroid-sparing therapeutic agent, low-dose methotrexate in the musculoskeletal manifestations of sarcoidosis, 642

Corticosteroids

differential regulation of LFA-1 and CD2 adhesion molecule gene expression in vitro, (A1) 146

high-dose pulse therapy effects on bone metabolism in rheumatoid arthritis, (A1) 128

inhibit lymphocyte adhesion to endothelial cells *in vitro*: confirmation using the SV40-transformed EC line LT4, (A1) 146

regulation of intracellular lipocortin-1 expression in murine leukocytes, (A1) 147

Corynebacterium pseudodiphtheriticum, discitis caused by following ear, nose and throat surgery, (L) 585

Cost effectiveness, treatment for vertebral fractures in women, 1167

COSTELLO P See Breathnach D COSTIGAN T See McGonagle D

COUGHLAN RJ See Flanagan FL

COUGHLAN RJ See Flanagan F COUMBE A See Gaffney K

Cox N See James D: Prouse P

Cox NL See James D; Pro

Coy SD See Wong WM

CRAFT A See Heslop P

CRAIG JP, TOMLINSON A, PATEL S, GRIERSON D, STURROCK RD. ABNORMAL TEAR EVAPORATION RATE AND OSMOLALITY IN RHEUMATOID ARTHRITIS. (A1) 91

Crawford F, Snaith M. Therapeutic ultrasound as a treatment of plantar heel pain, (A1) 42

CREAMER P, HUNT M, DIEPPE P. Pain mechanisms in osteoarthritis of the knee: effect of intra-articular anaesthetic, (A2) 45

CREE I See Veale DJ

CREST syndrome

with hypermobility: an unusual association, (A2) 54 and Wegener's granulomatosis, (L) 483, (L) 484

CROCKARD A See Casey ATH

CROFT P, BURT J, SCHOLLUM J, SILMAN A. Number of tender points and extent of reported pain, (A1) 38

CROFT P See also Campbell L; Hay EM; Lau E

CROFT PR, PAPAGEORGIOU A, FERRY S, THOMAS E, JAYSON MIV, SILMAN AJ. New episode incidence of low back pain: results from the south Manchester low back pain study, (A1) 71

CROFT PR See also Ferry S

Cronstein BN. The antirheumatic agents sulphasalazine and methotrexate share an anti-inflammatory mechanism, (S2) 30

CROSS S See Thomson W

Cross-sectional and longitudinal study of anti-ribonucleoprotein antibodies in lupus nephropathy, (A1) 8

CROSSMAN M See Lewis S

CROWLEY C See James IT

CROWSTON JG See Forbes JE

CRUWYS SC, GARRETT NE, KIDD BL. The consequences of sustained activation of small sensory fibres on pain sensation and neurogenic inflammation, (A1) 53

CRUWYS SC, GARRETT NE, PERKINS MN, KIDD BL. The neurogenic contribution to bradykinin-induced intra-articular plasma extravasation in the rat knee joint, (A1) 52

CRUWYS SC See also Jawed S

Cryoglobulinaemia, type 1 associated with a thymic tumour: successful treatment with plasma exchange, 285

CUADRADO MJ See Mujic F CURNOCK A See Seed MP

CURRAN MD See Kerr JR

CURRY V See Cawston TE

CURSON D See Pimm TJ

CURY SE, FERRAZ MB, DOS SANTOS JQ, MIRANDA C, REDKO C, VILAR MJP, ATRA E. The use of focus group interview in the evaluation of patients with ankylosing spondylitis, 150

Cutaneous leucocytoclastic vasculitis: a clinical and aetiological study, 942 Cutaneous lymphocyte antigen expression by peripheral blood and synovial fluid lymphocytes in psoriasis and psoriatic arthritis, (A1) 147

CUTOLO M, SULLI A, BARONE A, PICCIOTTO A, MANGRAVITI S, SERIOLO B, ACCARDO S. Macroamylasemia: a possible cause of unexplained hyperamylasemia in rheumatoid arthritis, 290

CUVELIER C See Mielants H

Cyclical pressure and metalloproteinase release from human articular cartilage, (A2) 28

Cyclophosphamide

intravenous in the treatment of systemic vasculitis: a five year follow-up study. (A1) 22

menstrual disorders in girls with systemic lupus erythematosus, 737 pulse therapy in systemic vasculitis, (A1) 21

Cyclosporin, effective treatment of anti Jo-1 antibody-positive polymyositis, (L) 1187

Cyclosporin A

in early rheumatoid arthritis patients: a 24 months chloroquine controlled follow up study, (A1) 99

in the treatment of rheumatoid associated pyoderma gangrenosum, (L) 182

Cytokine mRNA expression in the labial salivary gland from patients with primary Siögren's syndrome, 326

determination of various and type III procollagen aminopeptide levels in bronchoalveolar lavage fluid of the patients with pulmonary fibrosis: inverse correlation between type III procollagen aminopeptide and interferon- Γ in progressive patients, 31

effect of methotrexate alone or in combination with sulphasalazine on the production and circulating concentrations of cytokines and their antagonists, 747

inflammatory responses in juvenile chronic arthritis, 454

methotrexate action in rheumatoid arthritis: stimulation of cytokine inhibition and inhibition of chemokine production by peripheral blood mononuclear cells, 602

DABADGHAO S See Agrawal S

DACRE JE See Byrne J

DAI L, JONES H, ZHANG Z, EDMONDS SE, GAFFNEY K, BLAKE DR, WINYARD PG, MORRIS CJ. Oxidised LDL is present in rheumatoid synovial fluid. (A1) 49

DALTROY LH, ROBB-NICHOLSON C, IVERSEN MD, WRIGHT EA, LIANG MH. Effectiveness of minimally supervised home aerobic training in patients with systemic rheumatic disease, 1064

DALZELL N See Tobias JH

DALZIEL M, ALAVI A, AXFORD JS. α2,6 sialyltransferase and β1,4 galactosyltransferase activity are simultaneously reduced in CD19+ peripheral blood lymphocytes in rheumatoid arthritis, (A1) 62

DALZIEL M, BOND A, ALAVI A, HAY FC, AXFORD JS. α2,3 and α2,6 linked sialylation and Lewis X expression on serum IgG in rheumatoid arthritis, (A1) 63

Damage correlates with SLE disease duration in Afro-Caribbeans and Caucasians but not Asians, (A1) 17

DANNESKIOLD-SAMSØE B See Nørregaard J

DARKE C See Brown MA

DARLINGTON LG, JUMP A, MACRAE K, NOBBS BT, PANUSH RS, PETRIE A, RAMSEY DJC, RAMSEY NW, STREET P. Prospective, placebo-controlled study of single and double-blind food challenges in patients with rheumatoid arthritis after dietary manipulation therapy, (A1) 94 D'ARRIGO C See Veale DJ

DARTON K See Misra R

DASGUPTA B, BOWLES J. Scintigraphic localization of steroid injection site for plantar fasciitis, (A1) 42

DASGUPTA B, DOLAN L, PANAYI G, FERNANDES L. A controlled trial of intramuscular methylprednisolone against oral prednisolone in polymyalgia rheumatica: a 48 week follow up, (A1) 69

DASGUPTA B See also Dolan AL; Robinson C

DAURES J-P See Combe B

DAVID J See Murphy K; Rooney M; Vancheeswaran R

DAVIDSON F See Jobanputra P

DAVIES E, ORDI J, TEH LS, HILLARBY C, OLLIER WER. Mannose-binding protein and complement C4 polymorphism are associated with SLE in a Spanish population, (A1) 78

DAVIES E See also Ollier WER

DAVIES EJ, STEERS G, OLLIER WER, GRENNAN DM, COOPER RG, HAY EM, HILLARBY MC. Relative contributions of HLA-DQA and complement C4A loci in determining susceptibility to systemic lupus erythematosus, 221

DAVIES GJ See Mackworth-Young CG
DAVIES KA, BEYNON HLC, LOIZOU S; LAFFAN M, WALPORT MJ. Thrombosis in patients with anti-cardiolipin antibodies is not associated with activated protein-C resistance due to factor VLeiden mutation, (A1)

DAVIES KA. Complement, immune complexes and systemic lupus erythematosus, (A1) 82

DAVIES KA, NORSWORTHY P, WALPORT MJ. Anti-Clq antibodies are associated with hypo-complementaemia, defective opsonisation of gram-positive organisms, and death due to infection in patients with SLE, (A1) 10

DAVIES KA See also Athanassiou P; Hogarth MB

DAVIES P See James D

Davis J See Brown MA

Davis M See Dawes PT; Hassell A

Davis MJ See Borg AA; Deodhar AA

Davis P. Letter from abroad, 171 DAVIS P See also Prouse P

Davis SM See Symmons DPM

DAVIS W See Spitzer WO

DAWES P See Hassell A; Underwood MR

DAWES PT, DAVIS M, HOTHERSALL TE, SHADFORTH MF, HAY E, HASSELL A. Departmental outcome-forget the patients and staff-measure the

DAWES PT See also Ahmed S; Aung TS; Averns HL; Badcock LJ; Borg AA; Carpenter EH; Cheung NT; Close DR; Dziedzic K; Evans E; Fisher M; Hassell AB; Lyttle K; Mattey DL; Mayer B; Nixon NB; Plant MJ; Russell G; Salih AM

Dawes PW See Griffiths SM

Dawson JK. Advice to car drivers with rheumatoid arthritis, (L) 796
Dawson JK, Lynch MP. Urinary tract infection in patients with rheumatoid arthritis. A predictable problem?, (A1) 85

DAY A See Hilman S

DAYER J-M See Awad M

DAYMOND T See Cawston TE DAYMOND TJ See Scott DL

D'CRUZ D, CHESSER AMS, LIGHTOWLER C, COMER M, HURST MJ, BAKER LRI, RAINE AEG. Antineutrophil cytoplasmic antibody-positive crescentic glomerulonephritis associated with anti-thyroid drug treatment, 1090

D'CRUZ D, KESER G, KHAMASHTA MA, DIRESKENELI H, TARGOFF IN, MILLER F, HUGHES GRV. Antiendothelial cell antibodies in inflammatory myopathies: association with interstitial lung disease, (A1) 19

D'CRUZ D, TAUB NA, GIANOTTA L, MCALINDON TE, HUGHES GRV. Anti-Sm antibodies: a potential risk factor for lupus nephritis, (A1) 9

D'CRUZ D See also Froes L DE BENEDETTI F See Massa M

DE JONG-DE VOS VAN STEENWIJK C See Hogeweg JA

DE KLOET ER See Griep EN

DE LORD D. PUTHRASINGHAM S. MULLIGAN P. BROWN K. WALTERS G. PITT P. Broadband ultrasound characteristics in rheumatoid arthritis and diabetes, (A1) 123

DE POUVOURVILLE G. iatrogenic cost of non-steroidal anti-inflammatory drug therapy, (S1) 19

DE S. Pamidronate and uveitis, (L) 479

DE SAINT PIERRE V See SARAUX A

DE Vos M See Mielants H

DE VRIES RRP See ten Wolde S DE LA SAYETTE V. Acute necrotizing vasculitis (polyarteritis nodosa?)

confined to the nerve with spontaneous recovery, (L) 694 De Quervain's disease, mimicked by fluoroquinolone-induced tenosynovitis

of the wrist, (L) 583 Decreased DNA repair proficiency: a factor in susceptibility or a feature of disease in autoimmune conditions?, (A1) 4

Defective spontaneous and bacterial lipopolysaccharide-stimulated production of interleukin-1 receptor antagonist by polymorphonuclear neutrophils of patients with active systemic lupus erythematosus, 107

DEIGHTON C See Regan M

DEIGHTON CM See McDermott EM

DELMAS PD. Non-steroidal anti-inflammatory drugs and renal function, (S1)

DENTON CP. BICKERSTAFF MCM. SHIWEN X. CARULLI MT. HASKARD DO. DUBOIS RM, BLACK CM. Special circulating adhesion molecule levels reflect disease severity in systemic sclerosis, 1048

DENTON CP, SHIWEN X, WELSH KI, PEARSON JD, BLACK CM. Modulation of scleroderma and control fibroblast phenotype by co-culture with human endothelial cells, (A1) 66

DENTON CP See also Bartram SA; Black CM; Shiwen X

DEODHAR AA, BRABYN J, JONES PW, DAVIS MJ, WOOLF AD. Hand bone densitometry in early rheumatoid arthritis: a measure of progression despite clinical control, (A1) 108

Deoxycoformycin, a selective lymphocyte depleting drug, induces flare of rheumatoid arthritis, (A1) 102

Departmental outcome-forget the patients and staff-measure the notes, (A2) 49

Depo-methyl-prednisolone, intra-muscular versus intra-articular in the management of ambulatory patients with active rheumatoid arthritis, (A1) 102

DEQUECKER J See Boonen S

Dermatomyositis

irreversible visual loss, 1179

juvenile: results from a nation-wide study, 732

proton-NMR spectroscopy and the diagnosis, (A2) 27

Dermatomyositis-scleroderma overlap in childhood: case report, (A2) 54 Desensitization, low dose, does it affect the toxicity of sulphasalazine in the

treatment of rheumatoid arthritis?, (A1) 97 Design of a non constrained, non cemented, modular metacarpophalangeal prosthesis, (A2) 47

Destructive lymphadenopathy and T-lymphocyte activation in adult-onset Still's disease, 978

Detection

of antiribosomal P protein antibodies in patients with systemic sclerosis, 908

of nitrosyl haemoglobin in isolated synovial fluid and blood of rheumatoid patients, (A1) 46

Determinants of arthrogenic inhibition in knee osteoarthritis, (A1) 134 Determination

of the apparent synovial permeability in the knee joint of patients suffering from osteoarthritis and rheumatoid arthritis, 520

of urinary ribonucleosides in healthy individuals and those with rheumatic disease, (A1) 59

of various cytokines and type III procollagen aminopeptide levels in bronchoalveolar lavage fluid of the patients with pulmonary fibrosis: inverse correlation between type III procollagen aminopep tide and interferon- Γ in progressive patients, 31

Development of systemic lupus erythematosus

after chemotherapy and radiotherapy for malignant thymoma, 1175 features in patients with primary antiphospholipid syndrome, (A1) 16

DEVLIN J See James D; Prouse P

DEWALD B See Seitz M

DI GIOVINE F See Rooney M

Diabetes, broadband ultrasound characteristics, (A1) 123

Diagnostic associations with hypermobility

in new rheumatology referrals, (A1) 39

in rheumatology patients, 1157

Diagnostic criteria, sensitivity and specificity for Behçet's disease according to the latent class approach, 932

Diagnostic features of ankylosing spondylitis, (E) 301

Diarrhoea, further possible cause of caused by oral gold, (L) 395

Diastrophic dysplasia, abnormal collagen cross-linking in cartilage, 512 DICKIE A See Keir G

Diclofenac

double blind comparison of the endoscopic and clinical effects of tenoxicam and diclofenac in rheumatoid arthritis, (A1) 95 is a better analgesic than paracetamol in patients with osteoarthritis of

the hip, knee and lumbar spine, (A1) 137 and misoprostol fixed combination, efficacy and gastroduodenal safety of in the treatment of arthritis, (S1) 11

perforations, peptic ulceration and bleeding on NSAIDs. Two randomised controlled trials of nabumetone, diclofenac SR and piroxicam in the UK. (A1) 95

and tenidap, a cytokine-modulating anti-rheumatic drug, a comparative study in rheumatoid arthritis: a 24-week analysis of a 1-year clinical trial, 554

DIEPPE P See Creamer P; Rogers J; Shahid MS

DIEPPE PA See Dodd J; Hilman S; Lim K; Sharif M

Diet, elemental and subsequent food reintroduction effects on rheumatoid arthritis, 270

Dietary essential fatty acid supplements reduce red blood cell aggregation in healthy volunteers, (A1) 28

Dietary manipulation therapy, prospective, placebo-controlled study of single and double-blind food challenges in patients with rheumatoid arthritis, (A1) 94

Differences between adult and childhood onset scleroderma, (A1) 29

Different abdominal scintigraphy pattern in patients with ulcerative colitis, Crohn's disease and seronegative spondylarthropathies, 946

Differential diagnosis and management of hip pain in childhood, 78 Differential regulation by corticosteroids of LFA-1 and CD2 adhesion

molecule gene expression in vitro, (A1) 146

DUKMANS BAC. Folate supplementation and methotrexate, 1172

DIJKMANS BAC See also Kloppenburg M; ten Wolde S; van den Borne BEEM; van den Ende CHM

DILLON MJ See Nash MC

Diltiazem induces remission of calcinosis in scleroderma, 576

DINANT HJ See Kaandorp CJE; Lycklama à Nijeholt GJ

DIRESKENELI H See D'Cruz D

knee pain and psychological variables in a Nottingham population, (A1)

predictors one year after presentation with inflammatory polyarthritis, (A1) 86

relative importance of pain and X-ray damage to in long-standing RA, (A2) 12

in rheumatoid arthritis-it's relation to clinical and radiological courses over 3 years, (A2) 11

Discitis

acute calcific. A report of two cases with scintigraphy and SPECT, (A2)

caused by Corynebacterium pseudodiphtheriticum following ear, nose and throat surgery, (L) 585

Discordance between objective and subjective assessment of functional ability of patients with rheumatoid arthritis, 951

Disease activity

IL-2R levels in rheumatoid arthritis: poor correlation with clinical activity is due in part to disease duration, 1037

measurement of antibodies to reactive oxygen species damaged DNA as a simple marker in SLE, (A1) 10

Disease assessment, validated, Bath Ankylosing Spondylitis Radiology Index (BASRI), (A2) 37

Disease concordance, heterogeneity of disease phenotype in monozygotic twins concordant for rheumatoid arthritis, 216

Disease course, analysis of the discrepancy between the clinical and radiological course in rheumatoid arthritis, (A1) 105

Disease outcomes in rheumatology, (E) 704

Disease severity, rheumatoid arthritis patients not attending a specialist clinic have similar levels as clinic attenders, (A1) 88

Disease status, and impact of the illness, defining in ankylosing spondylitis, 663

Disease-modifying drugs

audit in rheumatoid arthritis, shared care between hospital and general practice, 673

combination second-line anti-rheumatic drug therapy in rheumatoid arthritis, (S2) 96

DISH, risk of vertebral fracture, (A1) 127

Disseminated penicillinase-producing Neisseria gonorrhoeae in a patient infected with human immunodeficiency virus, (L) 487

Diurnal variation

in 24 hour ambulatory recordings of patient activity is different in rheumatoid and osteoarthritis, (A2) 16

in soluble E-selectin, P-selectin and ICAM-1 levels in healthy volunteers, (A1) 87

DIXEY J See Doherty M; Elvins DM; Prouse P

DIXIE J See James D

DNA

damage in the rheumatoid peripheral blood mononuclear cell population assessment by single cell gel electrophoresis, (A1) 153

decreased repair proficiency: a factor in susceptibility or a feature of disease in autoimmune conditions?, (A1) 4

Do patients with SLE, SLE with Sjögren's syndrome and primary Sjögren's syndrome differ in damage or function?, (A2) 26

DOBBIE JW, HIND C, MEIJERS P, BODART C, TASIAUX N, PERRET J,

ANDERSON JD. Lamellar body secretion: ultrastructural analysis of an unexplored function of synoviocytes, 13

DODD J, HOLLY JMP, MEADOWS K, ELSON CJ, DIEPPE PA, SHARIF M. Insulin-like growth factor-1: serum and synovial fluid levels and associations with markers of cartilage turnover in osteoarthritis and rheumatoid arthritis, (A1) 55

Does low dose desensitization affect the toxicity of sulphasalazine in the treatment of rheumatoid arthritis?, (A1) 97

Does MCTD exist in India?, (A2) 26

Does mixed connective tissue disease exist in India?, 539

Does spiking of the tibial tubercles predict osteoarthritis of the knee?, (A1) 36 Does vasectomy predispose to osteoporosis?, (A2) 42

DOHERTY E. MULHERIN D. BRESNIHAN B. MCKANE R. TAGGART A. BELLA Analysis of childhood relationships in adults with juvenile chronic arthritis, (A1) 90

DOHERTY M, HUGHES A, McGIBBON D, WOODWARD E, DIXEY J, JONES R, HUNT A. Localisation of a gene for familial chondrocalcinosis in a UK kindred to chromosome 5P, (A2) 41

DOHERTY M See also Belcher C; Hodges E; Jones AC; Ledingham J; Lewis S; O'REILLY SC; O'Reilly SC; Regan M; Yaqub R

DOLAN AL, BAARN K, GRAHAME R, SPECTOR TD. Assessment of bone in Ehlers Danlos syndrome by ultrasound and densitometry, (A1) 130 DOLAN AL, DASGUPTA B, TODD P, FERNANDES L, PANAYI GS, MONIZ C.

Serial changes in bone mineral density in polymyalgia rheumatica, (A1) 74

DOLAN AL, KASSIMOS D, GIBSON T, KINGSLEY GH. Diltiazem induces

remission of calcinosis in scleroderma, 576 Dolan AL, Ryan PJ, Arden N, Stratton R, Hamann W, Wedley J, FOGELMAN I, GIBSON T. SPECT scanning can predict successful facet joint injection, (A1) 71

DOLAN AL, SWINBURN HM, PRICE TR. Ferritin is an acute phase protein in longstanding and recent onset rheumatoid arthritis, (A1) 83

DOLAN L See Dasgupta B

Dominant human leukocyte antigen DR4-Dw15 restricted mycobacterial 65 kDa antigen-specific T-cell immunity in Chinese patients with rheumatoid arthritis, 498

Dominant susceptibility gene effects on phenotype in BXSB lupus prone mice. (A1) 3

DONALD F, ESDAILE JM, KIMOFF J, FITZCHARLES M-A. Low prevalence of fibromyalgia syndrome in patients attending a sleep disorders clinic, (A1)

DONN RP, FARHAN A, THOMSON W, OLLIER WER, HOLT PJL, KAY RA. T-cell receptor variable gene segment TCRBV6S1 null allele frequencies in UK juvenile chronic arthritis, (A1) 144

DONN RP, THOMSON W, PEPPER L, CARTHY D, FARHAN A, RYDER C, SOUTHWOOD T, HOLT L, OLLIER W. Antinuclear antibodies in early onset pauciarticular juvenile chronic arthritis (JCA) are associated with HLA-DQB1*0603: a possible JVA-associated human leucocyte antigen haplotype, 461

DONNELLY S, JAWED S, HART DJ, DOYLE DJ, SPECTOR TD. Does spiking of

the tibial tubercles predict osteoarthritis of the knee?, (A1) 36 DONNELLY S, JAWED S, MEJIA A, DOYLE DV. Effect of syndesmophyte formation on lumbar spine bone density in patients with ankylosing spondylitis, (A1) 117

DONNELLY S, LONG A, HALL, MURPHY A, CASEY EB, KELLEHER D. Role of protein kinase C in cellular activation for signalling through CD44 and LFA-1, (A1) 156

DONNELLY S See also McGonagle D

DONOGHUE DO See Mulherin D

DORAN M, FENELON H, SANT SM, BREATNACH ES. High resolution computed tomography of the lung in asymptomatic SLE patients with normal chest radiographs, (A1) 14

DOS SANTOS JQ See Cury SE

Double blind comparison of the endoscopic and clinical effects of tenoxicam and diclofenac in rheumatoid arthritis, (A1) 95

Double blind placebo controlled trial of minocycline in patients with active rheumatoid arthritis, (A2) 16

Double-blind placebo-controlled study of ranitidine for the prevention of recurrent peptic ulcer disease in rheumatoid arthritis patients on nonsteroidal anti-inflammatory drugs, (A1) 94

Dougados M. Diagnostic features of ankylosing spondylitis, (E) 301

Dougados M See also Claudepierre P

DOYLE DV See Carr AJ; Comer M; Donnelly S; Hart DJ; Lloyd ME; Spector

DR1 and HLA-DR4, compound heterozygosity in Asian Indians increases the risk of extra-articular features in rheumatoid arthritis, 41 DREWES AM, GADE J, NIELSEN KD, BJERREGÅRD K, TAAGHOLT SJ,

SVENDSEN L. Clustering of sleep electroencephalographic patterns in patients with the fibromyalgia syndrome, 1151

DREWES AM, NIELSEN KD, TAAGHOLT SJ, BJERREGARD K, SVENDSEN L, GADE J. Sleep intensity in fibromyalgia: focus on the microstructure of the sleep process, 629

Dry eyes

assessment of primary Sjögren's syndrome, (A2) 25

and dry mouth, weak association between self-reported and objective tests of salivary and lacrimal function: a population study, (A1) 33

DRYLL A See Lioté F DU BOIS RM See Bartram SA; Shiwen X

Dual energy X-ray absorptiometry of the knee joint, (A1) 122

DUANE P See Raja KB

DUBOIS RM See Denton CP

DUDLEY HART F, SAVAGE MO. Oswald Arthur Savage, obituary, 591

DUGGLEBY S See Egger P DUKE O See Caulfield F

DUKE OL See Kaminski ER

DUNNE C See Jones HW

DUNNE CA. Generalized pruritis in the rheumatoid patient-consider all possible causes, (L) 295

DUNNE CA, THOMPSON PW, COLE J, DUNNING J, MARTYN CN, COGGON D, COOPER C. Validation of a new method of assessing median nerve conduction at the wrist in carpal tunnel syndrome, (A1) 43

DUNNING J See Dunne CA

DUNPHY J See Elvins DM

Duodenal iron uptake and serum and mucosal iron protein levels, with special reference to rheumatoid arthritis, 1041

DYER PA See Howarth S

DZIEDZIC K, HAY EM, JACKSON SE, JONES PW, ROLFE P, DAWES PT. The body chart: a meaningful assessment of pain compared to visual analogue scales in ankylosing spondylitis, (A2) 36

DZIEDZIC K See also Averns HL

cE-selectin, cICAM-1 and cICAM-3 in patients with rheumatoid vasculitis, (A1) 87

E-selectin

P-selectin and ICAM-1 diurnal variation in levels in healthy volunteers. (A1) 87

relationship with ICAM-1 in serum, synovial fluid and synovial membrane in inflammatory arthritis, (A1) 148

Early postmenopausal bone loss and polymorphisms of the vitamin D receptor, (A1) 79

EARNSHAW WC See Whyte J

EASTMOND CJ, GARTON M, ROBINS S, RIDDOCH S. The effects of alcoholic beverages on urate metabolism in gout sufferers, (A1) 120, 756

EBRAHIM E See Thompson PW EDMONDS J See Lassere M

EDMONDS SE See Dai L

EDNO L See Combe B

EDTA bis(ethyl phenylalaninate): a novel transition metal-ion chelating hydroxyl radical scavenger with a potential anti-inflammatory role, (A1)

Education in rheumatology: what do general practitioners want?, (A2) 49

EDWARDS J See Clunie G

EDWARDS JCW, MALIK J, WILKINSON LS. The synovial lining-associated epitope 67: tissue distribution and identification by Western blotting,

EDWARDS JCW, WILKINSON LS. Comparable co-expression of VCAM-1* and DAF* on synovial intimal cells and follicular dendritic cells explains synovial targeting in rheumatoid arthritis, (A2) 4

EDWARDS NL See Fiske DN

EEG activity, can we isolate it related to on-going pain in the brain?, (A2) 13 Effect

of 2nd line antirheumatic therapy on pyridinium crosslink excretion in patients with established RA, (A1) 99

of dominant susceptibility genes on phenotype in BXSB lupus prone mice, (A1) 3 of gold salts on monocytes and their pattern of cytokine secretion, (A1)

of growth factors on collagenase and TIMP production from human

tendon fibroblasts in culture, (A1) 138 of high-dose corticosteroid pulse therapy on bone metabolism in rheumatoid arthritis, (A1) 128

of hypermobility syndrome on knee joint proprioception, 121

of liposomally conjugated methotrexate upon mediator release from human peripheral blood monocytes, 241

of methotrexate alone or in combination with sulphasalazine on the production and circulating concentrations of cytokines and their antagonists, 747

of minocycline on acute phase proteins and serum interleukin-6 levels in rheumatoid arthritis patients, (A1) 96

of smoking on outcome of ankylosing spondylitis, (A2) 35

of sulphasalazine on neutrophil superoxide protection in RA, (A2) 14

of syndesmophyte formation on lumbar spine bone density in patients with ankylosing spondylitis, (A1) 117

of systemic lupus erythematosus and long-term steroid therapy on bone mass in pre-menopausal women, 742

of thiazide use on bone density: the Chingford study, (A1) 129

Effective treatment of anti Jo-1 antibody-positive polymyositis with cyclosporin, (L) 1187

Effectiveness of minimally supervised home aerobic training in patients with systemic rheumatic disease, 1064

of alcoholic beverages on urate metabolism in gout sufferers, (A1) 120.

of elemental diet and subsequent food reintroduction on rheumatoid arthritis, 270

of joint washout and steroid injection compared with either joint washout or steroid injection alone in rheumatoid knee effusion, 771

of reactive oxygen species on the ability of IgG to interact with Clq, (A1)

of rheumatoid arthritis on the patient's spouse, (A2) 10

of upper motor neurons on the cutaneous response to topical capsaicin, (A1) 53

Efficacy

and gastroduodenal safety of a fixed combination of diclofenac and misoprostol in the treatment of arthritis, (S1) 11 of methotrexate in rheumatoid arthritis, (S2) 43

EGGER P, FRITH S, DUGGLEBY S, FALL C, COOPER C. Obesity, occupational activity and osteoarthritis of the spine, (A1) 35

EGGER P See also Campbell L; Peacock DJ

Ehlers-Danlos syndrome

assessment of bone by ultrasound and densitometry, (A1) 130

type II, linkage of the gene that encodes the al chain of type V collagen (COL5A1) to, (A2) 40

EHRENSTEIN MR, CONROY SE, HEATH J, LATCHMAN DS, ISENBERG DA. The occurrence, nature and distribution of flares in a cohort of patients with systemic lupus erythematosus: a rheumatological view, 257

EHRENSTEIN MR. Combination therapy in rheumatoid arthritis, (L) 580
EHRENSTEIN MR, KATZ DR, GRIFFITHS M, WINKLER TH, KALDEN JR,
ISENBERG DA. Human IgG anti-DNA antibodies deposit in kidneys and induce proteinuria in SCID mice, (A1) 65 EHRENSTEIN MR See also Rankin ECC

Eight year outcome of patients presenting with early non-erosive rheumatoid arthritis. (A2) 8

EISMAN JA See Keen RW

EL MANZALAWI See Troughton PR

Elderly, rheumatoid arthritis: is peripheral oedema at onset a good prognostic feature?, (A2) 10

Elemental diet and subsequent food reintroduction effects on rheumatoid arthritis, 270

Elevated serum nitrate concentrations in active spondylarthropathies, 616

ELEWAUT D See Mielants H ELIAOU J-F See Combe B

ELL P See Clunie G

ELLIOTT JRM See Kerr JR

ELMORE JG See Spitzer WO

ELSON CJ See Dodd J; Hilman S; Perry MJ; Sharif M

ELVINS DM, MITRA D, DIXEY J, DUNPHY J, COLLINS A, RING EFJ, O'NEILL TW, BHALLA AK. Bone density and bone turnover in early AS, (A2) 34 EMERY P. Surgery to the shoulder-shoulder replacement, 653 EMERY P. Therapeutic approaches for early rheumatoid arthritis. How early?

How aggressive?, (S2) 87

EMERY P See also Gordon C; Gough A; Griffiths B; Griffiths HR; James D; McRorie ER; Prouse P; Reece R; Scott DL

Employment status of patients in rheumatology clinics, (L) 693

Endochondreal ossification, identification of genes expressed specifically during, (A1) 77

Endogenous opiates: are they responsible for the defective hypothalamic response to inflammatory stimuli in patients with rheumatoid arthritis?, (A1) 84

Endothelial activation, targeting in inflammation using 111In-labelled anti-Eselectin monoclonal antibody, (A1) 81

Endothelial cells

conditioned medium promotes contraction of collagen gels by scleroderma lung fibroblasts, (A1) 31

factors promoting apoptosis in vitro, (A1) 32

human model of ischaemia-reperfusion injury, (A1) 32

synovial microvascular in culture express active xanthine oxidase, (A1)

synovial microvascular exhibit enhanced shear rate dependent leucocyte adhesion in vitro, (A1) 51

Endothelial leucocyte adhesion molecule, levels are not increased and levels of circulating intercellular adhesion molecule-1 and-3 are increased in patients with rheumatoid arthritis, 311

Endothelium-derived haemostatic factors and the antiphospholipid syndrome, 201

Endothelium-derived relaxing factor/nitric oxide increased production in patients with systemic vasculitis, (A1) 23

ENGELHARDT G, THOMAE K. Meloxicam: a preferential inhibitor of COX-2,

ENGLAND A See Martin JC

Enhanced macrophage degradation of low density lipoprotein from synovial fluid of rheumatoid arthritis patients, (A1) 49

Enthesitis, antibodies to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of ankylosing spondylitis patients with/without iritis and enthesitis, 418

Environmental, life-style and medical factors influencing peak bone mass in women, 620

Environmental change and disease, 1105

Environmental lupus syndromes, (E) 597

Eosinophilia myalgia syndrome, continuing occurrence in Canada, 246

Epidemiology

and prevention of non-steroidal anti-inflammatory drug effects in the gastrointestinal tract, (S1) 5

of vertebral osteophytosis, (A1) 70

Epidural infiltration, steroid myopathy induced by triamcinolone injection,

Epithelial cell activation and interferon-gamma in Sjögren's syndrome, 226 Epitope 67, synovial lining-associated: tissue distribution and identification by Western blotting, (A1) 149

EPSTEIN O See Carulli MT

ERLACHER L, WINTERSBERGER W, MENSCHIK M, BENKE-STUDNICKA A, MACHOLD K, STANEK G, SÖLTZ-SZÖTS J, SMOLEN J, GRANINGER W Reactive arthritis: urogenital swab culture is the only useful diagnostic method for the detection of the arthritogenic infection in extraarticularly asymptomatic patients with undifferentiated oligoarthritis,

ERNST E. Complementary therapies, the baby and the bath water, (L) 479

ESCALONA M See López-Longo FJ

Escherichia coli

Klebsiella pneumoniae and Proteus mirabilis antibodies in the sera of ankylosing spondylitis patients with/without iritis and enthesitis,

Klebsiella pneumoniae and Proteus mirabilis antibodies in the sera of patients with axial and peripheral form of ankylosing spondylitis, 413

ESDAILE JM See Donald F: Hudson N

ESPINOSA G See Pons F

Essential fatty acid supplements reduce red blood cell aggregation in healthy volunteers, (A1) 28

ESSEX T See Walker DJ; Wright D

Estimates of knee pain and disability in the community with differing knee pain criteria, (A1) 36

Estimating fracture risk gradients from case-control studies, (A2) 43 ETHERINGTON J, BUCHANAN N, LIMA F, GREENSTAIN B, BROAD I, KERSLAKE

S, HUGHES GRV. Prolactin levels in lupus pregnancy do not differ significantly from those in controls. There is no correlation between disease activity and prolactin levels, (A1) 13

ETHERINGTON J, HARRIS P, NANDRA D, HART DJ, SPECTOR TD. Bone mineral density and exercise: a comparison of female ex-elite athletes and aged matched controls, (A1) 72

ETHERINGTON J, KEELING J, BRAMLEY R, MCCURDIE I, BAAN K, GARNETT R, SPECTOR TD. Changes in heel ultrasound bone measurements following strenuous exercise in army recruit training, (A1) 131

ETHERINGTON J, MOOTOO R, MATHEWS JA. An audit of the outpatient use of disease modifying and immunosuppressive anti-rheumatic drugs, (A1)

ETHERINGTON J See also Spector TD

Etidronate, cyclical and hormone replacement therapy, can changes in appendicular bone predict axial bone mineral density during treatment, (A2) 41

EUSTACE J, BROPHY D, GIBNEY R, BRESNIHAN B, FITZGERALD O. Comparison of the accuracy of steroid placement with clinical outcome in patients with shoulder symptoms, (A1) 41

Evaluation of microwave thermography of the hand and wrist in rheumatoid arthritis, (A2) 13

EVANS C See Thompson PW

Evans D See Vaghmaria A

EVANS E, HASSELL A, MATTEY D, GASTON H, DAWES PT. Salmonella-a forgotten infection in seronegative oligoarthritis?, (A2) 32

EVANS MD, PERRETT D, LUNEC J, NICHOL FE, HERBERT KE. DNA damage in the rheumatoid peripheral blood mononuclear cell population assessment by single cell gel electrophoresis, (A1) 153

EVANS MD, PERRETT D, LUNEC J, NICHOL FE, HERBERT KE. Determination of urinary ribonucleosides in healthy individuals and those with rheumatic disease, (A1) 59

EVANS MD See also Hill CM

EVANS WD See Byrne P

Evidence

for genetic anticipation in nodal osteoarthritis, (A1) 133

of omeprazole-induced small bowel bacterial overgrowth in patients with scleroderma, 976

for a possible gene pool difference between ethnic groups?-a controlled study of British and Algerian subjects, (A1) 143

Evidence-based medical practice: the Cochrane collaboration and osteoarthritis, (E) 403

and bone mineral density: a comparison of female ex-elite athletes and aged matched controls, (A1) 72

randomized controlled trial of four different programs in patients with rheumatoid arthritis, (A1) 93

strenuous, changes in heel ultrasound bone measurements in army recruit training, (A1) 131

EXLEY A See McRorie ER

Extracellular matrix environment and the human peripheral blood dendritic cells, in vitro maturation, (A1) 145

FITZCHARLES M-A See Donald F

Facet joint injection, SPECT scanning can predict successful, (A1) 71

Factor VIIA, increase of activated and haemostatic molecular markers in juvenile chronic arthritis, 466

Factor VIII von Willbrand factor antigen, assessing systemic lupus erythematosus activity using, (A1) 8

Factors

associated with the development of vasculitis in rheumatoid arthritis: results of a case-control study, (A1) 86

promoting endothelial cell apoptosis in vitro, (A1) 32 FAIN O. Aortitis: a new manifestation of primary antiphospholipid syndrome, (L) 686

FAIRBURN K See Jones HW: Nazhat NB

FAITH JM See Spitzer WO

FAKHOURY H, HILLARBY C, WEISS JB, JAYSON MIV. Neutral matrix metalloproteinases in scleroderma, (A1) 29

FALGA C See Marsal S

FALGA-TIRADO C. Lhermitte's sign in Behçet's disease, (L) 184

FALL C See Egger P; Peacock DJ

False positive IgA and IgG rheumatoid factor isotypes in infectious diseases, (A1) 109

Familial risk for rheumatoid arthritis is low in first degree relatives of probands with inflammatory arthritis ascertained from a populationbased register, (A1) 80

FARHAN A See Donn RP

FARHAN AJ, OLLIER WE, KAY RA. The TCRBV6S1*3 null allele may confer additional RA risk in HLA shared epitope positive individual, (A1) 141 FAROOQI A. Chronic abdominal sepsis and migratory polyarthritis, (L) 478 FARR M, BACON PA. How and when should combination therapy be used?

The role of an anchor drug, (S2) 100 FARR M See also Bacon PA; Popert AJ; Waterhouse L

FARRALL M See Brown MA

FARRELL AM See Hodges E

FAS, soluble is increased in patients with SLE, (A1) 5

Fatal vasculitis following treatment with co-trimoxazole, (L) 84

Fatigue relationship in ankylosing spondylitis to disease activity, sleep and other factors, (A2) 36

FATIMA Z See Hameed K

FAULDS G, CONROY S, MADAIO M, ISENBERG D, LATCHMAN D. Increased levels of antibodies to heat shock proteins with increasing age in MRL/ MP-LPR/LPR mice, 610

FAWTHROP F See Belcher C; Yaqub R

FAY M See Bailly S

FC/II receptor-mediated arrest under flow conditions is enhanced by Substance P, (A1) 52

FEAR J See Tennant A

Feasibility study of homeopathic remedy in rheumatoid arthritis, (A2) 17

FEEHALLY J See Casey MC

FEELY J See Wylie G

FELSENBERG D See Kiss Cs

Felty's syndrome, absence of exogenous retroviruses, (L) 185

FENELON H See Doran M

FERNANDES L See Dasgupta B; Dolan AL

FERRAZ MB See Cury SE

FERRELL WR See Hall MG

FERRI C See Berrettini S

Ferritin is an acute phase protein in longstanding and recent onset rheumatoid arthritis, (A1) 83

FERRY S, PRITCHARD T, CROFT PR, KEENAN J, SILMAN AJ. Use of a handpain diagram as a screening tool for carpal tunnel syndrome, (A1) 68 FERRY S See also Croft PR

Fetal calf serum in growth medium obscures the detection of early IgG antocardiolipin-antibody secreting clones, (A1) 12

Fibroblasts, human synovial, TGF-β1 induces myofibroblastic differentiation in in vitro: modulation by basic FGF, (A2) 29

Fibromyalgia

gynaecological symptoms in, (L) 888

hypothalamic-pituitary-adrenal function in patients with chronic fatigue syndrome, (A1) 69

muscle strength, voluntary activation and cross-sectional muscle area, 925

sleep intensity in: focus on the microstructure of the sleep process, 629 studies of sleep in; techniques, clinical significance and future directions, (E) 897

Fibromyalgia syndrome

diagnostic associations with hypermobility in rheumatology patients,

low prevalence in patients attending a sleep disorders clinic, (A1) 38 sleep electroencephalographic pattern clustering, 1151

FIELD M, NELSON S, CAMPBELL DA, GALLAGHER G. TNF NCO-1 RFLP is not a predictive risk factor in rheumatoid arthritis, (A1) 140

FIELD M See also Cook A; McInnes IB

FISHER A See Thompson PW

FISHER AD See Wright D

FISHER J See Badcock LJ; Carpenter EH; Hassell AB; Russell G

FISHER M, RALPHS A, SHAPLEY M, TUCKER R, DAWES PT, HAY EM. Outpost rheumatology: the general practitioners' perspective, (A2) 50

FISHMAN D. Successful management of reactive haemophagocytic syndrome in systemic-onset juvenile chronic arthritis, (L) 888

FISKE DN, BUSH CH, EDWARDS NL. Axial skeletal changes in paralysed patients may mimic ankylosing spondylitis, 174

FITZCHARLES M-A See Hudson N

FITZGERALD DA See Carpenter EH

FITZGERALD O See Breathnach D; Eustace J; Martin K; Mulherin D

Five-year follow-up of a controlled trial of an arthritis education programme,

FLANAGAN FL, SANT S, COUGHLAN RJ, O'CONNELL D. Symptomatic enlarged iliopsoas bursae in the presence of a normal pain hip radiograph, 365

Flares, occurrence, nature and distribution in a cohort of patients with systemic lupus erythematosus: a rheumatological view, 257

FLOWER RJ See Perretti M

Flucloxacillin and methotrexate, lack of interaction between in patients with rheumatoid arthritis, (A1) 100

Fluorophore assisted electrophoresis, oligosaccharide changes in rheumatoid arthritis determined by, (A1) 63

Fluoroquinolone-induced tenosynovitis of the wrist mimicking de Quervain's disease, (L) 583

Focus group interview use in the evaluation of patients with ankylosing spondylitis, 150

Foetally inherited paternal HLA does not explain post partum susceptibility to rheumatoid arthritis in the mother, (A1) 142

FOGELMAN I See Chikanza IC: Dolan AL

Folate supplementation and methotrexate, 1172

Folic acid and methotrexate, an audit of for rheumatoid arthritis. Experience from a teaching centre, 971

FONT J See Pons F

Food challenges, prospective, placebo-controlled study of single and doubleblind food challenges in patients with rheumatoid arthritis after dietary manipulation therapy, (A1) 94

FOOT V See Speed CA

Foot and ankle in rheumatoid arthritis, 443

FORBES J See Whitefield L

FORBES JE, CROWSTON JG, HOWARD M, BRAZIER DJ. Screening for chloroquine retinopathy, (A1) 18

FORDHAM JN See Speed CA

FOSTER H, GRIFFITHS I. A regional audit of adult onset Still's disease, (A2) 22

FOSTER H See also Heslop P FOULKES R See Ward PS

FOWLER PD See Hassell AB

Fracture risk gradient estimation from case-control studies, (A2) 43

FRAGA A See Medina F

FRANCIS RM, ANDERSON FH, TORGERSON DJ. A comparison of the effectiveness and cost of treatment for vertebral fractures in women, 1167 FRANCOMBE J, KINGSNORTH AN, TUNN E. Panniculitis, arthritis and

pancreatitis, 680 FRASER SM See Hamilton JD

FRASER WD See MacDonald AG

FREARS E See Zhang Z

Frequency of autoantibodies to a major epitope on the carboxyl terminal fragment of CENP-B in patients with autoimmune disease, 407

FRIMAN C See Julkunen H

FRITH S See Egger P

FROES L, D'CRUZ D, LLOYD ME, KHAMASHTA MA, HUGHES GRV. Intravenous cyclophosphamide in the treatment of systemic vasculitis: a five year follow-up study, (A1) 22

FRÖLICH JC See Stichtenoth DO

FRYER A See Ollier WER

FUENTES M See Medina F FUJIKAWA Y See Shingu M

FUJIMOTO M, SATO S, TAKEHARA K, NOJIMA Y, SOMA Y, KIKUCHI K, IHN H, IGARASHI A, TAMAKI K. Detection of antiribosomal P protein antibodies

in patients with systemic sclerosis, 908 Fuлимото M See also Ihn H

Functional ability, discordance between objective and subjective assessment of patients with rheumatoid arthritis, 951

FURST DE. Practical clinical pharmacology and drug interactions of low-dose methotrexate therapy in rheumatoid arthritis, (S2) 20

Further possible cause of diarrhoea caused by oral gold, (L) 395

GADE J See Drewes AM

Gadolinium-DTPA enhanced magnetic resonance imaging in rheumatoid knee synovitis: clinical, laboratory and histological correlation, (A1) 75

GAFFNEY K, COUMBE A, BLADES S, BLAKE D, COOKSON J. Gadolinium-DTPA enhanced magnetic resonance imaging in rheumatoid knee synovitis: clinical, laboratory and histological correlation, (A1) 75

GAFFNEY K, WILLIAMS R, JOLLIFFE V, BLAKE D. Intra-articular pressure changes in normal and rheumatoid peripheral joints, (A1) 46

GAFFNEY K See also Dai L; Naughton D; Winrow VR GAIKWAD P See Troughton PR

Galactosyltransferase isoenzyme changes occur in rheumatoid arthritis, (A1) 61

GALLAGHER G See Cook A; Field M

GALLATI H See Seitz M

GALLIVAN S See James D; Prouse P

GARBER S See Clunie G

GARCIA-CONSUEGRA J See Gonzalez-Crespo MR

GARDNER CR See Seed MP

GARDNER-MEDWIN JMM, MACDONALD IA. Reduced vasodilatation to acetylcholine in the finger microcirculation in primary Raynaud's phenomenon, (A1) 25

GARNETT R See Etherington J GARRETT NE See Cruwys SC

GARRETT SL, KENNEDY LG, CALIN A. The nature of surgical intervention in patients with ankylosing spondylitis, (A1) 115

GARRETT SL See also Jones SD; Kabasakal Y; Kennedy LG; Koh WH

GARTON M See EASTMOND CJ; Eastmond CJ

GARTON MJ, REID DM. Estimating fracture risk gradients from case-control studies, (A2) 43

GARTON MJ See also Keir G

GASTON H See Evans E

GASTON JSH. Involvement of the gut in the pathogenesis of inflammatory synovitis, (E) 801

Gastric ulcer patients, limited association of Helicobacter pylori with rheumatic disease, (L) 1188

Gastroduodenal safety and efficacy of a fixed combination of diclofenac and misoprostol in the treatment of arthritis, (S1) 11

Gastrointestinal tract, epidemiology and prevention of non-steroidal antiinflammatory drug effects in the gastrointestinal tract, (S1)

GATTORNO M, BUONCOMPAGNI A, MOLINARI AC, BARBANO GC, MORREALE G, STALLA F, PICCO P, MORI PG, PISTOIA V. Antiphospholipid antibodies in paediatric systemic lupus erythematosus, juvenile chronic arthritis and overlap syndromes: SLE patients with both lupus anticoagulant and high-titre anticardiolipin antibodies are at risk for clinical manifestations related to the antiphospholipid syndrome, 873 GEENEN R, JACOBS JWG, GODAERT G, KRAAIMAAT FW, BRONS MR, VAN

DER HEIDE A, BIJLSMA JWJ. Stability of health status measurement in rheumatoid arthritis, 1162

Gemfibrozil, spondylitis-like and symmetric polyarthralgia associated with

therapy, (L) 692

Gender bias in patient selection for total hip arthroplasty, (A1) 34
GENDI NST, GIBSON K, WORDSWORTH BP. The relationship between prolonged joint symptoms, hypocomplementaemia and HLA type in parvovirus arthritis, (A1) 111

GENDI NST. Lupus-like syndrome in patients treated for acne, (L) 584

Gene pool difference, evidence for between ethnic groups?—a controlled study of British and Algerian subjects, (A1) 143

General practice, musculoskeletal problems in: a prospective survey of consultations and management, (A2) 50

General practitioners

perspective of outpost rheumatology, (A2) 50 recognition of case descriptions of rheumatic diseases, assessing clinical competence, 375

what do they want in rheumatology education?, (A2) 49

Generalized pruritis in the rheumatoid patient-consider all possible causes, (L) 295

Genetic anticipation

evidence for in nodal osteoarthritis, (A1) 133 in familial rheumatoid arthritis, (A2) 6

Genetic susceptibility to rheumatoid arthritis and peptide transporter genes (TAP) polymorphisms, 207

GEORGANAS K See Athanassiou P

GERBER NJ See Beyeler C; Seitz M

GERRITS MI See Lems WF

GEUSENS P See Vandevyer C GIANOTTA L See D'Cruz D

Giant cell arteritis

aetiopathogenesis, 960

presenting with ischaemic skin lesions of the neck, 1182

GIBNEY R See Eustace J

GIBSON K See Brown MA; Gendi NST

GIBSON T, HAMEED K. Rheumatic diseases and rheumatoid arthritis amongst Pakistanis in London and Karachi, (A1) 33

GIBSON T See also Chikanza IC; Dolan AL; Gordon P; Hameed K

GIFFORD L. See Herrick AL.

GILLESPIE KM See Kiely PDW

GILLESPIE S See Kave SA

GILLET P. Fluoroquinolone-induced tenosynovitis of the wrist mimicking de Quervain's disease, (L) 583

GIRI TK See Uppal SS

GLANT TT See Brennan FR

GLENNAS A, KVIEN TK, MELBY K, REFVEM OK, ANDRUP O, KARSTENSEN B, THOEN JE. Acute sarcoid arthritis: occurrence, seasonal onset, clinical features and outcome, 45

GLENNON P, SHEERAN T, PRICE T. Irritable bone syndrome-a new concept in somatisation?, (A2) 52

Global tests for multiple outcomes in rheumatoid arthritis trials, (A1) 89 Glomerulonephritis

antineutrophil cytoplasmic antibody-positive crescentic associated with anti-thyroid drug treatment, 1090

type III mesangiocapillary, hypocomplementaemia and C3 nephritic factor progressing to systemic lupus erythematosus, (L) 90

Glutathione S-transferase (GSTM1 and GSTT1) polymorphisms influence on the incidence and severity of rheumatoid arthritis, (A2) 18

Glyceryl trinitrate, sustained-release transdermal patches as a treatment for primary and secondary Raynaud's phenomenon, 636

Glycolipid antibodies, synovial fluid in rheumatoid arthritis-synovial neuropathy or synovial damage, (A2) 4

Glycosaminoglycan, collagenase, tissue inhibitor of metalloproteinase (TIMP) and collagenase-TIMP complex levels in sequential samples of synovial fluid aspirated from patients with osteoarthritis, (A1) 138

GODAERT G See Geenen R GOEI THE HS See van den Borne BEEM

can we differentiate between radiological progression over one year in patients with rheumatoid arthritis randomly allocated to im gold or hydroxychloroquine?, (A1) 107

injections, neurological side effects in two patients with rheumatoid arthritis, 989

oral, further possible cause of diarrhoea caused by, (L) 395 parenteral

HLA phenotypes and therapeutic response in rheumatoid arthritis, (A1) 97

oral steroids as bridge therapy in rheumatoid arthritis patients starting with parenteral gold. A randomized double-blind placebo-controlled trial, 347

salt effects on monocytes and their pattern of cytokine secretion, (A1)

Gold-induced remission, and human leucocyte antigen phenotypes in patients with rheumatoid arthritis, 343

GOLIKA P See Vaghmaria A

GOMEZ-REINO JJ See Gonzalez-Crespo MR; Pablos JL

GONZÁLEZ CM, LÓPEZ-LONGO FJ, MONTEAGUDO I, GRAU R, MORENO AC, GONZÁLEZ M, ABAD MA, MÍNGUEZ M, ARIZA A, ALONSO JM, CARREÑO L. Association of anti-Ro/SS-A antibodies with progressive multifocal leukencephalopathy in HIV infected patients, (A1) 112

GONZÁLEZ CM See also Abad MA; Alonso JC; González M; López-Longo FJ; Monteagudo I; Moreno FC

GONZÁLEZ M, LÓPEZ-LONGO FJ, LÓPEZ-GOMEZ JM, JOFRE R, GONZÁLEZ CM, MORENO AC, GRAU R, ARIZA A, ALONSO JM, CARREÑO L. Crosssectional and longitudinal study of anti-ribonucleoprotein antibodies in lupus nephropathy, (A1) 8

GONZÁLEZ M See also Abad MA; Alonso JC; González CM; López-Longo FJ: Moreno FC

GONZÁLEZ-AMARO R See Laffón A

GONZALEZ-CRESPO MR, BLANCO FJ, RAMOS A, CIRUELO E, MATEO I, LOPEZ PINO MA, GOMEZ-REINO JJ. Magnetic resonance imaging of the brain in systemic lupus erythematosus, 1055

GONZALEZ-CRESPO MR, GOMEZ-REINO JJ, MERINO R, CIRUELO E, GOMEZ-REINO FJ, MULEY R, GARCIA-CONSUEGRA J, PINILLOS V, RODRIGUEZ-VALVERDE V. Menstrual disorders in girls with systemic lupus erythematosus treated with cyclophosphamide, 737

GONZALEZ-RUIZ A See Tinahones FJ

GOODACRE JA See Ross DA

Gooi HC See Bradley SM

GORDON C, MALIK R, STOLL T, SITUNAYAKE RD, EMERY P, BACON PA, ISENBERG DA. Damage correlates with SLE disease duration in Afro-Caribbeans and Caucasians but not Asians, (A1) 17

GORDON C See also Griffiths B

GORDON P, JONES H, GIBSON T. The relative importance of pain and X-ray damage to disability in long-standing RA, (A2) 12

GOSLING J See Shahid MS

GOUDIE RB See KARIM SN

GOUGEROT-POCADILO MA See Bailly S

GOUGH A. ANDREWS D. BACON PA. EMERY P. Evidence of omentazoleinduced small bowel bacterial overgrowth in patients with scleroderma, 976

GOUGH A See also James D; Prouse P

GOULDING NJ See Perretti M

Gout

effects of alcoholic beverages on urate metabolism, (A1) 120, 756 management, (L) 482

management in patients with renal insufficiency and a gastrointestinal bleed-an answer to Dr Dunne's letter, (L) 481

in patients attending the rheumatology unit of Lomé Hospital, 843 quantitative study of the phagocytosis of urate crystals in the synovial fluid of asymptomatic joints, 724

serum urate acute, (A2) 40

GOWLAND G See Willis D

GRABOWSKI PS, RALSTON SH. Cellular sources of nitric oxide production in the joint, (A1) 44

GRAHAM S See Jobanputra P

GRAHAME R, PYERITZ RE. The Marfan syndrome: joint and skin manifestations are prevalent and correlated, 126

GRAHAME R See also Dolan AL GRANFORS K See Mäki-Ikola O

GRANINGER W See Erlacher L; Koeller M

GRANT ME See Hillarby MC

GRANT SFA, REID DM, RALSTON SH. A polymorphism in an Sp1 consensus binding motif in the first intron of collagen I alpha 1 is related to postmenopausal osteoporosis, (A1) 78

Granulocyte macrophage colony stimulating factor, macrophage derived may augment HLA-DR expression on inflammatory cells in temporal arteritis, (A1) 151

GRAU R See González CM; González M; López-Longo FJ; Monteagudo I; Moreno FC

GREAVES M See Hill MB

GREEN H See Rafferty JA

GREENSTAIN B See Etherington J

GREER M See Veale DJ

GRENNAN DM See Davies EJ; Herrick AL

GRIEP EN, BOERSMA JW, LENTJES E, VERCOULEN JHMM, SWANINK CMA, VAN DER MEER JWM, PRINS APA, VAN DER KORST JK, DE KLOET ER. Hypothalamic-pituitary-adrenal function in patients with fibromyalgia and chronic fatigue syndrome, (A1) 69

GRIERSON D See Craig JP

GRIFFIN K See Herrick AL; Rafferty JA

GRIFFITH SM See Lyttle K

GRIFFITHS B, KLOVRZA L, GORDON C, WARING RH, EMERY P, SITUNYAKE RD. Comparison of S-oxidation status in Asian and Caucasian rheumatoid arthritis patients, (A2) 19

GRIFFITHS B, KLOVRZA L, GORDON C, WARING RH, SITUNAYAKE RD, EMERY P. Variation in sulphoxidation capacity in systemic lupus erythematosus patients and healthy controls of different ethnic origins, (A1) 6

GRIFFITHS G See Pickavance L

GRIFFITHS HR, EMERY P, LUNEC J. The effects of reactive oxygen species on the ability of IgG to interact with Clq, (A1) 45

GRIFFITHS I See Foster H

GRIFFITHS ID, STURROCK RD. Dr W. Carson Dick, obituary, 998

GRIFFITHS M See Ehrenstein MR

GRIFFITHS SM, HASSELL AB, JONES PW, HOTHERSALL TE, SHADFORTH MF. Audit of patients taking sulphasalazine and penicillamine individually, sequentially and concurrently in inflammatory arthritis, (A2) 17

GROOTVELD M See Naughton D

GROSSIN M See Kaye O GROSSMAN A See Kassimos D

Growth factor effects on collagenase and TIMP production from human tendon fibroblasts in culture, (A1) 138

Growth of scientific interest in rheumatology, (L) 796

GUAÑABENS N See Peris P; Pons F

GUBLER FM See Lycklama à Nijeholt GJ

GUEGUEN A See Claudepierre P

Gut

influence of sulphasalazine on the evolution of inflammation in the spondylarthropathies, (A1) 115

involvement in the pathogenesis of inflammatory synovitis, (E) 801

GUTHRIE BT See Marshall KW

GUY M See Brammah TB

Gynaecological symptoms in fibromyalgia, (L) 888

GYSELBRECHT G See Mielants H

HAAGSMA CJ See Barrera P; Van Gestel AM

HAAGSMA CJ, VAN RIEL PLCM, VAN DE PUTTE LBA. Combining sulphasalazine and methotrexate in rheumatoid arthritis: early clinical impressions, (S2) 104

HAGGERTY JL See Spitzer WO

HAJJAJ-HASSOUNI N See Claudepierre P

HÄKKINEN A, HANNONEN P, HÄKKINEN K. Muscle strength in healthy people and in patients suffering from recent-onset inflammatory arthritis, 355

HALL C See Roberts R HALL M See Bajocchi GL

HALL MG, FERRELL WR, STURROCK RD, HAMBLYN DL, BAXENDALE RH. The effect of hypermobility syndrome on knee joint proprioception, 121

HALL N See Donnelly S HALL ND See Whyte J

HALL NF See Hamilton JD

HALL-CRAGGS M See Clunie G

HAMA N See SATOH M

HAMBLYN DL See Hall MG

HAMEED K, GIBSON T, KADIR M, SULTANA S, FATIMA Z, SYED A. The prevalence of rheumatoid arthritis in affluent and poor urban commu-

nities of Pakistan, 252
HAMEED K, GIBSON T. Subcutaneous and periarticular calcification—a variant of limited scleroderma?, (A2) 53

HAMEED K See also Gibson T

HAMILTON JD, HALL NF, KYLE PM, LEE WR, FRASER SM. Orbital pseudotumour: a new ocular entity in RA?, (A2) 55 HAMMAN W See Dolan AL

HAMMOUDEH M. Spondylitis-like and symmetric polyarthralgia associated with gemfibrozil therapy, (L) 692

HAN S-H See Hsieh S-C

Hand bone densitometry in early rheumatoid arthritis: a measure of progression despite clinical control, (A1) 108

Hand-pain diagram as a screening tool for carpal tunnel syndrome, (A1) 68 HANSEN BL, JUNKER P. Giant cell arteritis presenting with ischaemic skin lesions of the neck, 1182

HARBUZ MS, PERVEEN-GILL Z, LIGHTMAN SL, JESSOP DS. A protective role for testosterone in adjuvant-induced arthritis, 1117 HARDWICK HJ, SLAVOTINEK A, WALSH S, BUTCHER S, WEEKS D, CARR AJ,

WORDSWORTH BP. Linkage analysis in multiple epiphyseal dysplasia, (A1) 144

HARDWICK LJ See Loughlin J

HARLEY SL See Abbot SE

HARMER I See Mackworth-Young CG

HARRALL RL See Bigg HF; Riley GP

HARRIS CM See Bruce IN HARRIS P See Etherington J; Keen RW

HARRIS PA See Spector TD

HARRISON BJ, SYMMONS DPM, BANKHEAD C, BARRETT EM, SCOTT DGI, SILMAN AJ. Predictors of resolution in inflammatory polyarthritis, (A1)

HARRISON BJ, SYMMONS DPM, BARRETT EM, SCOTT DGI, SILMAN AJ. Survival on sulphasalazine-results from the Norfolk Arthritis Register,

HARRISON BJ See also Thomson W

HART DJ, NANDRA D, McALINDON TE, WHEELER MJ, DOYLE DV, SPECTOR TD. The relationship of IGF-1 and osteoarthritis in women aged 45-64 in the general population: the Chingford study, (A1) 137

HART DJ See also Carr AJ; Donnelly S; Etherington J; Spector TD HARVEY GR, WHYTE J, MCHUGH NJ. Two distinct sites on the RNA

polymerase-II large subunit are recognized by scleroderma autoantibo-dies but one site is unique to the phosphorylated (IIO) form, (A1) 31 Has polyarteritis nodosa been classified out of existence?, (A2) 25

HASKARD DO See Chapman PT; Denton CP

HASLOCK I, MACFARLANE D, SPEED C. Intra-articular and soft tissue injections: a survey of current practice, 449

HASLOCK I See also Speed CA

HASSAN WU, KEANEY NP, HOLLAND CD, KELLY CA. Association of HLA-DR4, protease inhibitor phenotypes and keratoconjunctivitis sicca with pulmonary abnormalities in rheumatoid arthritis, 37

HASSELL A, PLANT M, DAVIS M, MATTEY D, DAWES P, OLLIER W, JONES P, ALLDERSEA J. STRANGE R. Influence of glutathione S-transferase (GSTM1 and GSTT1) polymorphisms on the incidence and severity of rheumatoid arthritis, (A2) 18

HASSELL A See also Dawes PT; Evans E
HASSELL AB, PLANT MJ, CLARKE S, FISHER J, JONES PW, SAKLATVALA J,
FOWLER PD, SHADFORTH MF, DAWES PT. Small joint synovitis in rheumatoid arthritis: should it be assessed separately?, 51

HASSELL AB See also Carpenter EH; Griffiths SM; Russell G

HASSELL AH See Mayer B

HASSON N See Vancheeswaran R

HAWKINS CP See Salih AM

HAY E See Dawes PT; Mayer B

HAY EM, VAGHMARIA A, SHELLEY M, CROFT P. Rheumatology education: what do general practitioners want?, (A2) 49

HAY EM See also Chambers H; Davies EJ; Dziedzic K; Fisher M

HAY FC See Alavi A; Dalziel M; Martin K

HAYEM G See Bailly S

HAYNES J See Hewlett SE

HAZENBERG MP See Melief MJ

HAZES JMW See ten Wolde S; van den Ende CHM; Vliet Vlieland TPM; Voskuyl AE

HAZLEMAN B See Barry R

HAZLEMAN BL See Adebajo AO; Cawston TE; Kavanagh R; Riley GP; Vecchio P

Health economics as an aspect of health outcome: basic principles and application in rheumatoid arthritis, 774

Health status

and joint damage divergence in rheumatoid arthritis, (A1) 107 measurement stability in rheumatoid arthritis, 1162

Heat shock proteins, increased levels of antibodies with increasing age in MRL/MP-LPR/LPR mice, 610

Heat-shock protein gene HSP70-2 polymorphism in systemic lupus erythcmatosus, 721

HEATH J See Ehrenstein MR

Heel ultrasound

bone measurement changes following strenuous exercise in army recruit training, (A1) 131

measurement, identification of an at risk group for osteoporosis in general practice, (A1) 37

HEINEGARD D See Sharif M

HELDERS PJM See Hogeweg JA

Helicobacter pylori in gastric ulcer patients, limited association with rheumatic disease, (L) 1188

HELIOVAARA M, AHO K, REUNANEN A, KNEKT P, AROMAA A. Parity and risk of rheumatoid arthritis in Finnish women, 625

HELLIWELL PS. Employment status of patients in rheumatology clinics, (L)

HELLIWELL PS, O'HARA M. Shared care between hospital and general practice: an audit of disease-modifying drug monitoring in rheumatoid arthritis, 673

HELLIWELL PS. Osteoarthritis and Paget's disease, (A2) 46, 1061 HELLIWELL PS. Should tests for proteinuria be included in the monitoring schedule of sulphasalazine?, 790

HELSLOOT J, VIRGO S, McGUIGAN L, STURGESS A. Antineutrophil

cytoplasmic antibodies in inflammatory arthritis-potential for misdiagnosis, 820

HELVE T See Paimela L

HEMERYCK L See McGonagle D

HENDERSON D See Hill C

HENDERSON KJ, HICKS R, WORRALL JG. Shape changes in synovial fibroblasts from RA patients, (A1) 145

Hepatitis B-associated polyarteritis nodosa, is interferon alpha a therapy for?,

Hepatitis C

viral infection and rheumatic diseases in HIV infected patients, (A1) 111 virus (HCV)-associated arthritis. Quantitative analysis of HCV RNA of the synovial fluid and the serum, (L) 691

HERBERT KE See Casey MC; Evans MD; Hill CM

Hereditary complement factor I deficiency, molecular basis, (A1) 155

Heritability of hip axis length and ultrasound of the calcaneus in postmenopausal females: a twin study, (A1) 128

HERMANS J See ten Wolde S; van den Ende CHM

HERNANDEZ MV See Peris P

HERNÁNDEZ RODRIGUEZ I. Primary pyomyositis. Two more cases with atypical presentation in diabetic hosts, (L) 482

HERRICK A See Blann AD; Bukhari M

HERRICK AL, BARLOW JD, BOWDEN A, WILLIAMS N, HOBSON AR, IRVING M, JAYSON MIV. Abnormal anal function in patients with systemic sclerosis, (A1) 26

HERRICK AL, GIFFORD L, GRIFFIN K, GRENNAN DM, AARONS L. Lack of interaction between methotrexate and flucloxacillin in patients with

rheumatoid arthritis, (A1) 100 HERRICK AL. Neurological involvement in systemic slerosis, (E) 1007

HERRICK AL See also Anderson ME; Rafferty JA

HESLOP P, FOSTER H, CRAFT A, WALKER D. NUMACT: a novel method of monitoring physical activity in JCA, (A2) 23

HESLOP P See also Walker DJ; Wright D

HESS EV. Environmental lupus syndromes, (E) 597

Heterogenity of disease phenotype in monozygotic twins concordant for rheumatoid arthritis, 215

HEWLETT SE, HAYNES J, SHEPSTONE L, KIRWAN JR. Rheumatoid arthritis patients cannot accurately report signs of inflammatory activity, 547

HEYMANN R See Bosi Ferraz M

HEYSE SP See Tilley BC

HICKLING P See Bacon PA

HICKS CA See Billingham MEJ

HICKS R See Henderson KJ

High diagnostic value of antiperinuclear factor-IgG: prospective analysis of 1004 sera diluted 1:100 and more, (L) 87

High free and latent collagenase activity in psoriatic arthritis synovial fluid, (L) 83, (L) 84

High frequency of parvovirus B19 in serum samples submitted for a rheumatoid factor test, (A2) 33

High prevalence of apical pulmonary fibrosis in ankylosing spondylitis by high resolution computerized tomography, (A1) 117

High resolution computed tomography of the lung in asymptomatic SLE patients with normal chest radiographs, (A1) 14

HILL C, PILE K, HENDERSON D, KIRKHAM B. Neurological side effects in two patients receiving gold injections for rheumatoid arthritis, 989

HILL CM, HERBERT KE, EVANS MD, LUNEC J. Neutrophil-derived tumour necrosis factor-α: a potential mechanism for promotion of inflammation in procainamide-induced lupus, (A1) 3

HILL MB, PHIPPS JL, MILFORD-WARD A, GREAVES M, HUGHES P. Characterisation of anti-endothelial cell membrane antibodies in systemic lupus erythematosus, (A1) 11

HILLARBY C See Davies E; Fakhoury H

HILLARBY MC, THOMAS JT, BRADY G, WALLIS GA, GRANT ME, BOOT-HANDFORD RP. The identification of genes expressed specifically during endochondreal ossification, (A1) 77

HILLARBY MC See also Davies EJ: Poulton K

HILLMAN M, WRIGHT A, RAJARATNAM G, TENNANT A, CHAMBERLAIN MA. Prevalence of low back pain and patterns of consultation, (A2) 38

HILLMAN M See also Tennant A

HILMAN S, DAY A, ELSON CJ, DIEPPE PA, SHARIF M. Bone-specific alkaline phosphatase as a marker of bone turnover in osteoarthritis and rheumatoid arthritis, (A1) 56

HIND C See Dobbie JW

arthroplasty, total, gender bias in patient selection, (A1) 34

axis length and ultrasound of the calcaneus in postmenopausal females: a twin study, (A1) 128

involvement, predictive factors of severity of spondylarthropathy in North Africa, 1139

involvement in juvenile chronic arthritis evaluation with contrastenhanced MR imaging, (A1) 124

osteoarthritis, acetabular dysplasia does not cause in a Chinese population, (A1) 34

pain, differential diagnosis and management in childhood, 78

radiograph, normal plain, symptomatic enlarged iliopsoas bursae, 365 replacement, total, outcome in ankylosing spondylitis: a thirteen year follow-up study, (A1) 114

transient regional osteoporosis of the hip. (L) 296

HIRAKATA M See SATOH M

Historical overview of non-steroidal anti-inflammatory drugs, (S1) 2

History of the use of sulphasalazine in rheumatology, (S2) 2

associations of different patterns of psoriatic arthritis, (A1) 143

associations of inflammatory polyarthritis developing post-immunisation, (A1) 76

class II associations of ankylosing spondylitis, (A1) 142

component of the genetic contribution to ankylosing spondylitis, (A1) 76 foetally inherited paternal does not explain post partum susceptibility to rheumatoid arthritis in the mother, (A1) 142 maternal antigens and antibodies to SS-A/Ro and SS-B/La. Comparison

with systemic lupus erythematosus and primary Sjögren's syndrome,

phenotypes and therapeutic response to parenteral gold in rheumatoid arthritis, (A1) 97

and rheumatoid arthritis linkage and penetrance in extended pedigrees, (A1) 139

HLA-A2, is remitting seronegative symmetrical synovitis with pitting oedema (RS3PE syndrome) associated with?, (L) 889

HLA-B27-related unclassifiable seronegative spondylarthropathy in females and its comparison with the profile in males, 137

HLA-DQA and complement C4A loci, relative contributions of in determining susceptibility to systemic lupus erythematosus, 221 HLA-DR4

and DR1 antigens, compound heterozygosity in Asian Indians increases the risk of extra-articular features in rheumatoid arthritis, 41 protease inhibitor phenotypes and keratoconjunctivitis sicca association

with pulmonary abnormalities in rheumatoid arthritis, 37 HLA-DRB1*0401 and DRB1*0404, polymyalgia association, (A1) 139

HML-1 expression by peripheral blood and synovial fluid lymphocytes in rheumatoid and psoriatic arthritis, (A1) 148

Ho M See Veale DJ

HOBBS R See Peacock DJ HOBSON AR See Herrick AL

HOCHBERG M See Lim K

HODGES E, QUIN C, FARRELL AM, CHRISTMAS S, SEWELL HF, DOHERTY M, POWELL RJ, SMITH JL. Arthropathy, leucopenia and recurrent infection associated with a TcR $\Gamma\delta$ population, 978

HOFER HO See Stoll T

HOGARTH MB, CHANDLER P, MORLEY BJ, SLINGSBY JH, NORSWORTHY P, SIMPSON E, DAVIES KA, WALPORT MJ. The effect of dominant susceptibility genes on phenotype in BXSB lupus prone mice, (A1) 3

HOGEWEG JA, KUIS W, HUYGEN ACJ, DE JONG-DE VOS VAN STEENWIJK C, BERNARDS ATM, OOSTENDORP RAB, HELDERS PJM. The pain threshold in juvenile chronic arthritis, 61

HOUER MA See Melief MJ

HOLLAND CD See Hassan WU

HOLLINGWORTH P. Differential diagnosis and management of hip pain in childhood, 78 HOLLMAN AS See Neilly JB

HOLLY JMP See Dodd J

HOLT L See Brammah TB; Donn RP

HOLT PJL See Baildam EM; Donn RP; Kay RA

Home exercises and out-patient hydrotherapy, (L) 991, (L) 992

Homeopathic remedy, feasibility study in rheumatoid arthritis, (A2) 17

HOMMA M See Satoh M

Homozygosity for glutathione-transferase GSTM1 and GSTT1 null alleles may confer risk for anti-Ro antibodies in SLE, (A1) 5

HOPKINS A See Comer M

HORIE K See Shingu M

Hormone replacement therapy and cyclical etidronate, can changes in appendicular bone predict axial bone mineral density during treatment, (A2) 41

HORSLEY SD See Wright D

HORWITZ RI See Spitzer WO

HOSKING DJ See Patel S; Yeap SS

Hospital notes, departmental outcome-forget the patients and staffmeasure the notes, (A2) 49

HOTHERSALL TE See Carpenter EH; Dawes PT; Griffiths SM; Mayer B

HOUSSEIN D See Lassere M

HOUSSEIN DA, CHOY EHS, BERRY H, SCOTT DL. Health status and joint damage divergence in rheumatoid arthritis, (A1) 107

HOUSSIEN DA, CHOY E, BERRY H, SCOTT DL. What quality of life assessments should be used in RA?, (A2) 11

HOUSSIEN DA See also Scott DL

How does neurophysiological assessment influence the management and outcome of patients with carpal tunnel syndrome?, (L) 581

How and when should combination therapy be used? The role of an anchor drug. (S2) 100

HOWARD M See Forbes JE

HOWARD Y See Silman AJ

HOWARTH S, RIDWAY JC, STEWART I, DYER PA, OLLIER WER. Polymyalgia is associated with both HLA-DRB1*0401 and DRB1*0404, (A1) 139

HOWELL WM See Wong WM Howie AJ See Lugmani RA

Hpertension, benign intracranial, skin hyper-reactivity in patients as an early manifestation of Behçet's disease, (L) 184

HSIEH S-C, TSAI C-Y, SUN K-H, TSAI Y-Y, TSAI S-T, HAN S-H, YU H-S, YU C-L. Defective spontaneous and bacterial lipopolysaccharide-stimulated production of interleukin-1 receptor antagonist by polymorphonuclear neutrophils of patients with active systemic lupus erythematosus, 107

HUBER E See Stoll T HUDSON N, STARR M, ESDAILE JM, FITZCHARLES M-A. Diagnostic

associations with hypermobility in new rheumatology referrals, (A1) 39 HUDSON N, STARR MR, ESDAILE JM, FITZCHARLES M-A. Diagnostic associations with hypermobility in rheumatology patients, 1157

HUGHES A See Doherty M

HUGHES GRV See Buchanan NMM; D'Cruz D; Etherington J; Froes L; Lima F; Mujic F; Rosman A

HUGHES P See Hill MB

HUGHES R See Humphries J

HUGHES RA See Stevens RJ

Hughes syndrome: antiphospholipid antibodies move closer to thrombosis in 1994. (E) 493

HUGUET M See Pons F

Human alpha-1-proteinase inhibitor, oxidative inactivation by simultaneous generation of superoxide and nitric oxide, (A1) 48

Human anti-chimaeric antibody response in rheumatoid arthritis patients after treatment with a chimaeric anti-CD4 monoclonal antibody, cM-T412, (A1) 101 Human IgG anti-DNA antibodies deposit in kidneys and induce proteinuria

in SCID mice, (A1) 65

Human immunodeficiency virus

antibodies to and to intermediate filaments, (A1) 112 disseminated penicillinase-producing Neisseria gonorrhoeae, (L) 487

and tuberculosis, multiple joint avascular necrosis-a rare but important cause, 387

Human immunodeficiency virus infection

Salmonella pyomyositis in, 568 arthritis associated with in Zambia, (A2) 32 association of anti-Ro/SS-A antibodies with progressive multifocal leukencephalopathy, (A1) 112

hepatitis C viral infection and rheumatic diseases, (A1) 111 staphylococcal arthritis in, (L) 397

Human leucocyte antigen

DR4-Dw15, dominant restricted mycobacterial 65 kDa antigen-specific T-cell immunity in Chinese patients with rheumatoid arthritis, 498

JVA-associated haplotype, antinuclear antibodies in early onset pauciarticular juvenile chronic arthritis (JCA) are associated with HLA-DOB1*0603, 461

phenotypes and gold-induced remissions in patients with rheumatoid arthritis, 343

Human model of ischaemia-reperfusion injury to endothelial cells, (A1) 32 Human peripheral blood dendritic cells and the extracellular matrix environment, in vitro maturation, (A1) 145

HUMPHRIES J, HUGHES R, TONG W, AKINTUNDE C, KEAT A. Screening of synovial fluid samples from patients with undifferentiated oligoarthritis for bacterial antigens using PCR, (A1) 116

HUNT A See Doherty M

HUNT B See Lima F

HUNT M See Creamer P

HUNTER J See Scott DL

HUNTER JA See Al-Mughales J; McInnes IB; Morrison E; Oien KA

HUNTER JO See Kavanagh R; King V

HUNTER M See Jobanputra P

HURLEY MV, NEWHAM DJ, JONES DW. Quadriceps activation and rehabilitation following anterior cruciate ligament rupture, (A1) 135

HURST MJ See D'Cruz D

HURST NP See Jobanputra P; Lambert CM; McRorie ER

HUSBY G See Stenstad T

HUSKISSON EC See Comer M

HUTCHINGS CJ See Kay RA

HUYGEN ACJ See Hogeweg JA

Hydrotherapy, out-patient, and home exercises, (L) 991, (L) 992

Hydroxychloroguine

can we differentiate between radiological progression over one year in patients with rheumatoid arthritis randomly allocated to im gold or hydroxychloroquine?, (A1) 107

safety in lupus pregnancy: experience in 27 pregnancies, (A1) 14 Hyperamylasemia, macroamylasemia: a possible cause of unexplained in

rheumatoid arthritis, 290 Hyperlipidaemia, increased VLDL levels and diminished renal excretion of

uric acid in hyperuricaemic-hypertriglyceridaemic patients, 920 Hyperlipoproteinaemia, musculoskeletal complaints, (L) 393

Hypermobility

with CREST syndrome: an unusual association, (A2) 54

diagnostic associations in new rheumatology referrals, (A1) 39 diagnostic associations rheumatology patients, 1157

Hypermobility syndrome, effect on knee joint proprioception, 121 Hyperuricaemia, increased VLDL levels and diminished renal excretion of

uric acid in hyperuricaemic-hypertriglyceridaemic patients, 920 Hypocomplementaemia, C3 nephritic factor and type III mesangiocapillary

glomerulonephritis progressing to systemic lupus erythematosus, (L) 90 Hypothalamic-pituitary-adrenal function in patients with fibromyalgia and chronic fatigue syndrome, (A1) 69

Hypoxia induces prolonged neutrophil adhesion to endothelial cells in vitro at physiological shear rates, (A1) 50

Hypoxic reperfusion injury and interleukin-8 (IL-8) in rheumatoid arthritis and seronegative polyarthritis, (A1) 50

Iatrogenic cost of non-steroidal anti-inflammatory drug therapy, (S1) 19 cICAM-1, cICAM-3 and cE-selectin levels in patients with rheumatoid vasculitis, (A1) 87 ICAM-1

E-selectin and P-selectin diurnal variation in levels in healthy volunteers, (A1) 87

relationship with E-selectin in serum, synovial fluid and synovial membrane in inflammatory arthritis, (A1) 148

ICHIKAWA Y See Satoh M

Identification

of an at risk group for osteoporosis using heel ultrasound measurement in general practice, (A1) 37

of genes expressed specifically during endochondreal ossification, (A1) 77 Identifying iron deficiency anaemia by simple laboratory measures in patients with rheumatoid arthritis, (A1) 83

Idiopathic juvenile osteoporosis: experience of twenty-one patients, 68 IgA deficiency, rheumatoid arthritis and primary selective IgA deficiency,

(A2) 3IgA-alpha-1-antitrypsin non-immune complex, inhibition of antibody dependent cellular cytotoxicity (ADCC) in rheumatoid arthritis, (A2) 30

IGARASHI A See Fujimoto M; Ihn H IGF-1, relationship with osteoarthritis in women aged 45-64 in the general population: the Chingford study, (A1) 137

IgG

1212

anti-DNA antibodies deposit in kidneys and induce proteinuria in SCID mice, (A1) 65

anticardiolipin antibodies

production and analysis of the binding characteristics of four human monoclonal, (A1) 12 sequence analysis of four human monoclonal, (A2) 28

anticardiolipin-antibody secreting clones, fetal calf serum in growth medium obscures the detection, (A1) 12 effects of reactive oxygen species on the ability of IgG to interact with

Clq, (A1) 45 reactivity, lack of temporal association of iridocyclitis with to core

histones and nucleosome subparticles in pauciarticular juvenile

chronic arthritis 507 IHN H, SHIMOZUMA M, FUJIMOTO M, SATO S, KIKUCHI K, IGARASHI A, SOMA Y, TAMAKI K, TAKEHARA K. Ultrasound measurement of skin thickness in systemic sclerosis, 535

IHN H See also Fujimoto M

Iliopsoas bursae, symptomatic enlarged in the presence of a normal plain hip radiograph, 365

Illness impact and disease status, defining in ankylosing spondylitis, 663 IM HOF V See Beyeler C

Immune complexes, complement and systemic lupus erythematosus, (A1) 82 Immunoblot analysis of precipitating and non-precipitating anti-ribonucleoprotein antibodies, (A1) 11

Immunoglobulin production clonality in rheumatoid synovium, (A1) 154 Impact of rheumatoid arthritis on body image, (A1) 89

Importance of musculo-skeletal pain in patients with irritable bowel syndrome, (A1) 119

In the rheumatoid patient: surgery to the cervical spine, 1078

In vitro duodenal iron uptake and serum and mucosal iron protein levels, with special reference to rheumatoid arthritis, 1041

In vitro effect of leflunomide on RA synovial fibroblasts proliferation and production of the matrix metallo-proteinase pro-MMP3, (A1) 60 In vitro immunomodulatory effect of tetracyclines in rheumatoid arthritis,

(A1) 152 In vitro maturation of human peripheral blood dendritic cells and the

extracellular matrix environment, (A1) 145 In-patient treatment for active rheumatoid arthritis: clinical course and

predictors of improvement, 847

INAMO Y, PEMBERTON S, TUDDENHAM EGD, WOO P. Increase of activated factor VIIA and haemostatic molecular markers in juvenile chronic arthritis, 466

Incidence

of juvenile dermatomyostitis: results from a nation-wide study, 732 of myopathy in patients treated with antimalarials. A report of three

cases and a review of the literature, 166 of rheumatoid arthritis is not related to socioeconomic factors, (A2) 7

of systemic vasculitis in the Norwich Health Authority, (A1) 21 Increase of activated factor VIIA and haemostatic molecular markers in

juvenile chronic arthritis, 466 Increased endothelium derived relaxing factor/nitric oxide production in

patients with systemic vasculitis, (A1) 23 Increased IL-1 receptor antagonist (IL-1ra) production and decreased IL-18/ IL-1ra ratio in mononuclear cells from rheumatoid arthritis patients, 24 Increased incidence of intermittent claudication in patients with systemic

sclerosis: a prospective study, (A1) 26 Increased interleukin-2 receptor expression on peripheral blood lymphocytes of patients with systemic lupus erythematosus, (A1) 6

Increased levels of antibodies to heat shock proteins with increasing age in

MRL/MP-LPR/LPR mice, 610

Increased levels of $\beta 2$ glycoprotein-1 antigen and $\beta 2$ glycoprotein-1 binding antibodies are associated with a history of thromboembolic complications in patients with SLE and primary antiphospholipid syndrome, 1031

Increased VLDL levels and diminished renal excretion of uric acid in hyperuricaemic-hypertriglyceridaemic patients, 920 India

Behçet's disease in: a clinical, immunological, immunogenetic and outcome study, 825

does MCTD exist?, (A2) 26 mixed connective tissue disease, 539

Indium-111 antimyosin antibody and muscle ultrasound scans in the evaluation of patients with suspected polymyositis, (A2) 27

Individual with ankylosing spondylitis: defining disease status and the impact of the illness, 663

Indomethacin, tiaprofenic acid and placebo in knee osteoarthritis: the long term outcome of therapy, (A2) 44

arthritogenic, reactive arthritis: urogenital swab culture is the only useful diagnostic method for detection in extra-articularly asymptomatic patients with undifferentiated oligoarthritis, 838

late, role of prophylactic antibiotics in the prevention of in prosthetic joints. Results of a questionnaire and review of the literature, 380

Infectious diseases, false positive IgA and IgG rheumatoid factor isotypes, (A1) 109

Inflammation

chronic OM-89 modulation of: relevance to clinical use, 525 neurogenic, consequences of sustained activation of small sensory fibres, (A1) 53

Inflammatory activity, rheumatoid arthritis patients cannot accurately report signs, 547

Inflammatory arthritis

antineutrophil cytoplasmic antibodies—potential for misdiagnosis, 820 luteinised unruptured follicles in women taking NSAIDs, (A2) 22 predictive value of proteoglycan antibodies for erosions, (A1) 104 predictors of resolution in, (A1) 85

relationship of ICAM-1 with E-selectin in serum, synovial fluid and synovial membrane, (A1) 148

Inflammatory back pain in primary care, 1074

Inflammatory bowel disease, risk factors for a patient with spondylarthropathy to develop, (A1) 116

Inflammatory cytokine responses in juvenile chronic arthritis, 454 Influence

of glutathione S-transferase (GSTM1 and GSTT1) polymorphisms on the incidence and severity of rheumatoid arthritis, (A2) 18

of knowledge, attitudes and personality characteristics on the outcome of acupuncture treatment, (A2) 51

physical, psychological and behavioural factors on consultations for back pain, 156

of sulphasalazine on the evolution of gut inflammation and locomotor manifestations in the spondylarthropathies, (A1) 115

INGMELO M See Pons F

Inhibition of antibody dependent cellular cytotoxicity (ADCC) in rheumatoid arthritis by the non-immune complex of IgA-alpha-1-antitrypsin, (A2) 30 Injection technique, intra-articular and soft tissue: a survey of current practice, 449

Inner ear involvement in mixed cryogolbulinaemia patients, 370

Insulin-like growth factor-1: serum and synovial fluid levels and associations with markers of cartilage turnover in osteoarthritis and rheumatoid arthritis, (A1) 55

Integrins, chondrocyte: increase in α5 and αV on stimulation with inflammatory synovial fluid, (A1) 57

Interferon-alpha

autoimmune phenomena associated with cutaneous aseptic necrosis during treatment for chronic myelogenous leukaemia, (L) 582 is it a therapy for hepatitis B-associated polyarteritis nodosa, (L) 486

Interferon-gamma

determination and of type III procollagen aminopeptide levels in bronchoalveolar lavage fluid of the patients with pulmonary fibrosis: inverse correlation between type III procollagen aminopeptide and interferon- Γ in progressive patients, 31 and epithelial cell activation in Sjögren's syndrome, 226

Interleukin IL-1\beta/IL-1ra ratio decrease and IL-1 receptor antagonist (IL-1ra) production increase in mononuclear cells from rheumatoid arthritis patients, 24

Interleukin-1 (IL-1) and IL-1ra production by monocyte/macrophage in rheumatoid arthritis synovial membranes, (A1) 150

Interleukin-1 receptor antagonist (IL-1ra)

defective spontaneous and bacterial lipopolysaccharide-stimulated production by polymorphonuclear neutrophils of patients with active systemic lupus erythematosus, 107 production increase and decreased IL-1\(\beta/\text{IL-1ra}\) ratio in mononuclear

cells from rheumatoid arthritis patients, 24

Interleukin-1α (IL-1α) intron 6 polymorphism, absence of correlation with rheumatoid arthritis, 1123 Interleukin-2 in psoriatic arthritis and seronegative spondylarthritis but not

in rheumatoid arthritis synovium, (A1) 151

Interleukin-2 receptor

IL-2R levels 'n rheumatoid arthritis: poor correlation with clinical activity is due in part to disease duration, 1037 increased expression on peripheral blood lymphocytes of patients with

systemic lupus erythematosus, (A1) 6

Interleukin-4 (IL-4) and IL-6 role in IL-1-dependent cartilage matrix degradation, 101 Interleukin-6 (IL-6)

effect of minocycline on serum levels in rheumatoid arthritis patients, (A1) 96

and IL-4 role in IL-1-dependent cartilage matrix degradation, 101 induces the proliferation of synovial fibroblastic cells in the presence of soluble IL-6 receptor, 321

soluble receptor, interleukin-6 (IL-6) induces the proliferation of synovial fibroblastic cells in the presence of, 321

Interleukin-8 (IL-8) and hypoxic reperfusion injury in rheumatoid arthritis and seronegative polyarthritis, (A1) 50

Intermittent claudication, increased incidence in patients with systemic sclerosis: a prospective study, (A1) 26

Intestinal strontium absorption and axial bone mineral density, (A1) 75 Intra-articular anaesthetic, pain mechanisms in osteoarthritis of the knee, (A2) 45

Intra-articular pressure changes in normal and rheumatoid peripheral joints, (A1) 46

Intra-articular and soft tissue injections: a survey of current practice, 449 Intra-muscular versus intra-articular depo-methyl-prednisolone in the management of ambulatory patients with active rheumatoid arthritis, (A1)

Intravenous cyclophosphamide in the treatment of systemic vasculitis: a five year follow-up study, (A1) 22

Introduction: decisions in non-steroidal anti-inflammatory prescribing, (S1) 1 Involvement of the gut in the pathogenesis of inflammatory synovitis, (E) 801 IOAKIMIDIS D See Athanassiou P

Iridocyclitis, lack of temporal association with IgG reactivities to core histones and nucleosome subparticles in pauciarticular juvenile chronic arthritis, 507

Iritis, antibodies to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of ankylosing spondylitis patients with/without iritis and enthesitis, 418

Iron, duodenal uptake and serum and mucosal iron protein levels, with special reference to rheumatoid arthritis, 1041

Iron deficiency anaemia, identifying by simple laboratory measures in patients with rheumatoid arthritis, (A1) 83

Irreversible visual loss in dermatomyositis, 1179

Irritable bone syndrome—a new concept in somatisation?, (A2) 52

Irritable bowel syndrome, importance of musculo-skeletal pain, (A1) 119

IRVEN C See Loughlin J IRVINE AE See Magill MK

IRVING M See Herrick AL

Is interferon alpha a therapy for hepatitis B-associated polyarteritis nodosa?, (L) 486

Is remitting seronegative symmetrical synovitis with pitting oedema (RS3PE syndrome) associated with HLA-A2?, (L) 889

Ischaemia-reperfusion injury to endothelial cells, (A1) 32

ISENBERG D See Casey MC; Faulds G

ISENBERG DA See Ehrenstein MR; Gordon C; Kausman D; McNally T; Menon S; Rahman MAA; Rankin ECC; Rose LM; Sutcliffe N; Tucker LB; Whitefield L

Isolated nail fold rheumatoid vasculitis-a benign condition?, (A1) 68

Isomaki P See Roivainen A

Isotype variation of TNF monoclonal antibodies. Is there a therapeutic difference in rheumatoid arthritis?, (A1) 153

IVERSEN MD See Daltroy LH

JACKSON SE See Dziedzic K

JACOBS JWG See Geenen R; Lems WF

JAHANGIER ZN See Lems WF

JAMAR F See Chapman PT

JAMES D, COX N, DAVIES P, DEVLIN J, DIXIE J, EMERY P, GALLIVAN S, GOUGH A, PROUSE P, WILLIAMS P, WINFIELD J, YOUNG A. Major life events as a result of rheumatoid arthritis. A comparison between nine centres in 446 patients in the first three years of disease, (A2) 9

JAMES D See also Prouse P JAMES IT, BIRCH A, CROWLEY C, PERRETT D. Stability of pyridinium crosslinks to photolysis, (A1) 54

JAMES MF See Byrne J

JANSSEN M See ten Wolde S

JARA LJ See Medina F

JAWAD ASM, AL NAHHAS A, McCREADY VR. The use of three-phase bone scan in the diagnosis of repetitive strain injury, (A2) 39
JAWED S, JOLLIFFE VA, CRUWYS SC, BLAKE DR, KIDD BL. The effects of

upper motor neurons on the cutaneous response to topical capsaicin, (A1) 53

JAWED S See also Byrne J; Donnelly S JAYSON MIV See Anderson ME; Bankhead CR; Blann AD; Brammah TB; Bukhari M; Chen ACN; Croft PR; Fakhoury H; Herrick AL; Teh LS; Wright D

JELLIS JE See Njobvu PD

JENKINS K See Jobanputra P

JENKINSON TR See Kennedy LG

JESSOP DS See Harbuz MS

JESSOP SJ. Cutaneous leucocytoclastic vasculitis: a clinical and aetiological study, 942

JEVTIC V, ROZMAN B, WATT I, PRESETNIK M. Use of a contrast enhanced MRI in the assessment of therapeutic response to a disease-modifying antirheumatic drug, 956

JEWKES RF See Misra R

JIAN-MEI L See Venables PJW

JOBANPUTRA P, DAVIDSON F, GRAHAM S, O'NEIL H, SIMMONDS P, YAP PL. High frequency of parvovirus B19 in serum samples submitted for a rheumatoid factor test, (A2) 33 JOBANPUTRA P, HUNTER M, CLARK D, LAMBERT CM, HURST NP. An audit of methotrexate and folic acid for rheumatoid arthritis. Experience from a teaching centre, 971

JOBANPUTRA P, LIN H, JENKINS K, BAVINGTON C, SALTER D, NUKI G. Chondrocyte integrins: increase in a5 and aV on stimulation with inflammatory synovial fluid, (A1) 57

JOBANPUTRA P See also Brennan FR; Wright M

JOFRE R See González M

JOHNS S See Seed MP

JOHNSON AE See Waterhouse L

JOHNSON ES See Avina-Zubieta JA

JOHNSON J See Kassimos D

JOHNSON S See Kirwan JR

JOHNSON SM, COLLINS DA, PATEL S, BRUCKNER FE. Mycoplasmal RNA and 5'-nucleotidase in the synovial fluid of rheumatoid arthritis patients, (A2)5

JOHNSTON GD See Bruce IN

Joint

bacterial infection in England & Wales 1990-1993, (A2) 31

damage and health status divergence in rheumatoid arthritis, (A1) 107 disease, pyridinoline and deoxypyridinoline may be influenced by

turnover of non-osseous tissues, (L) 1186 erosions and patients with early rheumatoid arthritis, (S2) 74

hyperlaxity: quantification according to age, (A1) 39

implant loosening, analysis of cell types and mediator production from tissues, 1127

inflammation, joints count: a review of old and new articular indices. (E) 1003

laxity quantification. The non-dominant (Spanish) modification, (L) 795 manifestations of Marfan syndrome: joint and skin manifestations are prevalent and correlated, 126

pain, proinflammatory role of tissue kallikrein in modulating in inflamed ioints. (L) 88

space width measured more accurately and precisely in semi-flexed than extended view of OA knees, (A1) 121

washout effects and steroid injection compared with either joint washout or steroid injection alone in rheumatoid knee effusion, 771

Joints count: a review of old and new articular indices of joint inflammation, (E) 1003

JOLLIFFE V See Gaffney K

JOLLIFFE VA See Jawed S; Watts RA

JONES AC, REGAN M, DOHERTY M. Psychological distress and local tenderness but not radiographic change or weakness are associated with disability in hospital referred patients with knee osteoarthritis, (A1) 136

JONES AC, REGAN M, DOHERTY M. Tibiofemoral osteophyte in the anteroposterior radiograph predicts joint space narrowing in the patellofemoral joint, (A1) 136

JONES AC, REGAN M, DOHERTY M. Determinants of arthrogenic inhibition in knee osteoarthritis, (A1) 134

JONES AC See also O'Reilly SC

JONES AKP See Chen ACN

JONES B See Lau CS

JONES C See Thompson PW JONES DW See Hurley MV

JONES G See Sambrook PN

JONES H See Dai L: Gordon P

JONES HW, BAILEY R, DUNNE C, ZHANG Z, COX NL, MORRIS CJ, WINYARD PG. Antithrombin III inactivation in rheumatoid arthritis, (A1) 84

JONES HW, KUS M, FAIRBURN K, LAMB DJ, LEAKE DS, COX NL, BLAKE DR, WINYARD PG. Enhanced macrophage degradation of low density lipoprotein from synovial fluid of rheumatoid arthritis patients, (A1) 49

JONES M, SILMAN A, WHITING S, BARRETT B, SYMMONS D. The familial risk for rheumatoid arthritis is low in first degree relatives of probands with inflammatory arthritis ascertained from a population-based register, (A1) 80

JONES P See Badcock LJ; Hassell A; Nazhat NB; Ollier WER

JONES PBB, MALIK S, BANKHEAD C. SCOTT DGI, CHAKRAVARTY K, SYMMONS D. Predictive value of proteoglycan antibodies for erosions in early inflammatory arthritis, (A1) 104

JONES PJ See Popert AJ

JONES PW See Averns HL; Borg AA; Cheung NT; Deodhar AA; Dziedzic K; Griffiths SM; Hassell AB; Plant MJ; Russell G

JONES R See Doherty M
JONES SD, GARRETT SL, CAPPER R, CANTER RJ, CALIN A. Ankylosing spondylitis, and tonsillectomy, (A2) 35

JONES SD, STEINER A, GARRETT SL, CALIN A. The Bath ankylosing spondylitis patient global score, (A1) 114

JONES SD See also Calin A; Koh WH

JORDI B See Beyeler C

JORGENSEN C, MOYNIER M, BOLOGNA C, YOUINOU P, SANY J. Rheumatoid factor associated with a secretory component in rheumatoid arthritis,

JOVANI V See Pascual E

JULKUNEN H, SIREN M-K, KAAJA R, KURKI P, FRIMAN C, KOSKIMIES S. Maternal HLA antigens and antibodies to SS-A/RO and SS-B/LA. Comparison with systemic lupus erythematosus and primary Sjögren's syndrome, 901

JUMP A See Darlington LG

JUNKER P See Hansen BL Juvenile chronic arthritis

analysis of childhood relationships in adults with, (A1) 90

antinuclear antibodies in early onset pauciarticular are associated with HLA-DQB1*0603: a possible JVA-associated human leucocyte antigen haplotype, 461 antiphospholipid antibodies in, 873

association between physical function and psychological problems in children, 470

hip involvement in evaluation with contrast-enhanced MR imaging, (A1) 124

increase of activated factor VIIA and haemostatic molecular markers. 466

inflammatory cytokine responses in, 454

NUMACT: a novel method of monitoring physical activity, (A2) 23 pain threshold, 61

pauciarticular, lack of temporal association of iridocyclitis with IgG reactivities to core histones and nucleosome subparticles, 507 systemic-onset, successful management of reactive haemophagocytic

syndrome, (L) 888 T-cell receptor variable gene segment TCRBV6S1 null allele frequencies in UK. (A1) 144

Juvenile dermatomyositis: results from a nation-wide study, 732

Juvenile onset ankylosing spondylitis-more girls than we thought?, (A2) 34 Juvenile onset Still's disease in Northern India: comparison with adult onset Still's disease, 429

KAAJA R See Julkunen H

KAANDORP CJE, DINANT HJ, VAN SCHAARDENBURG D, VAN DER LAAR MAFJ. Septic arthritis in Amsterdam: potential targets for prevention, (A1) 113

KABASAKAL Y, GARRETT SL, CALIN A. Outcome of total hip replacement in ankylosing spondylitis: a thirteen year follow-up study, (A1) 114 KADIR M See Hameed K

KAHN MF See Bailly S; Kaye O

KAILASH S See Uppal SS

KAKKER R See Uppal SS

KALDEN JR, MANGER B. New therapeutic approaches in autoimmune rheumatic diseases, with special emphasis on rheumatoid arthritis, (E)

KALDEN JR See also Ehrenstein MR

KALLA AA, SMITH PR, BROWN GMM, MEYERS OL, CHALTON D. Responsiveness of Keitel functional index compared with laboratory measures of disease activity in rheumatoid arthritis, 141

Kallikrein, tissue, proinflammatory role in modulating pain in inflamed joints, (L) 88

KAMINSKI A See Kaminski ER

KAMINSKI ER, KAMINSKI A, PEREIRA RS, DUKE OL. The in vitro immunomodulatory effect of tetracyclines in rheumatoid arthritis, (A1)

KANIS JA See O'Neill TW

KANTEREWICZ E. Pseudogout masking brucellar arthritis, (L) 294

KARIM SN, MURPHY EA, STURROCK RD, GOUDIE RB. T-cell clonality in synovial fluid from rheumatoid joints and after culture in interleukin-2, 232

KARSTENSEN B See Glennås A

KASSIMOS D, CHOY EHS, CHIKANZA I, GROSSMAN A, PANAYI GS. Endogenous opiates: are they responsible for the defective hypothalamic response to inflammatory stimuli in patients with rheumatoid arthritis?, (A1) 84

KASSIMOS D, JOHNSON J, KINGSLEY GH. Lumbar epidural steroid injection in sciatica: who benefits?, (A1) 40

KASSIMOS D See also Dolan AL; Rankin ECC

KASSIMOS DG, CHOY EHS, CHALLACOMBE SJ, PANAYI GS. The prevalence of xerostomia in a population with primary generalised osteoarthritis, (A1) 132

KATOH O See Kuroki S

KATZ DR See Binks MH; Ehrenstein MR

KAUSMAN D, ISENBERG DA. Thyroid autoimmunity in systemic lupus erythematosus: the clinical significance of a fluctuating course, 361

KAVANAGH R, RYAN MJ, WALL P, HAZLEMAN BL. Bacterial joint infection in England & Wales 1990-1993, (A2) 31

KAVANAGH R, WORKMAN E, NASH P, SMITH M, HAZLEMAN BL, HUNTER JO. The effects of elemental diet and subsequent food reintroduction on rheumatoid arthritis, 270

KAVANAGH R See also Vecchio P

Kawasaki disease

adult, symptomatic splenic infarction complicating, (L) 579

anti-neutrophil cytoplasmic antibodies and anti-endothelial cell antibodies are not increased, 882

KAY LJ, LILLEY M, WALKER DJ. HLA associations of different patterns of psoriatic arthritis, (A1) 143

KAY LJ, VALE S, BORG AA, WALKER DJ. Second-line drug experience of patients with rheumatoid arthritis receiving lower limb arthroplasty, (A2) 20

KAY RA, ROBINSON MA, OLLIER WER, HOLT PJL, BERNSTEIN RM, HUTCHINGS CJ. Mapping the genetic association between hypergammaglobulinaemic primary Sjögren's syndrome and the TRCB gene complex, (A1) 77

KAY RA See also Donn RP; Farhan AJ

KAY SPJ See Beevers DJ

KAYE O, PALAZZO E, GROSSIN M, BOURGEOIS P, KAHN MF, MALAISE MG. Low-dose methotrexate: an effective corticosteroid-sparing agent in the musculoskeletal manifestations of sarcoidosis, 642

KAYE S See Marguerie C

KAYE SA, LIM SG, TAYLOR M, PATEL S, GILLESPIE S, BLACK CM. Small bowel bacterial overgrowth in systemic sclerosis: detection using direct and indirect methods and treatment outcome, 265

KEANEY NP See Hassan WU

KEAT A See Humphries J KEATING NA See Yeap SS

KEELAN ETM See Chapman PT

KEELING J See Etherington J

KEEN RW, ARDEN NK, MAJOR PJ, MORRISON NA, KELLY PJ, BAKER JR, SAMBROOK PN, LANCHBURY JS, EISMAN JA, SPECTOR TD, VITAMIN D receptor gene alleles and bone density in postmenopausal twins, (A1) 82

KEEN RW, MAJOR PJ, HARRIS P, LANCHBURY JS, SPECTOR TD. Polymorphisms of the vitamin D receptor gene in postmenopausal female athletes, (A1) 79

KEEN RW, MAJOR PJ, NANDRA D, LANCHBURY JS, SPECTOR TD. Early postmenopausal bone loss and polymorphisms of the vitamin D receptor, (A1) 79 KEENAN J See Ferry S

KEIR G, DICKIE A, LIGGETT N, RENNIE JAN, GARTON MJ. Prevalence of serum paraprotein bands in rheumatoid arthritis, (A2) 5

Keitel functional index responsiveness compared with laboratory measures of disease activity in rheumatoid arthritis, 141

KELLEHER D See Donnelly S

KELLY CA See Hassan WU KELLY PJ See Keen RW

KENNEDY LG, BENLAHRACHE S, REMITA K, CALIN A. Evidence for a possible gene pool difference between ethnic groups?-a controlled study of British and Algerian subjects, (A1) 143

KENNEDY LG, JENKINSON TR, MALLORIE PA, WHITELOCK HC, GARRETT SL, CALIN A. Ankylosing spondylitis: the correlation between a new metrology score and radiology, 767

KENNEDY LG See also Brown MA; Calin A; Garrett SL

KENNEDY TD See Yeap SS

Keratan sulphate epitopes, chondroitin sulphate, glycosaminoglycans and hyaluronan in normal knee synovial fluid, (A1) 55 Keratoconjunctivitis sicca, HLA-DR4 and protease inhibitor phenotypes

association with pulmonary abnormalities in rheumatoid arthritis, 37 KERR JR, CARTRON JP, CURRAN MD, MOORE JE, ELLIOTT JRM, MOLLAN

RAB. A study of the role of parvovirus B19 in rheumatoid arthritis, 809 KERRY R See Averns HL

KERSLAKE S See Buchanan NMM; Etherington J; Lima F

KESER G See D'Cruz D

KHAMASHTA MA, ASHERSON RA. Hughes syndrome: antiphospholipid antibodies move closer to thrombosis in 1994, (E) 493 KHAMASHTA MA See also Buchanan NMM; D'Cruz D; Froes L; Lima F;

Mujic F; Rosman A

KHAN FM See Scott DL

KIDD BL See Cruwys SC; Jawed S

KIELY PDW, GILLESPIE KM, OLIVEIRA DBG. Oxpentifylline enhances the Th2 environment and inhibits TNFα mRNA production and inflammatory arthritis in mercuric chloride-induced autoimmunity, (A2) 29

KIENER H See Koeller M

KIFF P See Walker DJ; Wright D KIKUCHI K See Fujimoto M; Ihn H

KIMOFF J See Donald F

Kinetics of von Willebrand factor and angiotensin converting enzyme release by endothelial cells, (A1) 23

KING RH See Vecchio P

KING V, HUNTER JO. Importance of musculo-skeletal pain in patients with irritable bowel syndrome, (A1) 119

KINGSLEY G. They turned down my abstract again, (E) 303

KINGSLEY GH See Campbell F; Choy EHS; Dolan AL; Kassimos D; Rankin ECC

KINGSNORTH AN See Francombe J

KINGSTON AE See Billingham MEJ

KINNINMONTH A See McInnes IB

KIRBY JT See Close DR

KIRK AP See Camp AV; Collier SM

KIRK G See Maple C: Veale DJ

KIRKHAM B See Hill C

KIRWAN JR, ARC Low Dose Corticosteroid Study Group. Radiological progression in rheumatoid arthritis during a randomised controlled trial of prednisolone, (A1) 80

KIRWAN JR, COLES CJ, JOHNSON S. The relationship between synovitis and erosions in rheumatoid arthritis, (A1) 105

KIRWAN JR. Growth of scientific interest in rheumatology, (L) 796

KIRWAN JR See also Brooks P; Hewlett SE; Sharif M; Thompson PW

Кізнімото Т See Mihara M

KISKINIS D See Kitas GD

KISS CS, MITUSZOVA M, FELSENBERG D, O'NEILL T, SILMAN A, POÓR GY. Risk of vertebral fracture in DISH, (A1) 127

KITAS GD, SARATZIS N, MCBURNEY A, KISKINIS D, BACON PA, LUNEC J. A human model of ischaemia-reperfusion injury to endothelial cells, (A1)

KITAS GD, ZWEDE M, BACON PA. Factors promoting endothelial cell apoptosis in vitro, (A1) 32

KITAS GD See also McRorie ER; Zwede M

KITNEY RI See Carter ME

Klebsiella pneumoniae

Escherichia coli and Proteus mirabilis antibodies in the sera of patients with axial and peripheral form of ankylosing spondylitis, 413

Proteus mirabilis and Escherichia coli antibodies in the sera of ankylosing spondylitis patients with/without iritis and enthesitis, 418

KLENERMAN L. The foot and ankle in rheumatoid arthritis, 443

KLINKE J See Brown MA

KLOPPENBURG M, VERWEIJ CL, BREEDVELD FC, DIJKMANS BAC. Effect of minocycline on acute phase proteins and serum interleukin-6 levels in rheumatoid arthritis patients, (A1) 96

KLOVRZA L See Griffiths B Knee

painful associated with a septic focus, 862

rheumatoid effusion, joint washout effects and steroid injection compared with either joint washout or steroid injection alone, 771 Knee joint

determination of the apparent synovial permeability in patients suffering from osteoarthritis and rheumatoid arthritis. 520

dual energy X-ray absorptiometry of the knee joint, (A1) 122

effect of hypermobility syndrome on proprioception, 121

magnetic resonance images: enhancement by advanced 3D computer visualisation, (A1) 123

relationship between serum cartilage oligomeric matrix protein levels and disease progression in osteoarthritis, 306

space, radiographic within-subject variability is low when measuring with digital image analysis, (A1) 122

Knee osteoarthritis

determinants of arthrogenic inhibition. (A1) 134

psychological distress and local tenderness but not radiographic change or weakness are associated with disability in hospital referred patients, (A1) 136

shortly after anterior cruciate ligament transection dogs develop characteristics similar to human, (A1) 135 symmetrical compartmental involvement, (A1) 134

anterior in thalassaemia major treated with L1, a new iron chelating agent, (A1) 119

and disability estimates in the community with differing knee pain criteria, (A1) 36

disability and psychological variables in a Nottingham population, (A1) 35

quadriceps weakness and muscle activation in the community, (A2) 45 KNEKT P See Heliövaara M

Ko OKH See Lau CS

KOELLER M, KIENER H, SIMONITSCH I, ARINGER M, STEINER CW, MACHOLD K, Graninger W. Destructive lymphadenopathy and T-lymphocyte activation in adult-onset Still's disease, 978

KOH W-H. Atypical antineutrophil cytoplasmic antibodies in patients with ankylosing spondylitis, (L) 695

KOH WH, JONES SD, GARRETT SL, STEINER A, CALIN A. The relationship of fatigue in ankylosing spondylitis to disease activity, sleep and other factors, (A2) 36

KONTOMERKOS T See Athanassiou P

KOREICHI A See SARAUX A

KOSKIMIES S See Julkunen H KRAAIMAAT FW See Geenen R

KREMER JM. Possible mechanisms of action of methotrexate in patients with rheumatoid arthritis, (S2) 26

KÜÇÜKDEVEÇI AA. Tarsal tunnel syndrome in ankylosing spondylitis, (L) 488 Kuis W See Hogeweg JA

KUMAR A See Sood A; Uppal SS

KUNTZ D See Lioté F

KURKI P See Julkunen H

KUROKI S, OHTA A, SUEOKA N, KATOH O, YAMADA H, YAMAGUCHI M. Determination of various cytokines and type III procollagen aminopeptide levels in bronchoalveolar lavage fluid of the patients with pulmonary fibrosis: inverse correlation between type III procollagen aminopeptide and interferon-I in progressive patients, 31

Kus M See Jones HW

KUWANA M See Satoh M KVALE G See Brun JG

KVIEN TK See Glennås A

KYLE PM See Hamilton JD

KYLE VA See Cawston TE

L1, a new iron chelating agent, anterior knee pain in thalassaemia major treated with, (A1) 119

LA CIVITA L See Berrettini S

LAAN RFJM See Van Gestel AM

LAASONEN L See Scott DL

LACEY RW. The rheumato-microbiology interface, (E) 99

Lack

of interaction between methotrexate and flucloxacillin in patients with rheumatoid arthritis, (A1) 100

of temporal association of iridocyclitis with IgG reactivities to core histones and nucleosome subparticles in pauciarticular juvenile chronic arthritis, 507

LADJOUZE A See Claudepierre P

LAFEBER FPJG See van Valburg AA

LAFFAN M See Davies KA

LAFFÓN A, GONZÁLEZ-AMARO R. Cell adhesion molecules: an overview, (E) 1101

LAFFORGUE P See Combe B

LAI N-S, LAN J-L, YU C-L, LIN R-H. Dominant human leukocyte antigen DR4-Dw15 restricted mycobacterial 65 kDa antigen-specific T-cell immunity in Chinese patients with rheumatoid arthritis, 498 LAMB DJ See Jones HW

LAMBERT CM, HURST NP. Health economics as an aspect of health outcome: basic principles and application in rheumatoid arthritis, 774

LAMBERT CM See also Jobanputra P

Lamellar body secretion: ultrastructural analysis of an unexplored function of synoviocytes, 13

LAMERS CBHW See ten Wolde S

LAMPREAVE JL See Alonso JC

LAN J-L See Lai N-S

LANCHBURY J See Bajocchi GL; Campbell F LANCHBURY JS See Keen RW

LAND D See Munro R

LANDEWE RBM See van den Borne BEEM

LAPRISE R See Spitzer WO

Larsen's scoring methods for hand and wrist radiographs, proposed modification, 56

LASSERE M, EDMONDS J, HOUSSEIN D, SCOTT D. Centile curves of radiological damage: rheumatoid arthritis population, (A1) 106

LATCHMAN D See Faulds G

LATCHMAN DS See Ehrenstein MR; Rahman MAA; Rose LM

Late-onset and young-onset rheumatoid arthritis. A comparative clinical and radiological study, (A1) 106

LAU CS, CHAN EYT, KO OKH, WONG D. Increased interleukin-2 receptor expression on peripheral blood lymphocytes of patients with systemic lupus erythematosus, (A1) 6

LAU CS, JONES B, WONG RWS. CD5-positive and CD5-negative plaqueforming cells against poly-L-lysine treated sheep erythrocytes in patients with systemic lupus erythematosus, (A1) 6

LAU CS, WONG D, MOK CC, CHAN EYT. Assessing systemic lupus erythematosus activity using plasma factor VIII von Willebrand factor antigen, (A1) 8

LAU CS See also Lau YL; Mok CC

LAU E, LIN F, CROFT P. Acetabular dysplasia does not cause hip osteoarthritis in a Chinese population, ($\dot{A}\dot{I}$) 34 Lau YL, Chan SY, Lau CS. Mannose binding protein and systemic lupus

erythematosus in southern Chinese, (A1) 7

LAVERSUCH CJ, BROWN MM, CLIFTON A, BOURKE BE. Cerebral venous thrombosis and acquired protein S deficiency: an uncommon cause of headache in systemic lupus erythematosus, 572

LAVERSUCH CJ, COLLINS DA, CHARLES PJ, BOURKE BE. Sulphasalazine-

induced autoimmune abnormalities in patients with rheumatic disease, 435

LE CESSIE S See Van Den Ende CHM

LE GALLEZ P See Bradley SM

LE GOFF P See SARAUX A

LE LOET X See Wylie G LE THI HUONG D. Symptomatic splenic infarction complicating adult Kawasaki disease, (L) 579 LE THI HUONG D. Wegener's granulomatosis and CREST syndrome, (L) 484

LEAKE DS See Jones HW

LEAKER B See Worrall JG

LEDINGHAM J, REGAN M, DOHERTY M. Symmetrical compartmental involvement in knee osteoarthritis, (A1) 134

LEE A See Bajocchi GL; Pitzalis C

LEE WR See Hamilton JD

LEEMING MRG See Wylie G

Leflunomide effects on RA synovial fibroblasts proliferation and production of the matrix metallo-proteinase pro-MMP3, (A1) 60

LEHTINEN K See Mäki-Ikola O

LEIRISALO-REPO M See Mäki-Ikola O; Paimela L

LEMS WF, GERRITS MI, JACOBS JWG, VAN VUGT RM, VAN RIJN HJM, BILISMA JWJ. Effect of high-dose corticosteroid pulse therapy on bone metabolism in rheumatoid arthritis, (A1) 128

LEMS WF, JAHANGIER ZN, JACOBS JWG, BIJLSMA JWJ. Prevalence of vertebral fractures in patients with RA, treated with corticosteroids, (A1)

LENTJES E See Griep EN LETHBRIDGE-CEJKU M See Lim K

LETROUBLON MC See Tebib JG

Letter

from abroad, 171

from Australia, 57

Leucocyte adhesion, synovial microvascular endothelial cells exhibit enhanced shear rate dependent in vitro, (A1) 51

Leukaemia, chronic myelogenous, autoimmune phenomena associated with cutaneous aseptic necrosis during interferon-alpha treatment, (L) 582 Levels

of chondroitin sulphate and keratan sulphate epitopes, glycosaminoglycans and hyaluronan in normal knee synovial fluid, (A1) 55

of cICAM-1, cICAM-3 and cE-selectin in patients with rheumatoid vasculitis, (A1) 87

of circulating intercellular adhesion molecule-1 and -3 but not circulating endothelial leukocyte adhesion molecule are increased in patients with rheumatoid arthritis, 311

of collagenase, tissue inhibitor of metalloproteinase (TIMP), collagenase-TIMP complexes and glycosaminoglycan in sequential samples of synovial fluid aspirated from patients with osteoarthritis, (A1) 138

of urokinase and plasminogen activator inhibitor-1 in synovial fluids in normal, osteoarthritis and rheumatoid arthritis knees, (A1) 59

LÉVESQUE H. Autoimmune phenomena associated with cutaneous aseptic necrosis during interferon-alpha treatment for chronic myelogenous leukaemia, (L) 582

LEWIS RJ. Fatal vasculitis following treatment with co-trimoxazole, (L) 84 LEWIS S, CROSSMAN M, DOHERTY M, BAYLISS M, MASON R. Characterization by capillary electrophoresis of synovial fluid chondroitin sulphates

in arthritis, (A1) 54 Lewis X expression and α2,3 and α2,6 linked sialylation on serum IgG in rheumatoid arthritis, (A1) 63

Lhermitte's sign in Behçet's disease, (L) 184

LIANG MH See Daltroy LH

LIEW FY See McInnes IB

Life-style, environmental and medical factors influencing peak bone mass in women, 620

LIGGETT N See Keir G

LIGHTMAN SL See Harbuz MS LIGHTOWLER C See D'Cruz D

LILLEY M See Kay LJ

LIM K, SHEPSTONE L, LETHBRIDGE-CEJKU M, HOCHBERG M, TOBIN J, DIEPPE PA. The pattern of distribution of osteoarthritis of the hand joints: comparative study between a clinical and a community cohort, (A1) 133

LIM K See also Pallis M. LIM KL See Neilly JB

LIM SG See Kaye SA

LIMA F, KHAMASHTA MA, KERSLAKE S, BUCHANAN N, HUNT B, HUGHES GRV. A prospective study of 110 high-risk pregnancies in a lupus pregnancy clinic, (A1) 13

LIMA F See also Buchanan NMM; Etherington J

Limited association of Helicobacter pylori in gastric ulcer patients with rheumatic disease, (L) 1188

LIN F See Lau E

LIN H See Jobanputra P

Lin R-H See Lai N-S Lindroth Y, Bauman A, Brooks PM, Priestley D. A 5-year follow-up of a controlled trial of an arthritis education programme, 647

LINK GROUP See Scott DL

Linkage

analysis in multiple epiphyseal dysplasia, (A1) 144

of the gene that encodes the al chain of type V collagen (COL5A1) to type II Ehlers-Danlos syndrome, (A2) 40

and penetrance of HLA and rheumatoid arthritis in extended pedigrees.

LIOTÉ F. Lymphocyte subsets in pamidronate-induced lymphopenia, (L) 993 LIOTE F, VASERMAN N, CHICHEPORTICHE V, DRYLL A, KUNTZ D. Late-onset and young-onset rheumatoid arthritis. A comparative clinical and radiological study. (A1) 106

Lipocortin-1 intracellular expression, corticosteroid regulation in murine leukocytes, (A1) 147

Liposome, effect of liposomally conjugated methotrexate upon mediator release from human peripheral blood monocytes, 241

LLOYD M See Mujic F

LLOYD ME, ARDEN NK, NANDRA D, DOYLE DV, SPECTOR TD. Effect of thiazide use on bone density: the Chingford study, (A1) 129

LLOYD ME, ARDEN NK, NANDRA D, DOYLE DV, SPECTOR TD. Serum electrolytes and bone density: the Chingford study, (A1) 129

LLOYD ME See also Froes L

Localisation of a gene for familial chondrocalcinosis in a UK kindred to chromosome 5P, (A2) 41

LOCKE IC See Worrall JG

LOCKWOOD CM See Luqmani RA

Locomotor manifestations, influence of sulphasalazine on the evolution in the spondylarthropathies, (A1) 115

LOETSCHER P See Seitz M

LOGAN JA, MORRISON E, McGILL PE. Serum urate acute gout, (A2) 40 LOHMANDER LS. Molecular markers to monitor outcome and intervention in

osteoarthritis (promises, promises...), (L) 599 LOIZOU S See Davies KA; Mackworth-Young CG

LOMBARDINI F See Berrettini S

LONG A See Donnelly S

LONG F See Scott DL

Long-term outcomes in rheumatoid arthritis. (S2) 59

Long-term usage and side-effect profile of sulphasalazine in rheumatoid arthritis, (S2) 40

LONGHURST CM See Rahman MAA

LONGOMBARDO G See Berrettini S LOPEZ PINO MA See Gonzalez-Crespo MR

LÓPEZ-GOMEZ JM See González M

LÓPEZ-LONGO FJ See Abad MA; González CM; González M; Monteagudo I; Moreno FC

LOPEZ-LONGO FJ, ESCALONA M, SÁNCHEZ E, GRAU R, GONZÁLEZ CM, MORENO AC, GONZÁLEZ M, ALONSO JM, RODRÍGUEZ-MAHOU M, CARREÑO L. Anti-Sa antibodies in patients with systemic lupus erythematosus, (A1) 9

LOPEZ-LONGO FJ See also Alonso JC

LORIG K. Patient education: treatment or nice extra, (E) 703

LOUGHLIN J, IRVEN C, HARDWICK LJ, BUTCHER S, WALSH S, WORDSWORTH P, SYKES B. Linkage of the gene that encodes the al chain of type V collagen (COL5A1) to type II Ehlers-Danlos syndrome, (A2) 40

LOUTHRENOO W. Disseminated penicillinase-producing Neisseria gonorrho in a patient infected with human immunodeficiency virus, (L) 487 Low back pain

acute, preliminary audit of the management in the Kettering district, (L) 693 new episode incidence: results from the south Manchester low back pain

study, (A1) 71 prevalence and patterns of consultation, (A2) 38

Low density lipoprotein

enhanced macrophage degradation from synovial fluid of rheumatoid arthritis patients, (A1) 49

is present in rheumatoid synovial fluid, (A1) 49 Low prevalence of fibromyalgia syndrome in patients attending a sleep

disorders clinic, (A1) 38 Low-dose methotrexate: an effective corticosteroid-sparing agent in the

musculoskeletal manifestations of sarcoidosis, 642 LUKEY PT See Ratcliffe LT

Lumbar epidural steroid injection in sciatica: who benefits?, (A1) 40

LUNEC J See Casey MC; Evans MD; Griffiths HR; Hill CM; Kitas GD

high resolution computed tomography in asymptomatic SLE patients with normal chest radiographs, (A1) 14 interstitial disease, antiendothelial cell antibodies in inflammatory

myopathies, (A1) 19

anticoagulant and anticardiolipin antibodies, systemic lupus erythematosus patients with both are at risk for clinical manifestations related to the antiphospholipid syndrome, 873 environmental syndromes, (E) 597

minocycline-related, (L) 398

sulphasalazine-induced in psoriatic arthritis, 383 Lupus nephritis, anti-Sm antibodies: a potential risk factor for, (A1) 9

Lupus nephropathy, cross-sectional and longitudinal study of anti-ribonucleoprotein antibodies, (A1) 8

Lupus pregnancy

prolactin levels do not differ significantly from those in controls. There is no correlation between disease activity and prolactin levels, (A1) 13 prospective study of 110 high-risk pregnancies in a lupus pregnancy

clinic, (A1) 13

safety of hydroxychloroquine: experience in 27 pregnancies, (A1) 14 Lupus-like syndrome in patients treated for acne, (L) 584

LUOMANI R See Zwede M

LUQMANI RA, ADU D, SAVAGE COS, LOCKWOOD CM, WALDMANN H, PALL A, RICHARDS NT, HOWIE AJ, BACON PA. A comparison of antineutrophil cytoplasmic antibody positive with ANCA negative microscopic polyarteritis, (A1) 67

LUOMANI RA See also McRorie ER: Smith GW

Luteinised unruptured follicles in women with inflammatory arthritis taking NSAIDs, (A2) 22

LYCKLAMA À NIJEHOLT GJ, GUBLER FM, SCHOLTEN R, VAN SCHAARDEN BURG D, DINANT HJ. Subchondral cysts as a favourable prognostic sign in rheumatoid arthritis, (A1) 104

Lyme arthritis and neuroborrelosis, raised levels of IgA and IgM rheumatoid factor, (A1) 110

Lyme borrelosis presenting as a polymyalgia rheumatica-like syndrome, (L) 392

Lyme disease

serological diagnosis, (L) 992

the value of serological testing disease in the UK, 132

Lymphadenopathy

destructive and T-lymphocyte activation in adult-onset Still's disease,

proteinaceous in a patient with known rheumatoid arthritis-case report and review of literature, 1087 Lymphatic hypoplasia, congenital, lymphoedema associated with active

inflammatory arthritis, (L) 689

Lymphocyte adhesion molecule, quantitative analysis of deployment in systemic lupus erythematosus, (A1) 4

Lymphocytes

adhesion to endothelial cells, corticosteroids inhibit in vitro: confirmation using the SV40-transformed EC line LT4, (A1) 146

chemo-attractant effect of rheumatoid synovial fluid, (A1) 57

cutaneous lymphocyte antigen positive, selective accumulation in the skin but not in the joint of inflammatory arthropathies, (A1) 64

reactivity to Clostridium perfringens antigens and toxoid in patients with arthritis, (L) 1095

subsets in pamidronate-induced lymphopenia, (L) 993

Lymphoedema associated with active inflammatory arthritis in a patient with congenital lymphatic hypoplasia, (L) 689

LYNCH JA, MACFARLANE DG, BUCKLAND-WRIGHT JC. Measurement of trabecular organisation of the OA knee by fractal signature analysis of high definition macroradiographs, (A1) 121

LYNCH MP See Dawson JK

LYTTLE K, GRIFFITH SM, MATTEY D, DAWES PT, ROLFE P. Near infra-red laser in the assessment of bone structure and function, (A2) 42

MATSUKAWA Y . Limited association of Helicobacter pylori in gastric ulcer patients with rheumatic disease, (L) 1188

MCALINDON TE See D'Cruz D; Hart DJ

McBurney A See Kitas GD

McCann S See McGonagle D

McCarthy D See Mulherin D McCLOSKEY E See O'Neill TW

McCready VR See Jawad ASM

McCurdie I See Etherington J

MCDERMOTT EM, DEIGHTON CM. Genetic anticipation in familial rheumatoid arthritis, (A2) 6

McDonagh JE, Powell RJ, Walker DJ. Rheumatoid factor detection in a 10 year follow-up study of multicase rheumatoid families, (A1) 88 McDonagh JE, Stephenson AM, Walker DJ. Penetrance and linkage of

HLA and rheumatoid arthritis in extended pedigrees, (A1) 139

McDonagh JE See also Paice E

McDonald A See Morrison E

MACDONALD AG, SYKES H, PHELAN M, THOMPSON RN, FRASER WD. Effect of 2nd line antirheumatic therapy on pyridinium crosslink excretion in patients with established RA, (A1) 99

MACDONALD IA See Gardner-Medwin JMM

McElwain J See Masterson E

McFarlane XA, Robertson DAF, Bhalla AK. Intestinal strontium absorption and axial bone mineral density, (A1) 75

MACFARLANE D See Buckland-Wright JC; Haslock I MACFARLANE DG See Lynch JA

McGibbon D See Doherty M

McGill PE See Logan JA; Njobvu PD

McGonagle D, Costigan T, Hemeryck L, O'Riordan J, McCann S, CASEY EB. Rheumatological complications of bone marrow transplant, (A1) 120

McGonagle D, Donnelly S, Hemeryck L, Ryan R, O'Reilly P, Casey EB. An outbreak of parvovoirus B19 arthritis in North Dublin, (A1) 110 McGuigan L See Helsloot J

MACGREGOR AJ, BAMBER S, CARTHY D, VENCOVSKY J, MAGEED RA, OLLIER WER, SILMAN AJ. Heterogeneity of disease phenotype in monozygotic twins concordant for rheumatoid arthritis, 215

MCHUGH N See Whyte J

MCHUGH NJ See Harvey GR; WHYTE J

MACHIN D See Barry R

MACHIN SJ See McNally T

MACHOLD K See Erlacher L; Koeller M

McINERNEY D See Masterson E

MCINNES IB, CAPELL HA, MADHOK R, HUNTER JA, PULLAR T. Why should rheumatoid arthritis patients recently recruited to sulphasalazine studies have milder disease than those enrolled over the previous decade? Does this trend influence outcome?, (A1) 98

McInnes IB, Field M, Kinninmonth A, Telfer J, Sturrock RD, Liew FY. Nitric oxide production by primary synovial cultures from rheumatoid and osteoarthritis patients, (A1) 44

McInnes IB, FIELD M, STURROCK RD, LIEW FY. Clinical effects of a nitric oxide inhibitor on developing and established collagen induced arthritis in DBA/1 mice, (A1) 64

McInnes IB, Murphy EA, Porter DR, Thompson EA, Capell HA, HUNTER JA, MADHOK R, PULLAR T. Does low dose desensitization affect the toxicity of sulphasalazine in the treatment of rheumatoid arthritis?, (A1) 97

MCKANE JR See Magill MK

MCKANE R See Doherty E

MCKENNA F. Efficacy and gastroduodenal safety of a fixed combination of diclofenae and misoprostol in the treatment of arthritis, (S1) 11 MCKENNA F See also Arnold MM; Pickavance L

MCKENNA SP See Tennant A

MCKILLOP JH See Neilly JB

MACKAY K See Pande I MACKIE IJ See McNally T

MACKWORTH-YOUNG C See Marguerie C

MACKWORTH-YOUNG CG, ANDREOTTI F, HARMER I, LOIZOU S, POTTINGER BE, PEARSON JD, DAVIES GJ, MASERI A, WALPORT MJ. Endotheliumderived haemostatic factors and the antiphospholipid syndrome, 201

MACKWORTH-YOUNG CG. Antiphospholipid antibodies and disease, 1009

McLaren M See Campbell L; Maple C; Veale DJ

MCLAUGHLIN P See McRorie ER McMorrow M See Bunn CC

McNally JA See Bruce IN

McNally T, Mackie IJ, Machin SJ, Isenberg DA. Increased levels of β2 glycoprotein-1 antigen and β 2 glycoprotein-1 binding antibodies are associated with a history of thromboembolic complications in patients with SLE and primary antiphospholipid syndrome, 1031

McNally T See also Menon S Macpherson BCM See Canvin JMG

MACRAE K See Darlington LG

MCRORIE ER, LUQMANI RA, MOOTS RJ, WOLTER D, NAZIR M, EXLEY A, KITAS GD, EMERY P, BACON PA. The value of von Willebrand factor and angiotensin converting enzyme levels in vasculitis, (A1) 22

MCRORIE ER. Lymphoedema associated with active inflammatory arthritis in a patient with congenital lymphatic hypoplasia, (L) 689

McRorie ER, McLoughlin P, Russell T, Nuki G, Hurst NP. Outcome of surgery to stabilise the cervical spine in rheumatoid arthritis—pain, neurological status and patient satisfaction, (A1) 92

Macroamylasemia: a possible cause of unexplained hyperamylasemia in rheumatoid arthritis, 290

Macrophage

enhanced degradation of low density lipoprotein from synovial fluid of rheumatoid arthritis patients, (A1) 49

populations in the synovium: inter-relationships and correlation with articular damage, (A1) 149

Macrophage derived granulocyte macrophage colony stimulating factor may augment HLA-DR expression on inflammatory cells in temporal arteritis, (A1) 151

MADAIO M See Faulds G

MADHOK R See McInnes IB; Morrison E; Neilly JB; Oien KA

MAGEED RA See Adebajo AO; Macgregor AJ
MAGILL MK, MCKANE JR, IRVINE AE, BELL AL. Pathological synovial fluids promote apoptosis of peripheral blood and synovial neutrophils: a role in resolution of synovitis?, (A1) 58

Magnetic resonance images of knee joint: enhancement by advanced 3D computer visualisation, (A1) 123

Magnetic resonance imaging

atlanto-occipital subluxation in rheumatoid arthritis demonstrated by,

of the brain in systemic lupus erythematosus, 1055

characteristics of the cervical spine, canal and cord in rheumatoid arthritis patients with myelopathy, (A1) 124

gadolinium-DTPA enhanced

in rheumatoid knee synovitis: clinical, laboratory and histological correlation, (A1) 75

synovial tissue volume quantification, (A1) 125 use of a contrast enhanced in the assessment of therapeutic response to a disease-modifying antirheumatic drug, 956

MAGNUS JH See Stenstad T

MAHMUD T, RAFI S, SCOTT DL, WRIGGLESWORTH JM, BJARNASON I. Uncoupling oxidative phosphorylation by non-steroidal anti-inflamma-tory drugs related to their pKA, (A2) 23 MAINI RN See Adebajo AO; Brookes SM; Misra R; Rigby S; Venables PJW

MAJOR PJ See Keen RW

Major life events as a result of rheumatoid arthritis. A comparison between nine centres in 446 patients in the first three years of disease, (A2) 9

MAKI-IKOLA O, LEHTINEN K, TOIVANEN P, GRANFORS K. Antibodies to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of ankylosing spondylitis patients with/without iritis and enthesitis, 418

MÄKI-IKOLA O, NISSILÄ M, LEHTINEN K, LEIRISALO-REPO M, TOIVANEN P, GRANFORS K. Antibodies to Klebsiella pneumoniae, Escherichia coli and Proteus mirabilis in the sera of patients with axial and peripheral form of ankylosing spondylitis, 413

Malabsorption caused by coeliac disease in patients who have scleroderma,

MALAISE MG See Kaye O

MALAVIYA AN See Sood A; Uppal SS

MALIK J See Edwards JCW; Sutcliffe N

MALIK R See Gordon C MALIK S See Jones PBB

MALLORIE PA See Kennedy LG Management of gout, (L) 482

in patients with renal insufficiency and a gastrointestinal bleed-an answer to Dr Dunne's letter, (L) 481

Management of oral complications of disease modifying drugs in rheumatoid arthritis, (A2) 20

MANGER B See Kalden JR MANGRAVITI S See Cutolo M

Manipulative physiotherapy for spinal problems in primary care: outcomes of care, 1070

MANNING J See Bukhari M: Teh LS

Mannose binding protein

and complement C4 polymorphism are associated with SLE in a Spanish population, (A1) 78

and systemic lupus erythematosus in southern Chinese, (A1) 7 MANTHORPE R See Wylie G

MAPLE C, BELCH JJF. Dietary essential fatty acid supplements reduce red

blood cell aggregation in healthy volunteers, (A1) 28
MAPLE C, KIRK G, MCLAREN M, BELCH JJF. A diurnal variation in soluble E-selectin, P-selectin and ICAM-1 levels in healthy volunteers, (A1) 87

MAPLE C, KIRK G, VEALE D, BELCH JJF. Reduction in cell adhesion molecule levels following sulphasalazine in patients with rheumatoid

arthritis, (A1) 98 MAPP PI See Winrow VR

Mapping the genetic association between hypergammaglobulinaemic primary Sjögren's syndrome and the TRCB gene complex, (A1) 77

MARCOLONGO R See Wylie G

Marfan syndrome

assessment of bone mineral density in women, 516

joint and skin manifestations are prevalent and correlated, 126

MARGISON GP See Rafferty JA

MARGUERIE C, KAYE S, VYSE T, MACKWORTH-YOUNG C, WALPORT MJ, BLACK C. Malabsorption caused by coeliac disease in patients who have scleroderma, 858

MAROTEAUX P See Bailey AJ

MARSAL S, FALGA C, SIMEON CP, VILARDELL M, BOSCH JA. Behçet disease and pregnancy relationship study, (A1) 24

MARSHALL KW, GUTHRIE BT, MIKULIS DJ. Quantitative cartilage imaging, (S1) 29

MARTIN JC, ENGLAND A, REID DM. Can changes in appendicular bone predict axial bone mineral density during treatment with hormone replacement therapy and cyclical etidronate? (A2) 41

MARTIN K, BOND A, HAY FC, AXFORD JS. Oligosaccharide changes in rheumatoid arthritis determined by fluorophore assisted electrophoresis and lectin binding analysis, (A1) 63

MARTIN K, MULHERIN D, FITZGERALD O, AXFORD JS. Fluorophore assisted carbohydrate electrophoresis: a potential tool for the identification of arthritic diseases, (A1) 61

MARTIN S See Voskuyl AE

MARTÍN-VILLA JM See Pablos JL

MARTINEZ DE OSABA MJ See Peris P

MARTINI A. See Massa M

MARTYN CN See Dunne CA

MASALANKA W See Patel S

MASERI A See Mackworth-Young CG

MASON R See Lewis S

MASSA M, DE BENEDETTI F, PIGNATTI P, ALBANI S, MONESTIER M, MARTINI A. Lack of temporal association of iridocyclitis with IgG reactivities to core histones and nucleosome subparticles in pauciarticular juvenile chronic arthritis, 507

Massive pericardial effusion in scleroderma: a review of five cases, 564 Mast cell tryptase in the synovial fluid of patients with arthritis, (A1) 56

MASTERSON E, MULCAHY D, MCELWAIN J, McINERNEY D. The planovalgus rheumatoid foot-is tibialis posterior tendon rupture a factor?, 645

MATEO I See Gonzalez-Crespo MR

Maternal HLA antigens and antibodies to SS-A/Ro and SS-B/La. Comparison with systemic lupus erythematosus and primary Sjögren's syndrome, 901

MATHEWS JA. Acute calcific discitis. A report of two cases with scintigraphy and SPECT, (A2) 55

MATHEWS JA See also Etherington J

MATSUKAWA Y. Abducent and trochlear palsies in a patient with Sjögren's syndrome, (L) 484 MATTEY D See Borg AA; Cheung NT; Evans E; Hassell A; Lyttle K; Nixon

NB; Salih AM

MATTEY DL, NIXON NB, SLATER H, DAWES PT. TGF-β1 induces myofibroblastic differentiation in human synovial fibroblasts in vitro: modulation by basic FGF, (A2) 29 MATTEY DL See also Cheung NT; Close DR; Russell G

MAUGARS Y See Berthelot JM

MAUS S See Stucki G

MAYER B, HASSELL AH, CARPENTER EH, HOTHERSALL TE, SHADFORTH MF, HAY E, DAWES PT. Routine follow up of RA patients: what dictates periodicity of review?, (A2) 12

MEADOWS K See Dodd J

Measurement

of antibodies to reactive oxygen species damaged DNA as a simple marker of disease activity in SLE, (A1) 10

of articular cartilage compressive modulus under physiological loading conditions, (A2) 48

of nitric oxide in plasma and synovial fluid using a novel ESR-spin trapping technique, (A1) 45

of pain in rheumatoid arthritis, (A1) 91

of shoulder movement using FASTRAK 3-D tracking system, (A2) 47 of trabecular organisation of the OA knee by fractal signature analysis of high definition macroradiographs, (A1) 121

Measuring outcome in rheumatoid arthritiswhich measures are suitable for routine clinical use?, (E) 802

Medial joint space narrowing and meniscal lesions, radiological, (L) 991 Median nerve conduction, validation of a new method of assessing at the wrist in carpal tunnel syndrome, (A1) 43

Medical education

continuing for rheumatologists: the current picture, 172 continuing in rheumatology, (E) 98

Medical factors, life-style and environmental factors influencing peak bone mass in women, 620

Medical practice, evidence-based: the Cochrane collaboration and osteoarthritis, (E) 403

MEDINA F, FUENTES M, JARA LJ, BARILE L, MIRANDA JM, FRAGA A. Salmonella pyomyositis in patients with the human immunodeficiency virus, 568

MEDINA F. Staphylococcal arthritis in human immunodeficiency virus infection, (L) 397

MEHRA NK See Uppal SS MEIJERS P See Dobbie JW

MEJIA A See Donnelly S

MELBY K See Glennås A

MELCHERS I See Voskuyl AE

MELIEF MJ, HOIJER MA, VAN PAASSEN HC, HAZENBERG MP. Presence of bacterial flora-derived antigen in synovial tissue macrophages and dendritic cells, 1112

Meloxicam: a preferential inhibitor of COX-2, (A1) 48

Meningitis, primary nocardial in systemic lupus erythematosus, 178

Meniscal lesions, and medial joint space narrowing, radiological, (L) 991

MENKES CJ. Jaques Villiaumey, obituary, 592

MENON S, ISENBERG DA. Fetal calf serum in growth medium obscures the detection of early IgG antocardiolipin-antibody secreting clones, (A1) 12 MENON S, RAVIRAJAN C, ISENBERG DA. An analysis of thrombotic events in antiphospholipid antibody positive lupus patients, (A1) 15

MENON S, RAVIRAJAN C, McNally T, ISENBERG DA. The production and analysis of the binding characteristics of four human monoclonal IgG anticardiolipin antibodies, (A1) 12

MENON S See also Rahman MAA; Tucker LB

MENSCHIK M See Erlacher L

Menstrual disorders in girls with systemic lupus erythematosus treated with cyclophosphamide, 737

Mercuric chloride-induced autoimmunity, oxypentifylline enhances the Th2 environment and inhibits TNFa mRNA production and inflammatory arthritis, (A2) 29

MERINO R See Gonzalez-Crespo MR

Metacarpophalangeal prosthesis, design of a non constrained, non cemented, modular (A2) 47

Metal-ion chelating hydroxyl radical scavenger, novel transition, EDTA bis(ethyl phenylalaninate): potential anti-inflammatory role, (A1) 47 Metalloproteinas

neutral matrix in scleroderma, (A1) 29

release and cyclical pressure from human articular cartilage, (A2) 28 Methodological problems in comparing the severity of rheumatoid arthritis between populations, 781

Methotrexate

action in rheumatoid arthritis: stimulation of cytokine inhibition and inhibition of chemokine production by peripheral blood mononuclear cells, 602

adverse events in treated rheumatoid arthritis patients, (S2) 49

alone or in combination with sulphasalazine, effects on the production and circulating concentrations of cytokines and their antagonists, 747

efficacy in rheumatoid arthritis, (S2) 43

and flucloxacillin, lack of interaction between in patients with rheumatoid arthritis, (A1) 100

and folate supplementation, 1172

and folic acid, an audit of for rheumatoid arthritis. Experience from a teaching centre, 971

liposomally conjugated effects upon mediator release from human peripheral blood monocytes, 241

low dose: an effective corticosteroid-sparing agent in the musculoskeletal manifestations of sarcoidosis, 642

dose, practical clinical pharmacology and drug interactions in rheumatoid arthritis, (S2) 20

possible mechanisms of action in patients with rheumatoid arthritis, (S2)

pulmonary function in rheumatoid arthritis treated with low dose: a longitudinal study, (A1) 100

and sulphasalazine antirheumatic agents share an anti-inflammatory mechanism, (S2) 30

and sulphasalazine combination in the management of rheumatoid arthritis—view based on personal clinical experience, (S2) 109

and sulphasalazine combination in rheumatoid arthritis: early clinical impressions, (S2) 104 total and free pharmacokinetics, with and without piroxicam, in

rheumatoid arthritis patients, 421 Methylprednisolone, intramuscular, controlled trial against oral prednisolone

in polymyalgia rheumatica: a 48 week follow up, (A1) 69 Metrology index, ankylosing spondylitis: the correlation between a new

metrology score and radiology, 767 MEVORACH D. Management of gout, (L) 482

MEYER O See Combe B

MEYERS OL See Kalla AA; Ratcliffe LT

MHC encoded TN-X gene may be important in RA, (A1) 140

MICHAEL J See Luqmani RA

MICHEL BA See Stoll T; Stucki G

Microwave thermography evaluation of the hand and wrist in rheumatoid arthritis, (A2) 13

MIELANTS H, VEYS EM, CUVELIER C, DE VOS M, SCHATTEMAN G, GYSELBRECHT G. Risk factors for a patient with spondylarthropathy to develop ankylosing spondylitis or inflammatory bowel disease, (A1)

MIELANTS H, VEYS EM, SCHATTEMAN, G, GYSELBRECHT G, ELEWAUT D. Influence of sulphasalazine on the evolution of gut inflammation and locomotor manifestations in the spondylarthropathies, (A1) 115
MIHARA M, MORIYA Y, KISHIMOTO T, OHSUGI Y. Interleukin-6 (IL-6)

induces the proliferation of synovial fibroblastic cells in the presence of soluble IL-6 receptor, 321

MIJIYAWA M. Gout in patients attending the rheumatology unit of Lomé Hospital, 843

MIKECZ K See Brennan FR

MIKULIS DJ See Marshall KW

MILFORD-WARD A See Hill MB

MILLER F See D'Cruz D

MIMORI T See Satoh M

MINGUEZ C. New case of polyarthritis as a presenting sign of non-Hodgkin's lymphoma, (L) 794

MINGUEZ M See González CM

MINGUEZ MD See Abad MA; Monteagudo I; Moreno FC

Minocycline

double blind placebo controlled trial in patients with active rheumatoid arthritis, (A2) 16

effect on acute phase proteins and serum interleukin-6 levels in rheumatoid arthritis patients, (A1) 96

Minocycline-related lupus, (L) 398

MIRANDA C See Cury SE

MIRANDA JM See Medina F

Misoprostol and diclofenac fixed combination, efficacy and gastroduodenal safety of in the treatment of arthritis, (S1) 11

MISRA R, DARTON K, JEWKES RF, BLACK CM, MAINI RN. Arthritis in scleroderma, 831

MISRA R See also Agrawal S MITCHELL N See Murray JG

MITRA D See Elvins DM

MITROVIC DR See Pejovic M

MITUSZOVA M See Kiss Cs

Mixed connective tissue diseas does it exist in India?, (A2) 26

in India?, 539

Mixed cryoglobulinaemia, inner ear involvement, 370

MIYACHI S See Shingu M

MIYAZAKI K See Satoh M

Modulation of scleroderma and control fibroblast phenotype by co-culture with human endothelial cells, (A1) 66 MOK CC, LAU CS, POON SP. Primary nocardial meningitis in systemic lupus

erythematosus, 178 Mok CC See also Lau CS

Molecular basis of hereditary complement factor I deficiency, (A1) 155

Molecular markers to monitor outcome and intervention in osteoarthritis (promises, promises...), (L) 599

MOLINARI AC See Gattorno M

MOLLAN RAB. See Kerr JR MONEGAL A See Peris P

MONESTIER M See Massa M MONIZ C See Dolan AL

Monoclonal antibodies, therapeutic, 707

Monocytes

effect of gold salts and their pattern of cytokine secretion, (A1) 150 peripheral blood, effect of liposomally conjugated methotrexate upon mediator release, 241

Mononuclear cells, increased IL-1 receptor antagonist (IL-1ra) production and decreased IL-1B/IL-1ra ratio from rheumatoid arthritis patients, 24

Monozygotic twins, heterogeneity of disease phenotype is concordant for rheumatoid arthritis, 215

MONTALVO G See Pablos JL

MONTEAGUDO I, GRAU R, GONZÁLEZ M, MORENO AC, ABAD MA, MINGUEZ MD, LÓPEZ-LONGO FJ, ARIZA A, CARREÑO L. Antibodies to intermediate filaments and human immunodeficiency virus, (A1) 112

MONTEAGUDO I See also Abad MA; González CM; Moreno FC

MONTGOMERY B See Carpenter EH

MOORE AR See Willis D

MOORE D See Nisar M

MOORE JE See Kerr JR

MOORE T See Bukhari M; Teh LS MOORE TL See Anderson ME

Моотоо R See Etherington J

MOOTS R.I. See McRorie F.R.

MORENO AC, LÓPEZ-LONGO FJ, GRAU R, MONTEAGUDO I, GONZÁLEZ CM, GONZÁLEZ M, MÍNGUEZ MD, ABAD MA, CARREÑO L. Immunoblot analysis of precipitating and non-precipitating anti-ribonucleoprotein antibodies, (A1) 11

MORENO AC See also Abad MA; González CM; González M; López-Longo FJ: Monteagudo I

MORGAN S See Pal B

MORGAN SL See Sandoval DM

MORI PG See Gattorno M MORIYA Y See Mihara M

MORLEY BJ See Bartók I; Hogarth MB; Vaishnaw AK

MORLEY KD See Veale DJ

MORREALE G See Gattorno M MORRIS C See Naughton D

MORRIS CJ See Dai L; Jones HW

MORRISON E, MUNRO R, MCDONALD A, CAPELL HA, MADHOK R, HUNTER JA. Can we differentiate between radiological progression over one year in patients with rheumatoid arthritis randomly allocated to im gold or hydroxychloroquine?, (A1) 107

MORRISON E See also Logan JA

MORRISON NA See Keen RW

MORTON MJS See Baildam EM

MORTUZA FY See Perry MJ MOUTSOPOULOS HM See Boumba D

MOYNIER M See Jorgensen C

MUIR KR See O'Reilly SC

MUJIC F, CUADRADO MJ, LLOYD M, KHAMASHTA MA, HUGHES GRV. Development of systemic lupus erythematosus features in patients with primary antiphospholipid syndrome, (A1) 16

MUJIC F, LLOYD M, CUADRADO MJ, KHAMASHTA MA, HUGHES GRV. Prevalence and clinical significance of subungual splinter haemorrhages in patients with an antiphospholipid syndrome, (A1) 16

MULCAHY D See Masterson E

MULCAHY H See Mulherin D

MULEY R See Gonzalez-Crespo MR

MULHERIN D, FITZGERALD O, BRESNIHAN B. An analysis of the discrepancy between the clinical and radiological course in rheumatoid arthritis, (A1)

MULHERIN D, FITZGERALD O, BRESNIHAN B. Macrophage populations in the synovium: inter-relationships and correlation with articular damage, (A1) 149

MULHERIN D, MULCAHY H, SKELLY M, SAUNDERS A, DONOGHUE DO, MCCARTHY D, FITZGERALD O, BRESNIHAN B. Identifying iron deficiency anaemia by simple laboratory measures in patients with rheumatoid arthritis, (A1) 83

MULHERIN D, VEALE DJ, BELCH JJF, BRESNIHAN B, FITZGERALD O. Relationship of ICAM-1 with E-selectin in serum, synovial fluid and synovial membrane in inflammatory arthritis, (A1) 148

MULHERIN D See also Breathnach D; Doherty E; Martin K

MULLIGAN P See De Lord D

Multiple epiphyseal dysplasia, linkage analysis in, (A1) 144

Multiple joint avascular necrosis: beware of tuberculosis and human immunodeficiency virus-a rare but important cause, 387

MUÑOZ-GOMEZ J See Peris P; Pons F

MUNRO R, ARMSTRONG S, LAND D, STURROCK RD. Evaluation of microwave thermography of the hand and wrist in rheumatoid arthritis,

MUNRO R See also Morrison E

MURPHY A See Donnelly S

MURPHY EA See Karim SN; McInnes IB

MURPHY K, WESTON A, BRADLOW A, DAVID J. Intra-muscular versus intraarticular depo-methyle-prednisolone in the management of ambulatory patients with active rheumatoid arthritis, (A1) 102

MURRAY JG, RIDLEY N, MITCHELL N, WOO P, ROONEY M. Hip evaluation in juvenile chronic arthritis evaluation with contrast-enhanced MR imaging, (A1) 124

MURRAY PI See Yeo LMW

Muscle abscess, primary tuberculous in a patient with systemic lupus erythematosus, 1177

Muscle strength

in healthy people and in patients suffering from recent-onset inflammatory arthritis, 355

measurement, new quantitative and simple: interobserver reliability in healthy subjects, (A1) 20

voluntary activation and cross-sectional muscle area in patients with fibromyalgia, 925

Musculoskeletal complaints in patients with hyperlipoproteinaemia, (L) 393 Musculoskeletal pain importance in patients with irritable bowel syndrome, (A1) 119

Musculoskeletal problems in general practice: a prospective survey of consultations and management, (A2) 50

Mycobacterial stress protein 65 kDa, antibodies to in West Africans with rheumatoid arthritis, tuberculosis and malaria, 352

Mycoplasmal RNA and 5'-nucleotidase in the synovial fluid of rheumatoid arthritis patients, (A2) 5 Myeloperoxidase, autoantibodies to in systemic sclerosis, (A2) 24

MYLES AB. Non-surgical management of the carpal tunnel syndrome, (L) 890

Myofibroblastic differentiation, TGF-β1 induced in human synovial fibroblasts in vitro: modulation by basic FGF, (A2) 29

Myopathy incidence in patients treated with antimalarials. A report of three cases and a review of the literature, 166

Nabumetone, perforations, peptic ulceration and bleeding on NSAIDs. Two randomised controlled trials of nabumetone, diclofenac SR and piroxicam in the UK, (A1) 95

NADDAF HO See Al Rikabi AC

NAGAI Y See Shingu M

NAIK S See Agrawal S

Nail fold rheumatoid vasculitis, isolated-a benign condition?, (A1) 68

NAKIELNY RA See Romanowski CAJ

NANDRA D See Carr AJ; Etherington J; Hart DJ; Keen RW; Lloyd ME; Spector TD

NARAYAN P. Preliminary audit of the management of acute low back pain in the Kettering district, (L) 693

NASH MC, SHAH V, READER JA, DILLON MJ. Anti-neutrophil cytoplasmic antibodies and anti-endothelial cell antibodies are not increased in Kawasaki disease, 882

NASH P See Kavanagh R

Nature of surgical intervention in patients with ankylosing spondylitis, (A1)

NAUGHTON D, RHODES C, GAFFNEY K, BLAKE DR, GROOTVELD M. Detection of nitrosyl haemoglobin in isolated synovial fluid and blood

of rheumatoid patients, (A1) 46
NAUGHTON D, TRENNAM C, MORRIS C, BLAKE DR, GROOTVELD M. EDTA bis(ethyl phenylalaninate): a novel transition metal-ion chelating hydroxyl radical scavenger with a potential anti-inflammatory role, (A1) 47

NAUGHTON M, BOTTO M, WURZNER R, WALPORT MJ. Where is C7 produced?, (A1) 155

NAZHAT NB, FAIRBURN K, SAADALLA-NAZHAT RA, WINYARD PG, JONES P, BLAKE DR. Measurement of nitric oxide in plasma and synovial fluid using a novel ESR-spin trapping technique, (A1) 45

NAZHAT NB See Zhang Z

NAZIR M See McRorie ER

Near infra-red laser in the assessment of bone structure and function, (A2) 42 Necrotizing vasculitis, acute (polyarteritis nodosa?) confined to the nerve with spontaneous recovery, (L) 694

Needle arthroscopy, (E) 1102 NEILLY JB, LIM KL, HOLLMAN AS, BEHAN WMH, MADHOK R, STURROCK RD, McKillop JH. Indium-111 antimyosin antibody and muscle ultrasound scans in the evaluation of patients with suspected polymyositis, (A2) 27

NEILLY JB. Scintigraphic findings in synovial chondromatosis at presentation, (L) 687

Neisseria gonorrhoeae, disseminated penicillinase-producing in a patient infected with human immunodeficiency virus, (L) 487

NELSON JL See Østensen M

NELSON PN. Absence of exogenous retroviruses in Felty's syndrome, (L) 185 NELSON S See Field M

Nephrotoxicity, chronic in the treatment of arthritis, 162

Neuroborrelosis and Lyme arthritis, raised levels of IgA and IgM rheumatoid factor, (A1) 110

Neurogenic contribution to bradykinin-induced intra-articular plasma extravasation in the rat knee joint, (A1) 52

Neurological involvement in systemic slerosis, (E) 1007

Neurological side effects in two patients receiving gold injections for rheumatoid arthritis, 989

Neurophysiological assessment, how does it influence the management and outcome of patients with carpal tunnel syndrome?, (L) 581

Neurovascular control mechanisms in primary Raynaud's phenomenon and systemic sclerosis as assessed by iontophoresis and laser Doppler techniques, (A2) 24

Neutral matrix metalloproteinases in scleroderma, (A1) 29

Neutrophil adhesion, prolonged to endothelial cells, hypoxia induces at physiological shear rates, (A1) 50 Neutrophil superoxide protection in RA, sulphasalazine effect, (A2) 14

Neutrophil-derived tumour necrosis factor-α: a potential mechanism for promotion of inflammation in procainamide-induced lupus, (A1) 3

New case of polyarthritis as a presenting sign of non-Hodgkin's lymphoma,

New episode incidence of low back pain: results from the south Manchester low back pain study, (A1) 71

New quantitative and simple muscle strength measurement: interobserver reliability in healthy subjects, (A1) 20 New therapeutic approaches in autoimmune rheumatic diseases, with special

emphasis on rheumatoid arthritis, (E) 193

NEWHAM DJ See Hurley MV

NEWMAN J. BAMBER S. SILMAN AJ. Patients with rheumatoid arthritis not attending a specialist clinic have similar levels of disease severity as clinic attenders, (A1) 88

NEWTH J See Thompson PW

NICHOL FE See Evans MD NIELSEN KD See Drewes AM

NIKKARI S See Roivainen A

NILSSEN S See Brun JG

NISAR M. Cyclosporin A in the treatment of rheumatoid associated pyoderma gangrenosum, (L) 182

NISAR M, ROMANOWSKI C, MOORE D. MRI characteristics of the cervical spine, canal and cord in rheumatoid arthritis patients with myelopathy, (A1) 124

NISAR M See also Romanowski CAJ

NISSILÄ M See Mäki-Ikola O

Nitrate concentrations, elevated serum in active spondylarthropathies, 616 Nitric oxide

inhibitor, clinical effects on developing and established collagen induced arthritis in DBA/1 mice, (A1) 64

measurement in plasma and synovial fluid using a novel ESR-spin trapping technique, (A1) 45

production, cellular sources in the joint, (A1) 44

production by primary synovial cultures from rheumatoid and osteoarthritis patients, (A1) 44

Nitrosyl haemoglobin detection in isolated synovial fluid and blood of rheumatoid patients, (A1) 46

NIXON NB, SALIH AM, GAGAN RM, MATTEY D. DAWES PT. Presence of synovial fluid glycolipid antibodies in rheumatoid arthritis-synovial neuropathy or synovial damage, (A2) 4

NIXON NB See also Mattey DL; Salih AM

NJOBVU PD, JELLIS JE, McGILL PE. Arthritis associated with HIV infection in Zambia, (A2) 32

1-NMR spectroscopy, polymyositis patients show consistent changes, (A1) 125

NOBBS BT See Darlington LG

NOBUNAGA M See Shingu M

Nocardial meningitis, primary in systemic lupus erythematosus, 178

NOEL E See Tebib JG

NOJIMA T See Fujimoto M

Non-Hodgkin's lymphoma

new case of polyarthritis as a presenting sign, (L) 794 and polyarthritis, (L) 293

presenting as polyarthritis, (L) 182

Non-steroidal anti-inflammatory drug therapy

iatrogenic cost, (S1) 19

patient activity as an objective measure: an Arthrotec and NUMACT study, (A2) 15

Non-steroidal anti-inflammatory drug-induced gastrointestinal damage. uncoupling of oxidative phosphorylation is the initiating factor, (A1) 93 Non-steroidal anti-inflammatory drugs

characteristics of ulcers of the small bowel induced by in the rat: implications for clinical practice, 727

chondrodestructive and chondrosparing properties in a model of destructive mono-articular arthritis, (A2) 15

double-blind placebo-controlled study of ranitidine for the prevention of recurrent peptic ulcer disease in rheumatoid arthritis patients, (A1)

epidemiology and prevention of effects in the gastrointestinal tract, (S1) 5 historical overview, (S1) 2

luteinised unruptured follicles in women with inflammatory arthritis, (A2) 22

perforations, peptic unceration and bleeding. Two randomised controlled trials of nabumetone, diclofenac SR and piroxicam in the UK, (A1) 95

and renal function, (S1) 25

uncoupling oxidative phosphorylation by non-steroidal anti-inflammatory drugs related to their pKA, (A2) 23

Non-steroidal anti-inflammatory prescribing, decisions in, (S1) 1 Non-surgical management of the carpal tunnel syndrome, (L) 890

NONAKA S See Shingu M

Norfolk Arthritis Register, survival on sulphasalazine, (A2) 14

NØRREGAARD J, BÜLOW PM, VESTERGAARD-POULSEN P, THOMSEN C,

DANNESKIOLD-SAMSØE B. Muscle strength, voluntary activation and cross-sectional muscle area in patients with fibromyalgia, 925

NORSWORTHY P See Davies KA; Hogarth MB

NOTARIANNI LJ See Collins AJ

NUKI G See Brennan FR; Jobanputra P; McRorie ER; Roberts R; Wright M NUMACT: a novel method of monitoring physical activity in JCA, (A2) 23 Number of tender points and extent of reported pain, (A1) 38

ØSTENSEN M, NELSON JL. Bits and pieces in a puzzle-rheumatoid arthritis and pregnancy, (E) 1

Obesity, occupational activity and osteoarthritis of the spine, (A1) 35 Obituar

AT Richardson, 188

Dr W. Carson Dick, 998 Jaques Villiaumey, 592 Oswald Arthur Savage, 591

O'CONNELL D See Flanagan FL O'CONNELL S See Rees DHE

Occurrence, nature and distribution of flares in a cohort of patients with systemic lupus erythematosus: a rheumatological view, 257

OGAWA S See SATOH M O'HARA PS See Helliwell PS

OHSUGI Y See Mihara M

OHTA A See Kuroki S OIEN KA, BLACK A, HUNTER JA, MADHOK R. Pneumocytis carinii pneumonia in a patient with rheumatoid arthritis, not on immunosuppressive therapy and in the absence of human immunodeficiency virus

infection, 677 Oligoarthritis

screening of synovial fluid samples for bacterial antigens using PCR,

seronegative, Salmonella-a forgotten infection?, (A2) 32

undifferentiated, reactive arthritis: urogenital swab culture is the only useful diagnostic method for the detection of the arthritogenic infection in extra-articularly asymptomatic patients, 838

Oligosaccharide changes in rheumatoid arthritis determined by fluorophore assisted electrophoresis and lectin binding analysis, (A1) 63

OLIVEIRA DBG See Kiely PDW OLLIER W See Donn RP; Hassell A; Thomson W

OLLIER WE See Farhan AJ

OLLIER WER, ALLDERSEA J, DAVIES E, FRYER A, SNOWDON N, JONES P, STRANGE R. Homozygosity for glutathione-transferase GSTM1 and GSTT1 null alleles may confer risk for anti-Ro antibodies in SLE, (A1) 5 OLLIER WER See also Brennan P; Davies E; Davies EJ; Donn RP; Howarth

S; Kay RA; Macgregor AJ; Poulton K OM-89

bacterial extract, effects of tolerance and sensitization to, with a possible mode of action in polyarthritis in the rat, 1135

modulation of chronic inflammation: relevance to clinical use, 525 Omeprazole and small bowel bacterial overgrowth in patients with systemic sclerosis, (A1) 67

Omeprazole-induced small bowel bacterial overgrowth in patients with scleroderma, 976

Oncogene expression in synovial fluid cells in reactive and early rheumatoid arthritis: a brief report, 805

O'NEIL H See Jobanputra P

O'NEILL T See Kiss Cs

O'NEILL TW. McCloskey E. Kanis JA. Reid DM. Reeve J. Todd C. WOOLF AD. The epidemiology of vertebral osteophytosis, (A1) 70 O'NEILL TW See also Elvins DM

OOSTENDORP RAB See Hogeweg JA

Oral complications of disease modifying drugs, management in rheumatoid arthritis, (A2) 20

Oral steroids as bridge therapy in rheumatoid arthritis patients starting with parenteral gold. A randomized double-blind placebo-controlled trial, 347

inflammatory, Churg-Strauss syndrome, (L) 485

a new ocular entity in RA?, (A2) 55

ORDI J See Davies E

O'REILLY D See Teh LS

O'REILLY P See McGonagle D

O'REILLY SC, JONES AC, MUIR KR, DOHERTY M. Knee pain, quadriceps weakness and muscle activation in the community, (A2) 45

O'REILLY SC, MUIR KR, DOHERTY M. Estimates of knee pain and disability in the community with differing knee pain criteria, (A1) 36

O'REILLY SC, MUIR KR, DOHERTY M. Knee pain, disability and psychological variables in a Nottingham population, (A1) 35

O'RIORDAN J See McGonagle D

Osteoarthritis

and the Cochrane collaboration, evidence-based medical practice, (E) 403

determination of the apparent synovial permeability in the knee joint,

diclofenac is a better analgesic than paracetamol in patients with osteoarthritis of the hip, knee and lumbar spine, (A1) 137

diurnal variation in 24 hour ambulatory recordings of patient activity is different from rheumatoid arthritis. (A2) 16

knee

determinants of arthrogenic inhibition, (A1) 134

does spiking of the tibial tubercles predict?, (A1) 36

joint space width measured more accurately and precisely in semiflexed than extended view, (A1) 121

measurement of trabecular organisation by fractal signature analysis of high definition macroradiographs, (A1) 121

pain mechanisms in: effect of intra-articular anaesthetic, (A2) 45 psychological distress and local tenderness but not radiographic change or weakness are associated with disability in hospital referred patients, (A1) 136

shortly after anterior cruciate ligament transection dogs develop characteristics similar to human, (A1) 135

symmetrical compartmental involvement, (A1) 134

tiaprofenic acid, indomethacin and placebo in: the long term outcome of therapy, (A2) 44 levels of collagenase, tissue inhibitor of metalloproteinase (TIMP),

collagenase-TIMP complexes and glycosaminoglycan in sequential samples of synovial fluid aspirated from patients, (A1) 138

of the lumbar spine, utility of lateral bone density measurements, (A1) 127

molecular markers to monitor outcome and intervention in (promises, promises...), (L) 599

nitric oxide production by primary synovial cultures, (A1) 44

nodal, evidence for genetic anticipation, (A1) 133

and osteophytosis, relationship in joints, (A1) 132

and Paget's disease, (A2) 46, 1061

pattern of distribution of the hand joints: a comparative study between a clinical and a community cohort, (A1) 133

primary generalised, prevalence of xerostomia in a population, (A1) 132 relationship between serum cartilage oligomeric matrix protein levels and disease progression in the knee joint, 306

relationship with IGF-1 in women aged 45-64 in the general population: the Chingford study, (A1) 137

risk associated with running: a radiological survey of female ex-athletes and population controls, (A1) 70

spinal and osteoporosis: evidence for a negative association independent of obesity, (A1) 74

of the spine, obesity and occupational activity, (A1) 35

Osteophyte, tibiofemoral in the anteroposterior radiograph predicts joint space narrowing in the patellofemoral joint, (A1) 136

Osteophytosis

epidemiology of vertebral, (A1) 70

and osteoarthritis, relationship in joints, (A1) 132

Osteoporosis

aetiology and presenting symptoms in males, 936

corticosteroid, 8

does vasectomy predispose to?, (A2) 42

identification of an at risk group using heel ultrasound measurement in general practice, (A1) 37

idiopathic juvenile: experience of twenty-one patients, 68

postmenopausal, polymorphism in an Sp1 consensus binding motif in the first intron of collagen I alpha 1 is related to, (A1) 78

randomized controlled trial of brisk walking in the prevention of and of and falls, (A1) 72

should close relatives of patients be investigated?, (A1) 73

and spinal osteoarthritis: evidence for a negative association independent of obesity, (A1) 74

transient regional of the hip, (L) 296

Outbreak of parvovoirus B19 arthritis in North Dublin, (A1) 110

conceptualizing and defining, (E) 899

of surgery to stabilise the cervical spine in rheumatoid arthritis-pain, neurological status and patient satisfaction. (A1) 92

of total hip replacement in ankylosing spondylitis: a thirteen year followup study, (A1) 114

Outpost rheumatology: the general practitioners' perspective, (A2) 50

Overlap syndromes, antiphospholipid antibodies in, 873

Overview of combination second-line or disease-modifying antirheumatic drug therapy in rheumatoid arthritis, (\$2) 96

Oxidative inactivation of human alpha-1-proteinase inhibitor by simultaneous generation of superoxide and nitric oxide, (A1) 48

Oxidative phosphorylation uncoupling by non-steroidal anti-inflammatory drugs related to their pKA, (A2) 23 is the initiating factor in NSAID-induced gastrointestinal damage, (A1)

Oxidised LDL is present in rheumatoid synovial fluid, (A1) 49

OXTOBY J See Averns HL

Oxypentifylline enhances the Th2 environment and inhibits TNFa mRNA production and inflammatory arthritis in mercuric chloride-induced autoimmunity, (A2) 29

PEARCE D. Urinary levels of pyridinoline and deoxypyridinoline may be influenced by turnover of non-osseous tissues in patients with joint disease, (L) 1186

P-selectin, E-selectin and ICAM-1 diurnal variation in levels in healthy volunteers, (A1) 87

PABLOS JL, CARREIRA PE, MARTÍN-VILLA JM, MONTALVO G, ARNAIZ-VILLENA A, GOMEZ-REINO JJ. Polymorphism of the heat-shock protein gene HSP70-2 in systemic lupus erythematosus, 721 Paediatric rheumatology, BPA report, 60

PAGEORGIOU A See Croft PR

Paget's disease

of bone, changes in bone mineral density with bisphosphonate treatment, (A1) 130

and osteoarthritis, (A2) 46, 1061

PAICE E, McDonagh JE. Anterior knee pain in thalassaemia major treated with L1, a new iron chelating agent, (A1) 119

PAIMELA L, PALOSUO T, LEIRISALO-REPO M, HELVE T, AHO K. Prognostic value of quantitative measurement of rheumatoid factor in early rheumatoid arthritis, 1146

Pain

the body chart: a meaningful assessment of pain compared to visual analogue scales in ankylosing spondylitis, (A2) 36 can we isolate the EEG activity related to on-going pain in the brain?,

measurement in rheumatoid arthritis, (A1) 91

mechanisms in osteoarthritis of the knee: effect of intra-articular anaesthetic. (A2) 45

reported, extent and number of tender points, (A1) 38

sensation, consequences of sustained activation of small sensory fibres, (A1) 53

single photon emission computerised tomography identifies sites in ankylosing spondylitis, (A1) 126

threshold in juvenile chronic arthritis, 61

Painful knees associated with a septic focus, 862

Pakistan, prevalence of rheumatoid arthritis in affluent and poor urban communities, 252

PAL B, MORGAN S. An assessment of primary Sjögren's syndrome in patients with dry eyes, (A2) 25

PALAZZO E See Kave O

PALEY M See Clunie G

PALL A See Luqmani RA

PALLIS M, LIM K, ROBINS RA, POWELL RJ. Quantitative analysis of lymphocyte adhesion molecule deployment in systemic lupus erythematosus, (A1) 4

PALMER W See Spitzer WO

PALOSUO T See Paimela L

Pamidronate and uveitis, (L) 479

Pamidronate-induced lymphopenia, lymphocyte subsets, (L) 993

PANAYI G See Bajocchi GL; Dasgupta B; Pitzalis C; Robinson C

PANAYI GS See Awad M; Campbell F; Cauli A; Choy EHS; Dolan AL; Kassimos D; Kassimos DG; Pitzalis C; Rankin ECC

Pancreatitis, panniculitis and arthritis, 680

Pancytopenia, toxic epidermal necrolysis and adult respiratory syndrome, (L)

PANDE I, MACKAY K, CHATFIELD C, CALIN A. The Bath Ankylosing Spondylitis Radiology Index (BASRI): a new validated approach to disease assessment, (A2) 37

PANDE I, UPPAL SS, KAILASH S, KUMAR A, MALAVIYA AN. Behcet's disease in India: a clinical, immunological, immunogentic and outcome study, 825

PANDE I See also Sood A; Uppal SS

PANDE IRA See Uppal SS

Panniculitis, arthritis and pancreatitis, 680

PANUSH RS See Darlington LG

PAPASAVVAS G See Raja KB

Paralysed patients, axial skeletal changes may mimic ankylosing spondylitis,

Paraproteins, prevalence of serum bands in rheumatoid arthritis, (A2) 5

Parity and risk of rheumatoid arthritis in Finnish women, 625

PARKER FP See Seed MP

PARKER KH See Whyte J

PARTSCH G. High free and latent collagenase activity in psoriatic arthritis synovial fluid, (L) 84

Parvovirus B19

arthritis, outbreak in North Dublin, (A1) 110

brachial plexitis in a patient with acute infection, (L) 480

in serum samples submitted for a rheumatoid factor test, (A2) 33 study of the role of in rheumatoid arthritis, 809

Parvovorus arthritis, relationship between prolonged joint symptoms, hypocomplementaemia and HLA type, (A1) 111

PASCUAL E, JOVANI V. Quantitative study of the phagocytosis of urate crystals in the synovial fluid of asymptomatic joints of patients with gout, 724

PASCUAL E. Management of gout in patients with renal insufficiency and a gastrointestinal bleed—an answer to Dr Dunne's letter, (L) 481

PATEL S, PEARSON D, BHALLAH A, MASALANKA W, HOSKING DJ. Changes in bone mineral density with bisphosphonate treatment of Paget's disease of bone, (A1) 130

PATEL S See also Craig JP; Johnson SM; Kaye SA

Patellofemoral joint, tibiofemoral osteophyte in the anteroposterior radiograph predicts joint space narrowing, (A1) 136 Pathological synovial fluids promote apoptosis of peripheral blood and

synovial neutrophils: a role in resolution of synovitis?, (A1) 58

activity as an objective measure of NSAID therapy: an Arthrotec and NUMACT study, (A2) 15

with rheumatoid arthritis and microscopic polyarteritis, (L) 1185

Patient education

arthritis programme, five-year follow-up of a controlled trial, 647 treatment or nice extra, (E) 703

Patients

with anti-RNA polymerase antibodies have diffuse scleroderma and exhibit a speckled pattern anti-nuclear antibody distinct from anti-Scl-70, (A1) 66

with rheumatoid arthritis not attending a specialist clinic have similar levels of disease severity as clinic attenders, (A1) 88

Pattern of distribution of osteoarthritis of the hand joints: a comparative study between a clinical and a community cohort, (A1) 133 PAULUS HE. Clinical trial design for evaluating combination therapies, (S2)

PAYTON A See Brennan P

PEACOCK DJ, EGGER P, HOBBS R, FALL C, COOPER C. Spinal osteoarthritis and osteoporosis: evidence for a negative association independent of obesity, (A1) 74

PEACOCK DJ, EGGER P, TAYLOR P, CAWLEY MID, COOPER C. Utility of lateral bone density measurements in osteoarthritis of the lumbar spine. (A1) 127

Peak bone mass in women, life-style, environmental and medical factors influencing, 620

PEARSON D See Patel S

PEARSON JD See Denton CP; Mackworth-Young CG; Shiwen X

PEASE CT See Bhakta BB

PEGO JM See Belzunegui J

PEJOVIC M, STANKOVIC A, MITROVIC DR. Determination of the apparent synovial permeability in the knee joint of patients suffering from osteoarthritis and rheumatoid arthritis, 520

PEMBERTON S See Inamo Y

Penetrance and linkage of HLA and rheumatoid arthritis in extended pedigrees, (A1) 139

Penicillamine and sulphasalazine, audit of patients taking them individually, sequentially and concurrently in inflammatory arthritis, (A2) 17

PEPPER L See Donn RP: Thomson W Peptic ulcer disease, recurrent, double-blind placebo-controlled study of ranitidine for prevention in rheumatoid arthritis patients on nonsteroidal anti-inflammatory drugs, (A1) 94

Peptide transporter genes (TAP) polymorphisms and genetic susceptibility to rheumatoid arthritis, 207

PEREIRA RS See Kaminski ER

Perforations, peptic ulceration and bleeding on NSAIDs. Two randomised controlled trials of nabumetone, diclofenac SR and piroxicam in the UK, (A1) 95

Pericardial effusion, massive in scleroderma: a review of five cases, 564

Peripheral blood mononuclear cells

methotrexate action in rheumatoid arthritis: stimulation of cytokine inhibition and inhibition of chemokine production, 602

rheumatoid population, DNA damage assessment by single cell gel electrophoresis, (A1) 153

unfractionated, BCL-2 expression is unaltered in patients with systemic lupus erythematosus, 316

Peripheral blood T cell receptor Vβ family expression in multicase rheumatoid arthritis families, (A1) 141

Peripheral joints, intra-articular pressure changes in normal and rheumatoid, (A1) 46

Peripheral neuropathy, anti ganglioside antibodies, (A2) 3

Peripheral oedema, rheumatoid arthritis in the elderly: is it a good prognostic feature at onset?, (A2) 10

Peris P, Guañabens N, Monegal A, Suris X, Alavarez L, Martinez de Osaba MJ, Hernandez MV, Muñoz-Gomez J. Actiology and presenting symptoms in male osteoporosis, 936

PERIS P See also Pons F

PERKINS MN See Cruwys SC

PERRET J See Dobbie JW PERRETT D See Evans MD; James IT

PERRETTI M, WHELLER SK, FLOWER RJ, GOULDING NJ. Corticosteroid regulation of intracellular lipocortin-1 expression in murine leukocytes,

(A1) 147 PERRY MJ, MORTUZA FY, PONSFORD FM, ELSON CJ, ATKINS RM. Analysis of cell types and mediator production from tissues around loosening

joint implants, 1127 Personal illness models and the self management of arthritis, (A1) 90

PERVEEN-GILL Z See Harbuz MS

PETERS AM See Chapman PT

PETERS TJ See Raja KB

PETRIE A See Darlington LG PHELAN M See MacDonald AG

PHILIPS DI See Camp AV; Collier SM PHIPPS JL See Hill MB

Phrenicus palsy in progressive systemic sclerosis, 684

Physical activity monitoring, NUMACT: a novel method in JCA, (A2) 23 Physical function, association with psychological problems in children with iuvenile chronic arthritis, 470

Physiotherapy, manipulative for spinal problems in primary care: outcomes of care, 1070

PICCIOTTO A See Cutolo M

Picco P See Gattorno M

PICKAVANCE L, GRIFFITHS G, MCKENNA F. Diclofenac is a better analgesic than paracetamol in patients with osteoarthritis of the hip, knee and lumbar spine, (A1) 137

PIGNATTI P See Massa M

PILE K See Hill C

PILE KD See Brown MA

PILLEMER S See Tilley BC

PIMM TJ, BYRON MA, CURSON D. Personal illness models and the self management of arthritis, (A1) 90

PINCUS T, CALLAHAN LF. Prognostic markers of activity and damage in rheumatoid arthritis: why clinical trials and inception cohort studies indicate more favourable outcomes than studies of patients with established disease, (E) 196

PINCUS T. Long-term outcomes in rheumatoid arthritis, (S2) 59

PINEDA M See Tinahones FJ

PINILLOS V See Gonzalez-Crespo MR

PIPITONE N See Bajocchi GL; Pitzalis C

Piroxicam

perforations, peptic ulceration and bleeding on NSAIDs. Two randomised controlled trials of nabumetone, diclofenac SR and piroxicam in the UK, (A1) 95

total and free methotrexate pharmacokinetics, in rheumatoid arthritis

patients, 421
PISTOIA V See Gattorno M

PITT P See De Lord D

PITZALIS C, CAULI A, YANNI G, PIPITONE N, PANAYI GS. The selective accumulation of cutaneous lymphocyte antigen positive lymphocytes in the skin but not in the joint of inflammatory arthropathies, (A1) 64

PITZALIS C, LEE A, PIPITONE N, BAJOCCHI GL, PANAYI G. Corticosteroids inhibit lymphocyte adhesion to endothelial cells in vitro: confirmation using the SV40-transformed EC line LT4, (A1) 146

PITZALIS C See also Bajocchi GL

Place of acupuncture in medicine today, (E) 404

Planovalgus rheumatoid foot—is tibialis posterior tendon rupture a factor?, 645

PLANT M See Hassell A

PLANT MJ, JONES PW, DAWES PT. Eight year outcome of patients presenting with early non-erosive rheumatoid arthritis, (A2) 8

PLANT MJ See also Aung TS; Borg AA; Carpenter EH; Hassell AB Plantar fasciitis, scintigraphic localization of steroid injection site, (A1) 42

Plantar heel pain, therapeutic ultrasound as a treatment, (A1) 42

Plasma exchange, type 1 cryoglobulinaemia associated with a thymic tumour: successful treatment, 285

Plasminogen activator inhibitor-1 and urokinase levels in synovial fluids in normal, osteoarthritis and rheumatoid arthritis knees, (A1) 59

PLATT R See Troughton PR

PLAZAOLA I See Belzunegui J Pneumocystis carinii pneumonia in a patient with rheumatoid arthritis, not on immunosuppressive therapy and in the absence of human immunodefi-

ciency virus infection, 677 Polyarteritis, microscopio

comparison of anti-neutrophil cytoplasmic antibody positive with ANCA negative, (A1) 67

and rheumatoid arthritis, (L) 1185

Polyarteritis nodosa, has it been classified out of existence?, (A2) 25 Polyarthralgia, spondylitis-like and symmetric associated with gemfibrozil

therapy, (L) 692 Polyarthritis

inflammatory

HLA associations developing post-immunisation, (A1) 76 predictors of disability one year after presentation with inflamma-tory polyarthritis, (A1) 86

masquerading as the bends, (A2) 53

migratory and chronic abdominal sepsis, (L) 478

new case as a presenting sign of non-Hodgkin's lymphoma, (L) 794 and non-Hodgkin's lymphoma, (L) 293

non-Hodgkin's lymphoma presenting as, (L) 182

in the rat: effects of tolerance and sensitization to the bacterial extract OM-89 with a possible mode of action, 1135 Polymorphism in an Spl consensus binding motif in the first intron of

collagen I alpha 1 is related to postmenopausal osteoporosis, (A1) 78 Polymorphism of the heat-shock protein gene HSP70-2 in systemic lupus erythematosus, 721

Polymorphisms of the vitamin D receptor gene in postmenopausal female athletes, (A1) 79

Polymorphonuclear neutrophils, defective spontaneous and bacterial lipopolysaccharide-stimulated production of interleukin-1 receptor antagonist in patients with active systemic lupus erythematosus, 107

Polymyalgia is associated with both HLA-DRB1*0401 and DRB1*0404, (A1) 139

Polymyalgia rheumatica

controlled trial of intramuscular methylprednisolone against oral prednisolone: a 48 week follow up, (A1) 69 serial changes in bone mineral density, (A1) 74

Polymyalgia-like syndrome, Lyme borrelosis presenting as, (L) 392

Polymyositis

anti Jo-1 antibody-positive, effective treatment with cyclosporin, (L) 1187

changing autoantibody profiles with variable clinical manifestations, 915 indium-111 antimyosin antibody and muscle ultrasound scans in the evaluation of patients, (A2) 27

patients show consistent changes on 1H-NMR spectroscopy, (A1) 125 proton-NMR spectroscopy and the diagnosis, (A2) 27

PONS F. PERIS P. GUANABENS N. FONT J. HUGUET M. ESPINOSA G. INGELMO M, MUÑOZ-GOMEZ J, SETOAIN. The effect of systemic lupus erythematosus and long-term steroid therapy on bone mass in pre-menopausal women, 742

PONSFORD FM See Perry MJ

POON SP See Mok CC Poor Gy See Kiss Cs

POPE D See Chambers H

POPERT AJ, JONES PJ, FARR M. Rheumatoid arthritis: relation of treatment to remission, relapse, progression, (A2) 18

PORTER DR See McInnes IB

Possible mechanisms of action of methotrexate in patients with rheumatoid arthritis, (S2) 26

Postmenopausal bone loss, early and polymorphisms of the vitamin D receptor, (A1) 79

POTTINGER BE See Mackworth-Young CG

POULTON K, HILLARBY MC, OLLIER WER. Soluble FAS is increased in patients with SLE, (A1) 5

POWELL RJ See Hodges E; McDonagh JE; Pallis M

Practical clinical pharmacology and drug interactions of low-dose methotrexate therapy in rheumatoid arthritis, (S2) 20

Predictive factors of severity of spondylarthropathy in North Africa, 1139 Predictive value

of proteoglycan antibodies for erosions in early inflammatory arthritis, (A1) 104

of von Willebrand factor and capillary microscopy in systemic sclerosis (scleroderma)—a prospective study, (A1) 27

Predictors

of disability one year after presentation with inflammatory polyarthritis, (A1) 86

of resolution in inflammatory polyarthritis, (A1) 85

Prednisolone

oral, controlled trial against intramuscular methylprednisolone in polymyalgia rheumatica: a 48 week follow up, (A1) 69 radiological progression in rheumatoid arthritis during a randomised controlled trial, (A1) 80

and Behçet disease relationship study, (A1) 24

induced glycosylation changes in rheumatoid arthritis, (A1) 62 and rheumatoid arthritis—bits and pieces in a puzzle, (E) 1

Preliminary audit of the management of acute low back pain in the Kettering district, (L) 693

of bacterial flora-derived antigen in synovial tissue macrophages and dendritic cells, 1112

of synovial fluid glycolipid antibodies in rheumatoid arthritis-synovial neuropathy or synovial damage, (A2) 4

Presentation, clinical patterns in 722 patients with early rheumatoid arthritis, examining the influences of age at onset and gender, (A2) 8

PRESETNIK M See Jevtic V

Prevalence

and clinical significance of subungual splinter haemorrhages in patients with an antiphospholipid syndrome, (A1) 16

of low back pain and patterns of consultation, (A2) 38

of rheumatoid arthritis in affluent and poor urban communities of Pakistan, 252

of serum paraprotein bands in rheumatoid arthritis, (A2) 5

of vertebral fractures in patients with RA, treated with corticosteroids, (A1) 73

of xerostomia in a population with primary generalised osteoarthritis,

Prevention and epidemiology of non-steroidal anti-inflammatory drug effects in the gastrointestinal tract, (S1) 5

PRICE EJ See Brookes SM; Rigby S

PRICE T See Glennon P

PRICE TR See Dolan AL

PRIESTLEY S See Lindroth Y

Primary care

inflammatory back pain, 1074

outcome, manipulative physiotherapy for spinal problems, 1070 Primary nocardial meningitis in systemic lupus erythematosus, 178

Primary pyomyositis. Two more cases with atypical presentation in diabetic hosts, (L) 482

Primary tuberculous muscle abscess in a patient with systemic lupus erythematosus, 1177

PRINS APA See Griep EN

PRITCHARD T See Ferry S

Pro-MMP1, serological joint markers-a comparison of serum proMMP-3, proMMP-1 and disease activity in RA, (A1) 60

Pro-MMP3

matrix metallo-proteinase, in vitro effect of leflunomide on RA synovial fibroblasts proliferation, (A1) 60

serological joint markers—a comparison of serum proMMP-3, proMMP-1 and disease activity in RA, (A1) 60

Procainamide-induced lupus, neutrophil-derived tumour necrosis factor-α: a potential mechanism for promotion of inflammation, (A1) 3

Procollagen aminopeptide type III, and various cytokine levels determination in bronchoalveolar lavage fluid of the patients with pulmonary fibrosis: inverse correlation between type III procollagen aminopeptide and interferon- Γ in progressive patients, 31

Production and analysis of the binding characteristics of four human monoclonal IgG anticardiolipin antibodies, (A1) 12

Profile of HLA-B27-related unclassifiable seronegative spondylarthropathy in females and its comparison with the profile in males, 137

Prognostic factors in rheumatoid arthritis. Comparative study of two subsets of patients according to severity of articular damage, 529

Prognostic markers of activity and damage in rheumatoid arthritis: why clinical trials and inception cohort studies indicate more favourable outcomes than studies of patients with established disease, (E) 196

Prognostic value of quantitative measurement of rheumatoid factor in early rheumatoid arthritis, 1146

Proinflammatory role of tissue kallikrein in modulating pain in inflamed joints, (L) 88

Prolactin levels in lupus pregnancy do not differ significantly from those in controls. There is no correlation between disease activity and prolactin levels, (A1) 13

Proposed modification to Larsen's scoring methods for hand and wrist radiographs, 56

Proprioception, effect of hypermobility syndrome on knee joint, 121

Prospective, placebo-controlled study of single and double-blind food challenges in patients with rheumatoid arthritis after dietary manipulation therapy, (A1) 94

Prospective study of 110 high-risk pregnancies in a lupus pregnancy clinic, (A1) 13

PROST A See Berthelot JM

Prostacyclin, use of intravenous to treat pulmonary hypertension associated with systemic sclerosis, (A1) 30

Prostanoid modulation of synovial antigen-specific CD4+ T-cell cytotoxic function in rheumatoid arthritis, 113

Prosthetic joints, role of prophylactic antibiotics in the prevention of late infection. Results of a questionnaire and review of the literature, 380

Protease inhibitor phenotypes, HLA-DR4 and keratoconjunctivitis sicca association with pulmonary abnormalities in rheumatoid arthritis, 37 Protective role for testosterone in adjuvant-induced arthritis, 1117

Protein S deficiency, acquired, and cerebral venous thrombosis: an uncommon cause of headache in systemic lupus erythematosus, 572

Proteinaceous lymphadenopathy in a patient with known rheumatoid arthritis-case report and review of literature, 1087

Proteinuria

should tests for be included in the monitoring schedule of sulphasalazine?, 790

suppression and prolongation of survival in MRL-lpr/lpr mice treated with sulphasalazine, (L) 85

Proteoglycan antibodies, predictive value for erosions in early inflammatory arthritis, (A1) 104

Proteoglycans associated with experimental reactive AA amyloidosis, (A1) 118

Proteus mirabilis

Escherichia coli and Klebsiella pneumoniae antibodies in the sera of ankylosing spondylitis patients with/without iritis and enthesitis, 418

Klebsiella pneumoniae and Escherichia coli antibodies in the sera of patients with axial and peripheral form of ankylosing spondylitis,

Proton-NMR spectroscopy and the diagnosis of polymyositis and dermatomyositis, (A2) 27

PROUSE P, COX N, DAVIS P, DEVLIN J, DIXEY J, EMERY P, GALLIVAN S, GOUGH A, JAMES D, WILLIAMS P, WINFIELD J, YOUNG A. Clinical patterns at presentation in 722 patients with early rheumatoid arthritis, examining the influences of age at onset and gender, (A2) 8

PROUSE P See also James D

Pruritis, generalized in the rheumatoid patient-consider all possible causes, (L) 295

Pseudogout masking brucellar arthritis, (L) 294

Psoriasis, cutaneous lymphocyte antigen expression by peripheral blood and synovial fluid lymphocytes, (A1) 147

Psoriatic arthritis

cutaneous lymphocyte antigen expression by peripheral blood and synovial fluid lymphocytes, (A1) 147

high free and latent collagenase activity in synovial fluid, (L) 83, (L) 84 HLA associations of different patterns, (A1) 143

HML-1 expression by peripheral blood and synovial fluid lymphocytes, (A1) 148

interleukin-2 in and in seronegative spondylarthritis but not in rheumatoid arthritis synovium, (A1) 151

sulphasalazine-induced lupus, 383

Psoriatic arthropathy after BCG immunotherapy for bladder carcinoma, (L)

Psychological distress and local tenderness but not radiographic change or weakness are associated with disability in hospital referred patients with knee osteoarthritis, (A1) 136

Psychological problems, association with physical function in children with juvenile chronic arthritis, 470

PULLAR T See McInnes IB

Pulmonary fibrosis

apical, high prevalence in ankylosing spondylitis by high resolution computerized tomography, (A1) 117

determination of various cytokines and type III procollagen aminopeptide levels in bronchoalveolar lavage fluid of the patients: inverse correlation between type III procollagen aminopeptide and interferon- Γ in progressive patients, 31

Pulmonary function in rheumatoid arthritis treated with low dose methotrexate: a longitudinal study, (A1) 100

Pulmonary hypertension, use of intravenous prostacyclin to treat in systemic sclerosis, (A1) 30

Pulse cyclophosphamide therapy in systemic vasculitis, (A1) 21

PUNN YL See Williams AS PUTHRASINGHAM S See De Lord D

PYERITZ RE See Grahame R

PYKE S See Sutcliffe N

Pyoderma gangrenosum, rheumatoid associated, cyclosporin A in treatment, (L) 182

Pyomyositis

Salmonella in patients with the human immunodeficiency virus, 568 primary. Two more cases with atypical presentation in diabetic hosts, (L) 482

Pyridinium crosslinks

effect of 2nd line antirheumatic therapy on excretion in patients with established RA, (A1) 99

stability to photolysis, (A1) 54

urinary, receiver operating characteristic analysis of in arthritis, (L) 792 Pyridinoline and deoxypyridinoline may be influenced by turnover of nonosseous tissues in patients with joint disease, (L) 1186

Quadriceps activation and rehabilitation following anterior cruciate ligament rupture, (A1) 135

Quality of life

assessments used in RA, (A2) 11

relationship with back pain and psychological status in women in the general population: the Chingford study, (A1) 40

Quantification of joint laxity. The non-dominant (Spanish) modification, (L)

Quantitative analysis of lymphocyte adhesion molecule deployment in systemic lupus erythematosus, (A1) 4

Quantitative cartilage imaging, (S1) 29

Quantitative study of the phagocytosis of urate crystals in the synovial fluid of asymptomatic joints of patients with gout, 724

QUEIRO R. Psoriatic arthropathy after BCG immunotherapy for bladder carcinoma, (L) 1097

QUIN C See Hodges E

REECE R, EMERY P. Needle arthroscopy, (E) 1102

Radiographic within-subject variability is low when measuring knee joint space with digital image analysis, (A1) 122

Radiological assessment, proposed modification to Larsen's scoring methods for hand and wrist radiographs, 56

Radiological damage, centile curves: rheumatoid arthritis population, (A1)

Radiological medial joint space narrowing and meniscal lesions, (L) 991 Radiological progression

can we differentiate between over one year in patients with rheumatoid arthritis randomly allocated to im gold or hydroxychloroquine?, (A1) 107

in rheumatoid arthritis during a randomised controlled trial of prednisolone, (A1) 80

Radiology, ankylosing spondylitis: the correlation between a new metrology score and, 767

RAFFERTY JA, GREEN H, GRIFFIN K, MARGISON GP, HERRICK AL. Decreased DNA repair proficiency: a factor in susceptibility or a feature of disease in autoimmune conditions?, (A1) 4

RAFI S, SOMASUNDARAM S, SCOTT DL, WRIGGLESWORTH J, BJARNASON I. Uncoupling of oxidative phosphorylation is the initiating factor in NSAID-induced gastrointestinal damage, (A1) 93

RAFI S See also Mahmud T

RAHMAN MM. Proinflammatory role of tissue kallikrein in modulating pain in inflamed joints, (L) 88

RAHMAN MAA. MENON S. LONGHURST CM. RAVIRAJAN CT. LATCHMAN DS, ISENBERG DA. Sequence analysis of four human monoclonal IgG anti-cardiolipin antibodies, (A2) 28

RAHMATALLA AA See Aung TS

RAINE AEG See D'Cruz D

Raised levels of IgA and IgM rheumatoid factor in patients with Lyme arthritis and neuroborrelosis, (A1) 110

RAJA KB, DUANE P, SNAPE SD, SIMPSON RJ, PAPASAVVAS G, PETERS TJ. in vitro duodenal iron uptake and serum and mucosal iron protein levels, with special reference to rheumatoid arthritis, 1041

RAJAN KT See Byrne P

RAJAPAKSE C, RAJAPAKSE Y, TREADWELL B, CORKILL M. Slow acting drug therapy of rheumatoid arthritis: treatment duration and side effect profile, (A1) 103

RAJAPAKSE Y See Rajapakse C

RAJARATNAM G See Hillman M

RALPHS A See Fisher M

RALSTON SH See Grabowski PS; Grant SFA

RAMOS A See Gonzalez-Crespo MR

RAMSEY DJC See Darlington LG RAMSEY NW See Darlington LG

RAMSEY S See Cawston TE

Randomized controlled trial of brisk walking in the prevention of osteoporosis and falls, (A1) 72

Randomized controlled trial of four different exercise programs in patients with rheumatoid arthritis, (A1) 93

Ranitidine, double-blind placebo-controlled study for the prevention of recurrent peptic ulcer disease in rheumatoid arthritis patients on nonsteroidal anti-inflammatory drugs, (A1) 94

RANKIN ECC, CHOY EHS, KASSIMOS D, KINGSLEY GH, SOPWITH AM, ISENBERG DA, PANAYI GS. The therapeutic effects of an engineered human anti-tumour necrosis factor alpha antibody (CDP571) in rheumatoid arthritis, 334

RANKIN ECC, CHOY EHS, KASSIMOS D, KINGSLEY GH, SOPWITH AM, ISENBERG DA, PANAYI GS. Repeated dose of an engineered anti-TNFa antibody, CDP571 in patients with rheumatoid arthritis, (A1) 81

RANKIN ECC, RAVIRAJAN CT, EHRENSTEIN MR, BODMAN KB, CHOY EHS, PANAYI GS, ISENBERG DA. Serological effects following treatment with an engineered human anti-TNFa antibody CDP571, in patients with rheumatoid arthritis, (A1) 101

RASPE H, KOHLMANN T, ENGSTER M, BROCKOW T. Back pain-more than pain in the back II: tender points in graded back pain, (A2) 38

RATCLIFFE LT, LUKEY PT, MEYERS OL, RESS SR. Prostanoid modulation of synovial antigen-specific CD4+ T-cell cytotoxic function in rheumatoid arthritis, 113

RAUS J See Vandevyer C

RAVIRAJAN C See Menon S

RAVIRAJAN CT See Rahman MAA; Rankin ECC

Raynaud's phenomenon

primary

neurovascular control mechanisms and in systemic sclerosis as assessed by iontophoresis and laser Doppler techniques, (A2)

reduced vasodilatation to acetylcholine in the finger microcirculation, (A1) 25

and systemic sclerosis, capillary measurements by video capillary microscopy, (A1) 25

primary and secondary, sustained-release transdermal glyceryl trinitrate patches as a treatment, 636

Reactive arthritis

analysis of Chlamydia trachomatis specific synovial fluid derived T cell clones, (A1) 152

oncogene expression in synovial fluid cells: a brief report, 805

urogenital swab culture is the only useful diagnostic method for the detection of the arthritogenic infection in extra-articularly asymptomatic patients with undifferentiated oligoarthritis, 838

Reactive haemophagocytic syndrome, successful management in systemiconset juvenile chronic arthritis, (L) 888

Reactive oxygen species, effects on the ability of IgG to interact with C1q, (A1) 45

READER JA See Nash MC

Receiver operating characteristic analysis of urinary pyridinium cross-links in arthritis, (L) 792

Recent-onset inflammatory arthritis, muscle strength in healthy people and in patients suffering from, 355

Red blood cell aggregation, dietary essential fatty acid supplements reduce in healthy volunteers, (A1) 28

REDKO C See Cury SE

Reduced vasodilatation to acetylcholine in the finger microcirculation in primary Raynaud's phenomenon, (A1) 25

Reduction in cell adhesion molecule levels following sulphasalazine in patients with rheumatoid arthritis, (A1) 98

REES DHE, O'CONNELL S, BROWN MM, ROBERTSON J, AXFORD JS. The

value of serological testing for Lyme disease in the UK, 132
REES DHE, STEERE AC, AXFORD JS. Raised levels of IgA and IgN rheumatoid factor in patients with Lyme arthritis and neuroborrelosis,

REEVE J See O'Neill TW

REEVES WH See Satoh M

Referral guidelines assessment to a bone densitometry service, (A1) 37

REFVEM OK See Glennås A

REGAN M, DEIGHTON C, DOHERTY M. Evidence for genetic anticipation in nodal osteoarthritis, (A1) 133

REGAN M See also Jones AC; Ledingham J

Regional audit of adult onset Still's disease, (A2) 22

Regulation of human complement component C4 gene expression, (A1) 156 Rehabilitation, Sixth European Regional Conference of Rehabilitation International, 283

REID DM See Garton MJ; Grant SFA; Martin JC; O'Neill TW; Torgerson

REID J See Collins AJ

Reiter's syndrome, sexually acquired arthritis in nineteenth century London, (A2) 31

Relationship

between back pain, psychological status and quality of life in women in the general population: the Chingford study, (A1) 40

between osteophytosis and osteoarthritis in joints, (A1) 132

between serum cartilage oligomeric matrix protein levels and disease progression in osteoarthritis of the knee joint, 306

between synovitis and erosions in rheumatoid arthritis, (A1) 105

bewteen prolonged joint symptoms, hypocomplementaemia and HLA type in parvovirus arthritis, (A1) 111

of fatigue in ankylosing spondylitis to disease activity, sleep and other factors, (A2) 36

of ICAM-1 with E-selectin in serum, synovial fluid and synovial membrane in inflammatory arthritis, (A1) 148

of IGF-1 and osteoarthritis in women aged 45-64 in the general population: the Chingford study, (A1) 137
Relative contributions of HLA-DQA and complement C4A loci in

determining susceptibility to systemic lupus erythematosus, 221

Relative importance of pain and X-ray damage to disability in long-standing RA, (A2) 12

REMITA K See Kennedy LG

Renal function and non-steroidal anti-inflammatory drugs, (S1) 25

RENNIE JAN See Keir G

Repeated dose of an engineered anti-TNFa antibody CDP571 in patients with rheumatoid arthritis, (A1) 81

Repetitive strain injury, use of three-phase bone scan in the diagnosis, (A2) 39 Reproductive factors, breast feeding, and rheumatoid arthritis. A prospective study, 542

Responsiveness of Keitel functional index compared with laboratory measures of disease activity in rheumatoid arthritis, 141

RESS SR See Ratcliffe LT

Retinol binding protein, urinary excretion in SLE, (A1) 17

Retrovirus envelope glycoprotein, antibodies to a human endogenous in rheumatoid arthritis, (A1) 154

Retroviruses, absence of exogenous in Felty's syndrome, (L) 185

REUNANEN A See Heliövaara M

Review, routine follow up of RA patients: what dictates periodicity?, (A2) 12 Rheumatic diseases

recognition of case descriptions by general practitioners, assessing clinical competence, 375

and rheumatoid arthritis amongst Pakistanis in London and Karachi, (A1) 33

sulphasalazine-induced autoimmune abnormalities, 435, (L) 1097

Rheumato-microbiology interface, (E) 99

Rheumatoid arthritis

in vitro duodenal iron uptake and serum and mucosal iron protein levels, 1041

in vitro immunomodulatory effect of tetracyclines. (A1) 152

abnormal tear evaporation rate and osmolality, (A1) 91

absence of correlation with IL-1a intron 6 polymorphism, 1123

active, in-patient treatment: clinical course and predictors of improvement, 847

adverse events in methotrexate-treated patients, (S2) 49

advice to car drivers, (L) 796

altered levels of soluble adhesion molecules, 814

analysis of the discrepancy between the clinical and radiological course,

anti ganglioside antibodies, (A2) 3

anti-CD44 antibody induces receptor shedding from synovial leukocytes and fibroblasts, (A2) 30 antibodies to the 65 kDa mycobacterial stress protein in West Africans

with rheumatoid arthritis, tuberculosis and malaria, 352

antibodies to a human endogenous retrovirus envelope glycoprotein, (A1) 154

antithrombin III inactivation, (A1) 84

are slow-acting anti-rheumatic drugs monitored too often? An audit of current clinical practice, 966

association of HLA-DR4, protease inhibitor phenotypes and keratoconjunctivitis sicca with pulmonary abnormalities, 37

atlanto-occipital subluxation demonstrated by magnetic resonance imaging, 787

audit of methotrexate and folic acid for. Experience from a teaching centre, 971

audit of thyroid function tests in a rheumatic disease clinic-results and impression that thyroxine excerbates RA, (A2) 21 breast feeding and other reproductive factors. A prospective study, 542

broadband ultrasound characteristics, (A1) 123

can we differentiate between radiological progression over one year in patients randomly allocated to im gold or hydroxychloroquine?, (A1) 107

centile curves of radiological damage, (A1) 106

clinical patterns at presentation in 722 patients, examining the influences of age at onset and gender, (A2) 8

combination of sulphasalazine and methotrexate in the management of-view based on personal clinical experience, (S2) 109

combination therapy in, (L) 580

combining sulphasalazine and methotrexate: early clinical impressions, (S2) 104

comparable co-expression of VCAM-1* and DAF* on synovial intimal

cells and follicular dendritic cells explains synovial targeting, (A2) 4 comparative study of tenidap, a cytokine-modulating anti-rheumatic drug, and diclofenae: a 24-week analysis of a 1-year clinical trial, 554

comparison of intravenous, intramuscular and oral pulse steroid therapies in the management of, (A2) 19

comparison of S-oxidation status in Asian and Caucasian patients, (A2)

compound heterozygosity of HLA-DR4 and DR1 antigens in Asian Indians increases the risk of extra-articular features, 41

cyclosporine A in early patients: a 24 months chloroquine controlled follow up study, (A1) 99

deoxycoformycin, a selective lymphocyte depleting drug, induces flare, (A1) 102

determination of the apparent synovial permeability in the knee joint, 520

disability in-it's relation to clinical and radiological courses over 3 vears, (A2) 11

discordance between objective and subjective assessment of functional ability, 951

diurnal variation in 24 hour ambulatory recordings of patient activity is different from osteoarthritis, (A2) 16

dominant human leukocyte antigen DR4-Dw15 restricted mycobacterial 65 kDa antigen-specific T-cell immunity in Chinese patients, 498 double blind placebo controlled trial of minocycline, (A2) 16

early hand bone densitometry: a measure of progression despite clinical control, (A1) 108

and joint erosions, (S2) 74

prognostic value of quantitative measurement of rheumatoid factor, 1146

therapeutic approaches. How early? How aggressive?, (S2) 87 effect of high-dose corticosteroid pulse therapy on bone metabolism, (A1) 128

effect of methotrexate alone or in combination with sulphasalazine on the production and circulating concentrations of cytokines and their antagonists, 747

effect of sulphasalazine on neutrophil superoxide protection, (A2) 14 effects of elemental diet and subsequent food reintroduction, 270

effects of joint washout and steroid injection compared with either joint washout or steroid injection alone in rheumatoid knee effusion, 771 effects on the patient's spouse, (A2) 10

efficacy of methotrexate, (S2) 43

eight year outcome of patients presenting with early non-erosive, (A2) 8 in the elderly: is peripheral oedema at onset a good prognostic feature?, (A2) 10

endogenous opiates: are they responsible for the defective hypothalamic response to inflammatory stimuli, (A1) 84

evaluation of microwave thermography of the hand and wrist in rheumatoid arthritis, (A2) 13

factors associated with the development of vasculitis in rheumatoid arthritis: results of a case-control study, (A1) 86

familial, genetic anticipation in, (A2) 6

familial risk is low in first degree relatives of probands with inflammatory arthritis ascertained from a population-based register, (A1) 80 feasibility study of homeopathic remedy, (A2) 17

ferritin is an acute phase protein in longstanding and recent onset, (A1)

foetally inherited paternal HLA does not explain post partum susceptibility in the mother, (A1) 142

foot and ankle in, 443

galactosyltransferase isoenzyme changes occur in, (A1) 61

global tests for multiple outcomes in trials, (A1) 89

health economics as an aspect of health outcome: basic principles and application, 774

health status and joint damage divergence, (A1) 107

heterogeneity of disease phenotype in monozygotic twins concordant for,

and HLA linkage and penetrance in extended pedigrees, (A1) 139 HLA phenotypes and therapeutic response to parenteral gold, (A1) 97 HML-1 expression by peripheral blood and synovial fluid lymphocytes, (A1) 148

human anti-chimaeric antibody response after treatment with a chimaeric anti-CD4 monoclonal antibody, cM-T412, (A1) 101

human leucocyte antigen phenotypes and gold-induced remissions, 343 identifying iron deficiency anaemia by simple laboratory measures, (A1) 83

IL-2R levels: poor correlation with clinical activity is due in part to disease duration, 1037

impact on body image, (A1) 89

incidence is not related to socioeconomic factors, (A2) 7

increased IL-1 receptor antagonist (IL-1ra) production and decreased IL-1β/IL-1ra ratio in mononuclear cells, 24

influence of glutathione S-transferase (GSTM1 and GSTT1) polymorphisms on the incidence and severity, (A2) 18 isotype variation of TNF monoclonal antibodies. Is there a therapeutic

difference?, (A1) 153 late-onset and young-onset. A comparative clinical and radiological

study. (A1) 106 levels of circulating intercellular adhesion molecule-1 and -3 but not circulating endothelial leucocyte adhesion molecule are increased, 311

long-term outcomes, (S2) 59

long-term usage and side-effect profile of sulphasalazine, (S2) 40 macroamylasemia: a possible cause of unexplained hyperamylasemia,

major life events as a result. A comparison between nine centres in 446 patients in the first three years of disease, (A2) 9

management of oral complications of disease modifying drugs, (A2) 20 measurement of pain, (A1) 91

measuring outcome in-which measures are suitable for routine clinical

methodological problems in comparing the severity between populations, 781

methotrexate action in: stimulation of cytokine inhibition and inhibition of chemokine production by peripheral blood mononuclear cells,

MHC encoded TN-X gene may be important, (A1) 140

with microscopic polyarteritis, (L) 1185 mycoplasmal RNA and 5'-nucleotidase in the synovial fluid, (A2) 5

with myelopathy, MRI characteristics of the cervical spine, canal and cord, (A1) 124

neurological side effects in two patients receiving gold injections, 989 new therapeutic approaches in autoimmune rheumatic diseases, with special emphasis on rheumatoid arthritis, (E) 193 nitric oxide production by primary synovial cultures, (A1) 44

oligosaccharide changes determined by fluorophore assisted electrophoresis and lectin binding analysis, (A1) 63

oncogene expression in synovial fluid cells: a brief report, 805

oral steroids as bridge therapy in rheumatoid arthritis patients starting with parenteral gold. A randomized double-blind placebo-controlled trial, 347

orbital pseudo-tumour: a new ocular entity?, (A2) 55

outcome of surgery to stabilise the cervical spine-pain, neurological status and patient satisfaction, (A1) 92

overview of combination second-line or disease-modifying antirheumatic drug therapy, (S2) 96

parity and risk in Finnish women, 625

patients cannot accurately report signs of inflammatory activity, 547 patients not attending a specialist clinic have similar levels of disease severity as clinic attenders, (A1) 88 peptide transporter genes (TAP) polymorphisms and genetic suscept-

ibility to, 207

peripheral blood T cell receptor VB family expression in multicase families, (A1) 141

Pneumocystis carinii pneumonia in a patient, not on immunosuppressive therapy and in the absence of human immunodeficiency virus infection, 677

possible mechanisms of action of methotrexate in patients, (S2) 26

practical clinical pharmacology and drug interactions of low-dose methotrexate therapy in rheumatoid arthritis, (S2) 20

and pregnancy-bits and pieces in a puzzle, (E) 1

pregnancy induced glycosylation changes, (A1) 62

presence of synovial fluid glycolipid antibodies--synovial neuropathy or synovial damage, (A2) 4

prevalence in affluent and poor urban communities of Pakistan, 252

prevalence of serum paraprotein bands, (A2) 5

prevalence of vertebral fractures in patients, treated with corticosteroids, (A1) 73

and primary selective IgA deficiency, (A2) 3

prognostic factors in. Comparative study of two subsets of patients according to severity of articular damage, 529 prognostic markers of activity and damage: why clinical trials and

inception cohort studies indicate more favourable outcomes than studies of patients with established disease, (E) 196

prospective, placebo-controlled study of single and double-blind food challenges after dietary manipulation therapy, (AI) 94 prostanoid modulation of synovial antigen-specific CD4+ T-cell

cytotoxic function, 113 proteinaceous lymphadenopathy-case report and review of literature,

1087 pulmonary function in treated with low dose methotrexate: a long-

itudinal study, (A1) 100 radiological progression during a randomised controlled trial of

prednisolone, (A1) 80 randomized controlled trial of four different exercise programs, (A1) 93

recent-onset, short-term outcomes, (S2) 79

relation of treatment to remission, relapse, progression, (A2) 18 relationship between synovitis and erosions, (A1) 105

relative importance of pain and X-ray damage to disability in longstanding, (A2) 12

repeated dose of an engineered anti-TNFα antibody CDP571, (A1) 81 responsiveness of Keitel functional index compared with laboratory measures of disease activity, 141

and rheumatic diseases amongst Pakistanis in London and Karachi, (A1) 33

rheumatoid factor associated with a secretory component, 236 routine follow up: what dictates periodicity of review?, (A2) 12

second-line drug experience of patients receiving lower limb arthroplasty, (A2) 20

serological effects following treatment with an engineered human anti-TNFα antibody CDP571, (A1) 101 shape changes in synovial fibroblasts, (A1) 145

shared care between hospital and general practice: an audit of diseasemodifying drug monitoring, 673

slow acting drug therapy: treatment duration and side effect profile, (A1) 103

small joint synovitis: should it be assessed separately, 51

specificity of antiperinuclear factor for in rheumatoid factor-positive sera, 716

stability of health status measurement, 1162

study of the role of parvovirus B19, 809

subchondral cysts as a favourable prognostic sign, (A1) 104

sulphasalazine: mechanism of action, (S2) 7 TCRBV6S1*3 null allele may confer additional risk in HLA shared epitope positive individual, (A1) 141

therapeutic effects of an engineered human anti-tumour necrosis factor alpha antibody (CDP571), 334

TNF Nco-1 RFLP is not a predictive risk factor, (A1) 140

total and free methotrexate pharmacokinetics, with and without piroxicam, 421

urinary tract infection. A predictable problem?, (A1) 85

use of a contrast enhanced MRI in the assessment of therapeutic response to a disease-modifying antirheumatic drug, 956 and vascular risk factors, (A2) 7

what quality of life assessments should be used?, (A2) 11

why should patients recently recruited to sulphasalazine studies have milder disease than those enrolled over the previous decade? Does this trend influence outcome?, (A1) 98

wrist predominance in seronegative: fact, fiction or natural history, (A2)

Rheumatoid associated pyoderma gangrenosum, cyclosporin A in treatment, (L) 182

Rheumatoid factor

associated cross reactive idiotypes in patients with tropical infections,

associated with a secretory component in rheumatoid arthritis, 236 detection in a 10 year follow-up study of multicase rheumatoid families, (A1) 88

false positive IgA and IgG isotypes in infectious diseases, (A1) 109 IgA and IgM, raised levels in patients with Lyme arthritis and neuroborrelosis, (A1) 110

prognostic value of quantitative measurement in early rheumatoid arthritis, 1146

test, high frequency of parvovirus B19 in serum samples, (A2) 33 Rheumatoid foot, planovalgus—is tibialis posterior tendon rupture a factor?, 645

Rheumatoid joints, T-cell clonality in synovial fluid and after culture in interleukin-2, 232

Rheumatoid knee synovitis, gadolinium-DTPA enhanced magnetic resonance imaging: clinical, laboratory and histological correlation, (A1) 75

generalized pruritis in-consider all possible causes, (L) 295 surgery to the cervical spine, 1078

Rheumatoid vasculitis, isolated nail fold—a benign condition?, (A1) 68 Rheumatological complications of bone marrow transplant, (A1) 120 Rheumatologists, bush rheumatology or rheumatologist gone bush?, 785

disease outcomes, (E) 704

education: what do general practitioners want?, (A2) 49

RHODES C See Naughton D

Rheumatology

Rhupus, clinical and serological uniform condition, (A1) 18

Ribonucleosides, determination of urinary in healthy individuals and those with rheumatic disease, (A1) 59

RICHARDS AJ. How does neurophysiological assessment influence the management and outcome of patients with carpal tunnel syndrome?, (L) 581

RIDDOCH S See Eastmond CJ

RIDLEY N See Murray JG

RIDWAY JC See Howarth S

RIGBY S, PRICE EJ, VENABLES PJW, WALKER JM, MAINI RN. Asymptomatic mothers of babies with congenital heart block with features of Sjögren's syndrome, (A1) 19

RILEY GP, HARRALL RL, CAWSTON TE, HAZLEMAN BL. Basic calcium phosphate salts and supraspinatus tendon degeneration, (A1) 41

RILEY GP See also Bigg HF RING EFJ See Elvins DM

Risk

factors for a patient with spondylarthropathy to develop ankylosing spondylitis or inflammatory bowel disease, (A1) 116

of osteoarthritis associated with running: a radiological survey of female ex-athletes and population controls, (A1) 70

of vertebral fracture in DISH, (A1) 127

RNA polymerase-II large subunit, two distinct sites are recognized by scleroderma autoantibodies but one site is unique to the phosphorylated (IIO) form, (A1) 31

ROBB-NICHOLSON C See Daltroy LH

ROBERTS R, SMITH GW, HALL C, NUKI G. Luteinised unruptured follicles in women with inflammatory arthritis taking NSAIDs, (A2) 22

ROBERTSON DAF See McFarlane XA

ROBERTSON J See Rees DHE

ROBINS RA See Pallis M

ROBINS S See Eastmond CJ

ROBINSON C, DASGUPTA B, CORRIGALL V, YOUNG S, PANAYI G. Deoxycoformycin, a selective lymphocyte depleting drug, induces flare of rheumatoid arthritis, (A1) 102

ROBINSON MA See Kay RA

RODRIGUEZ ALBAN M See Rodriguez de la Serna A

RODRÍGUEZ SANCHEZ JL See Rodríguez de la Serna A

RODRÍGUEZ-MAHOU M See López-Longo FJ

RODRIGUEZ-VALVERDE V See Gonzalez-Crespo MR

RODRÍGUEZ DE LA SERNA, RODRÍGUEZ ALBÁN M, RODRÍGUEZ SANCHEZ JL. Rhupus, clinical and serological uniform condition, (A1) 18

ROGERS J. WATT I. DIEPPE P. The relationship between osteophytosis and osteoarthritis in joints, (A1) 132

ROIVAINEN A, ISOMÄKI P, NIKKARI S, SAARIO R, VUORI K, TOIVANEN P. Oncogene expression in synovial fluid cells in reactive and early rheumatoid arthritis: a brief report, 805

ROLDAN MR. Non-Hodgkin's lymphoma presenting as polyarthritis, (L) 182 Role

of IL-4 and IL-6 in IL-1-dependent cartilage matrix degradation, 101 of prophylactic antibiotics in the prevention of late infection of prosthetic joints. Results of a questionnaire and review of the literature, 380

ROLFE P See Cheung NT; Dziedzic K; Lyttle K

ROMANOWSKI C See Nisar M

ROMANOWSKI CAJ, NISAR M, NAKIELNY RA. Atlanto-occipital subluxation in rheumatoid arthritis demonstrated by magnetic resonance imaging,

ROONEY M, DAVID J, SYMONS J, DI GIOVINE F, VARSANI H, WOO P. Inflammatory cytokine responses in juvenile chronic arthritis, 454

ROONEY M See also Murray JG

RORDORF C See Seitz M

ROSE LM, LATCHMAN DS, ISENBERG DA. BCL-2 expression is unaltered in unfractionated peripheral blood mononuclear cells in patients with systemic lupus erythematosus, 316

ROSMAN A. ATSUMI T. KHAMASHTA MA, AMES PRJ, HUGHES GRV. Development of systemic lupus erythematosus after chemotherapy and radiotherapy for malignant thymoma, 1175

Ross DA, Boylston A, Carney T, Goodacre JA. Peripheral blood T cell receptor $V\beta$ family expression in multicase rheumatoid arthritis families, (A1) 141

Routine follow up of RA patients: what dictates periodicity of review?, (A2) 12

ROWLEY DI. Surgery to the ankle, 854

ROZMAN B See Jevtic V

Running, risk of osteoarthritis associated with: a radiological survey of female ex-athletes and population controls, (A1) 70

RUSINOVA M See Tilley BC

RUSSELL AS See Avina-Zubieta JA

RUSSELL G, FISHER J, CLARKE S, JONES PW, CHEUNG NT, MATTEY DL, HASSELL AB, DAWES PT. Vascular risk factors and rheumatoid arthritis, (A2)7

RUSSELL T See McRorie ER

RYAN MJ See Kavanagh R RYAN PJ See Chikanza IC; Dolan AL

RYAN R See McGonagle D

RYAN S See Carpenter EH

RYDER C See Donn RP

STEUER A. A patient with rheumatoid arthritis and microscopic polyarteritis, (L) 1185

S-oxidation status comparison in Asian and Caucasian rheumatoid arthritis patients. (A2) 19

SAADALLA-NAZHAT RA See Nazhat NB

SAKLATVALA J See Borg AA; Cheung NT SAARIO R See Roivainen A

Safety of hydroxychloroquine in lupus pregnancy: experience is ? pregnancies, (A1) 14

SAHINOGLU T. BHATT B. ABBOT SE. BLAKE DR. STEVENS CR. Synovial microvascular endothelial cells in culture express active xanthine oxidase, (A1) 51

SAKLATVALA J See Hassell AB

SALAHELDIN IB See Troughton PR

SALIH AM, NIXON NB, HAWKINS CP, DAWES PT, MATTEY D. Anti ganglioside antibodies in patients with rheumatoid arthritis and peripheral neuropathy, (A2) 3 SALIH AM See also Nixon NB

Salivary glands, cytokine mRNA expression from patients with primary Sjögren's syndrome, 326

Salmonella pyomyositis in patients with the human immunodeficiency virus, 568

Salmonella—a forgotten infection in seronegative oligoarthritis?, (A2) 32 SALTER D See Jobanputra P

SALTER DM See Wright M

SAMAD SA See Segasothy M

SAMBROOK PN, JONES G. Corticosteroid osteoporosis, 8

SAMBROOK PN See also Keen RW

SANCHEZ E See López-Longo FJ

SANCHEZ GUIJO P See Tinahones FJ

SANDOVAL DM, ALARCÓN GS, MORGAN SL. Adverse events in methotrexatetreated rheumatoid arthritis patients, (S2) 49

SANT S See Flanagan FL

SANT SM See Doran M

SANY J See Combe B; Jorgensen C

SARATZIS N See Kitas GD

SARAUX A, DE SAINT PIERRE V, BARON D, VALLS I, KOREICHI A, YOUINOU P, LE GOFF P. The antiperinuclear factor in spondylarthropathies, 912

Sarcoid arthritis, acute: occurrence, seasonal onset, clinical features and outcome, 45

Sarcoidosis, low-dose methotrexate: an effective corticosteroid-sparing agent in the musculoskeletal manifestations, 642

SATO S See Fujimoto M; Ihn H

SATOH M, MIYAZAKI K, MIMORI T, AKIZUKI M, ICHIKAWA Y, HOMMA M, AJMANI AK, REEVES WH. Changing autoantibody profiles with variable clinical manifestations in a patient with relapsing systemic systemic lupus erythematosus and polymyositis, 915

SATOH M, TOKUCHIRA M, HAMA N, HIRAKATA M, KUWANA M, AKIZUKI M, ICHIKAWA Y, OGAWA S, HOMMA M. Massive pericardial effusion in

scleroderma: a review of five cases, 564
SAUNDERS A See Mulherin D

SAVAGE COS See Luqmani RA

Savage, Oswald Arthur, obituary, 591

SAVILL J. Apoptosis: from worm to wonder drug, (E) 95

SAXNE T See Sharif M

SCHAEVERBEKE T. Is remitting seronegative symmetrical synovitis with pitting oedema (RS3PE syndrome) associated with HLA-A2?, (L) 889

SCHALLER JG See Tucker LB SCHANTZ A See Choy EHS

SCHATTEMAN G See Mielants H

SCHAUFELBERGER C, BENGTSSON B-A, ANDERSSON R. Epidemiology and mortality in 220 patients with polymyalgia rheumatica, 261
SCHEJA A, WOLLHEIM FA, ÅKESSON A. The predictive value of von

Willebrand factor and capillary microscopy in systemic sclerosis (scleroderma)-a prospective study, (A1) 27

SCHOLLUM J See Croft P

SCHOLTEN R See Lycklama à Nijeholt GJ

SCHUWIRTH LWT See Blaauw AAM

SCHWARTZBERG M. Lyme borrelosis presenting as a polymyalgia rheumaticalike syndrome, (L) 392

Sciatica, lumbar epidural steroid injection: who benefits?, (A1) 40

Scientific interest growth in rheumatology, (L) 796

Scintigraphic findings in synovial chondromatosis at presentation, (L) 687 Scintigraphic localization of steroid injection site for plantar fasciitis, (A1) 42 Scleroderma

arthritis in, 831

autoantibodies, two distinct sites on the RNA polymerase-II large subunit are recognized but one site is unique to the phosphorylated (IIO) form, (A1) 31

and control fibroblast phenotype modulation by co-culture with human endothelial cells, (A1) 66

differences between adult and childhood onset, (A1) 29

diltiazem induces remission of calcinosis, 576

evidence of omeprazole-induced small bowel bacterial overgrowth, 976 limited, subcutaneous and periarticular calcification—a variant of?, (A2)

lung fibroblasts, endothelial cell conditioned medium promotes contraction of collagen gels, (A1) 31

malabsorption caused by coeliac disease, 858

massive pericardial effusion in: a review of five cases, 564

neutral matrix metalloproteinases, (A1) 29

skin score correlation to change in soluble ICAM-1 levels reflects endothelial activation, (A1) 28

survival following onset. Results from a UK clinic population, (A1) 27 SCOTT D See Lassere M

SCOTT DGI. Antigen specificity of ANCA in systemic vasculitis, (L) 394 SCOTT DGI See also Bankhead CR; Brennan P; Carruthers DM; Harrison BJ; Jones PBB; Watts RA

SCOTT DL, BERRY H, HUNTER J, LINK GROUP. Tiaprofenic acid, indomethacin and placebo in knee osteoarthritis: the long term outcome of therapy, (A2) 44

SCOTT DL, BERRY H, WILLIAMS P, KHAN FM, EMERY P, DAYMOND TJ. Perforations, peptic ulceration and bleeding on NSAIDs. Two randomised controlled trials of nabumetone, diclofenac SR and piroxicam in the UK. (A1) 95

SCOTT DL, HOUSSIEN DA, LAASONEN L. Proposed modification to Larsen's scoring methods for hand and wrist radiographs, 56

SCOTT DL, LONG F, SILMAN A. Disease outcomes in rheumatology, (E) 704 SCOTT DL See also Comer M; Houssein DA; Houssien DA; Mahmud T; Rafi S; Smith EC; Storey GO

Screening

for chloroquine retinopathy, (A1) 18

of synovial fluid samples from patients with undifferentiated oligo-arthritis for bacterial antigens using PCR, (A1) 116

Second-line drug experience of patients with rheumatoid arthritis receiving lower limb arthroplasty, (A2) 20

SEED MP, BOWDEN AR, PARKER FP, JOHNS S, CURNOCK A, GARDNER CR. Chondrodestructive and chondrosparing properties of non-steroidal anti-inflammatory drugs in a model of destructive mono-articular arthritis, (A2) 15

SEEDHOM BB See Beevers DJ; Shepherd DET

SEGASOTHY M, CHIN GL, SIA KK, ZULFIQAR A, SAMAD SA. Chronic nephrotoxicity in the treatment of arthritis, 162 SEGNINI G See Berrettini S

SEIFERT B See Stoll T

SEITZ M, LOETSCHER P, DEWLAD B, TOWBIN H, RORDORF C, GALLATI H,
BAGGIOLINI M, GERBER NJ. Methotrexate action in rheumatoid arthritis: stimulation of cytokine inhibition and inhibition of chemokine production by peripheral blood mononuclear cells, 602

SEKHARAN NG See Uppal SS

Selective accumulation of cutaneous lymphocyte antigen positive lymphocytes in the skin but not in the joint of inflammatory arthropathies, (A1)

Self management and personal illness models of arthritis, (A1) 90

comparison of the validity and reliability of self-reported articular indices, 760

rheumatoid arthritis patients cannot accurately report signs of inflammatory activity, 547

SELLAMI S See Claudepierre P

SELLARI-FRANCESCHINI S See Berrettini S

Sensitivity and specificity of different diagnostic criteria for Behçet's disease according to the latent class approach, 932

Septic arthritis in Amsterdam: potential targets for prevention, (A1) 113

Sequence analysis of four human monoclonal IgG anti-cardiolipin antibodies,

Serial changes in bone mineral density in polymyalgia rheumatica, (A1) 74 SERIOLO B See Cutolo M

Serological diagnosis of Lyme disease, (L) 992

Serological effects following treatment with an engineered human anti-TNF α antibody CDP571, in patients with rheumatoid arthritis, (A1) 101

Serological joint markers—a comparison of serum proMMP-3, proMMP-1 and disease activity in RA, (A1) 60

Seronegative arthritis mailbase discussion group, (L) 1185

Serum electrolytes and bone density: the Chingford study, (A1) 129

Serum urate acute gout, (A2) 40

SETOAIN J See Pons F

SEWELL HF See Hodges E

Sexually acquired arthritis in nineteenth century London, (A2) 31 SHADFORTH MF See Carpenter EH; Dawes PT; Griffiths SM; Hassell AB; Mayer B

SHAH V See Nash MC

SHAHID MS, GOSLING J, SHEPSTONE L, WATT I, DIEPPE P. Dual energy X-ray absorptiometry of the knee joint, (A1) 122

Shape changes in synovial fibroblasts from RA patients, (A1) 145

SHAPLEY M See Fisher M

Shared care between hospital and general practice: an audit of diseasemodifying drug monitoring in rheumatoid arthritis, 673

SHARIF M, SAXNE T, SHEPSTONE L, KIRWAN JR, ELSON CJ, HEINEGARD D, DIEPPE PA. Relationship between serum cartilage oligomeric matrix protein levels and disease progression in osteoarthritis of the knee joint,

SHARIF M See also Dodd J; Hilman S

SHEERAN T See Glennon P

SHEERAN TP. Hypocomplementaemia, C3 nephritic factor and type III mesangiocapillary glomerulonephritis progressing to systemic lupus erythematosus, (L) 90

SHEERAN TP See also Birrell FN

SHELLEY M See Hay EM

SHEPHERD DET, SEEDHOM BB. Measurement of articular cartilage compressive modulus under physiological loading conditions, (A2) 48

SHEPSTONE L See Hewlett SE; Lim K; Shahid MS; Sharif M

SHIMOZUMA M See Ihn H

SHINGU M, FUJIKAWA Y, WADA T, NONAKA S, NOBUNAGA M. Increased IL-1 receptor antagonist (IL-1ra) production and decreased IL-1β/IL-1ra ratio in mononuclear cells from rheumatoid arthritis patients, 24

SHINGU M, MIYAUCHI S, NAGAI Y, YASUTAKE C, HORIE K. The role of IL-4 and IL-6 in IL-1-dependent cartilage matrix degradation, 101

SHIROKY JB. Combination of sulphasalazine and methotrexate in the management of rheumatoid arthritis-view based on personal clinical experience, (S2) 109

SHIWEN X, DENTON CP, PEARSON JD, DU BOIS RM, BLACK CM. Endothelial cell conditioned medium promotes contraction of collagen gels by scleroderma lung fibroblasts, (A1) 31

SHIWEN X See also Denton CP

Short-term outcomes in recent-onset rheumatoid arthritis, (S2) 79

Shortly after anterior cruciate ligament transection dogs develop characteristics similar to human moderate knee osteoarthritis, (A1) 135

Should close relatives of patients with osteoporosis be investigated?, (A1) 73 Should tests for proteinuria be included in the monitoring schedule of sulphasalazine?, 790

movement measurement using FASTRAK 3-D tracking system, (A2) 47 pain in a community-based rheumatology clinic, 440 replacement-surgery to the shoulder, 653

symptoms, comparison of the accuracy of steroid placement with clinical outcome, (A1) 41

SIA KK See Segasothy M

Sialylation, α2.3 and α2.6 linked and Lewis X expression on serum IgG in rheumatoid arthritis, (A1) 63

Sialyltransferase α2,6 and β1,4 galactosyltransferase activity are simultaneously reduced in CD19+ peripheral blood lymphocytes in rheumatoid arthritis (A1) 62

Signal transduction pathways in human chondrocytes activated by intermittent pressurisation, (A1) 58

SILLS J. The BPA report on paediatric rheumatology, 60

SILLS JA See Symmons DPM

SILMAN A See Ali-Gombe A; Brennan P; Croft P; Jones M; Kiss Cs; Scott DL

SILMAN AJ, HOWARD Y, BRYAN C, BRENNAN P, BLACK CM. Survival following the onset of scleroderma. Results from a UK clinic population, (A1) 27

SILMAN AJ See also Bankhead CR; Brennan P; Chambers H; Croft PR; Ferry S; Harrison BJ; Macgregor AJ; Newman J

SILVERMAN B See Cawston TE SIMEON CP See Marsal S

SIMMONDS P See Jobanputra P

SIMONITSCH I See Koeller M

SIMPSON E See Hogarth MB SIMPSON RJ See Raja KB SIMS TJ See Bailey AJ

SINGH G See Uppal SS

Single photon emission computerised tomography identifies pain sites in ankylosing spondylitis, (A1) 126

SIREN M-K See Julkunen H

SITUNAYAKE RD See Gordon C; Griffiths B; Yeo LMW

Sixth European Regional Conference of Rehabilitation International, 283 Sjögren's syndrome

abducent and trochlear palsies in, (L) 484

asymptomatic mothers of babies with congenital heart block with features of, (A1) 19

cytokine mRNA expression in the labial salivary gland, 326

do patients with SLE, SLE with Sjögren's syndrome and primary Sjögren's syndrome differ in damage or function?, (A2) 26

hypergammaglobulinaemic primary, mapping the genetic association with the TRCB gene complex, (A1) 77

interferon-gamma and epithelial cell activation, 226 maternal HLA antigens and antibodies to SS-A/Ro and SS-B/La. Comparison with systemic lupus erythematosus and primary Sjögren's syndrome, 901

primary, assessment of in patients with dry eyes, (A2) 25

VIP in salivary glands, (L) 891

SKELLY M See Mulherin D

hyper-reactivity in patients with benign intracranial hypertension as an early manifestation of Behçet's disease, (L) 184

manifestations of Marfan syndrome: joint and skin manifestations are prevalent and correlated, 126

ultrasound thickness measurement in systemic sclerosis, 535

SKOPOULI FN See BOUMBA D

SLATER H See Mattey DL SLAVOTINEK A See Hardwick LJ

Sleen

electroencephalographic pattern clustering in patients with the fibromyalgia syndrome, 1151

intensity in fibromyalgia: focus on the microstructure of the sleep process, 629

low prevalence of fibromyalgia syndrome in patients attending a sleep disorders clinic, (A1) 38

studies in fibromyalgia; techniques, clinical significance and future directions, (E) 897

SLINGSBY JH See Hogarth MB

Slow acting anti-rheumatic drugs, are they monitored too often? An audit of current clinical practice, 966

Slow acting drug therapy of rheumatoid arthritis: treatment duration and side effect profile, (A1) 103

Small bowel, characteristics of ulcers of induced by non-steroidal antiinflammatory drugs in the rat: implications for clinical practice, 727

Small bowel bacterial overgrowth evidence of imeprazole-induced in patients with scleroderma, 976 and omeprazole in patients with systemic sclerosis. (A1) 67

in systemic sclerosis: detection using direct and indirect methods and treatment outcome, 265

Small joint synovitis in rheumatoid arthritis: should it be assessed separately, 51

Small sensory fibres, consequences of sustained activation on pain sensation and neurogenic inflammation, (A1) 53

SMEDEDÅRD G, BJÖRK J. SULPHASALAZINE: MECHANISM OF ACTION IN RHEUMA-TOID ARTHRITIS, (S2) 7

SMITH DA See Carpenter EH

SMITH EC, WILLIAMS S, WASSIF W, SCOTT DL. Proton-NMR spectroscopy and the diagnosis of polymyositis and dermatomyositis, (A2) 27
SMITH EC, WILLIAMS S, BERRY H, SCOTT DL. Polymyositis patients show

consistent changes on 1H-NMR spectroscopy, (A1) 125

SMITH GW, SMITH NM, UTTLEY WS, LUQMANI RA. Case report: dermatomyositis-scleroderma overlap in childhood, (A2) 54

SMITH GW See also Roberts R

SMITH JL See Hodges E; Wong WM Sмітн M See Barry R; Kavanagh R

SMITH NM See Smith GW

SMITH PR See Kalla AA

SMITH R. Idiopathic juvenile osteoporosis: experience of twenty-one patients,

SMITH RW. Polyarthritis masquerading as the bends, (A2) 53

SMITS F See Blaauw AAM

Smoking effects on outcome of ankylosing spondylitis, (A2) 35

SMOLEN J See Erlacher L; Wylie G

SMYTHE HA. Studies of sleep in fibromyalgia; techniques, clinical significance and future directions, (E) 897

SNAITH M See Crawford F SNAPE SD See Raja KB

SNOWDON N See Ollier WER

Soft tissue and intra-articular tissue injections: a survey of current practice, 449

SOHAIBANI MO See Al Rikabi AC

SOLTYS AJ, ALAVI A, AXFORD JS. Galactosyltransferase isoenzyme changes occur in rheumatoid arthritis, (A1) 61

SÖLTZ-SZÖTS J See Erlacher L

Soluble FAS is increased in patients with SLE, (A1) 5

Soluble immunoglobulin inhibits FCIII receptor-mediated neutrophil arrest to immune complexes under flow conditions: a possible mechanism of action in treatment of vasculitis, (A1) 24

SOMA Y See Ihn H

SOMASUNDARAM S See Rafi S

Somatisation, irritable bone syndrome—a new concept in?, (A2) 52

SONNTAG K-C. Is interferon alpha a therapy for hepatitis B-associated polyarteritis nodosa, (L) 486

SOOD A, KUMAR A, PANDE I, MALAVIYA AN. Does mixed connective tissue disease exist in India?, 539

SOOD A, PANDE I, KUMAR A, MALAVIYA AN. Does MCTD exist in India?, (A2) 26

SOPER CJ See Collins AJ

SOPWITH AM See Rankin ECC

SORIANO E See WHYTE J SOUTHWOOD T See Donn RP

SPÄTH PJ See Bartók I

Special circulating adhesion molecule levels reflect disease severity in systemic sclerosis, 1048

Specificity of antiperinuclear factor for rheumatoid arthritis in rheumatoid factor-positive sera, 716

SPECT scanning can predict successful facet joint injection, (A1) 71

SPECTOR TD, HARRIS PA, HART DJ, CICUTTINI F, NANDRA D, ETHERINGTON J. Wolman R, Doyle DV. Risk of osteoarthritis associated with running: a radiological survey of female ex-athletes and population controls, (A1) 70

SPECTOR TD See also Alavi A; Arden NK; Dolan AL; Donnelly S; Etherington J; Hart DJ; Keen RW; Lloyd ME

SPEED C See Haslock I

SPEED CA, FOOT V, HASLOCK I, FORDHAM JN. A comparison of intravenous, intramuscular and oral pulse steroid therapies in the management of active rheumatoid arthritis, (A2) 19

Spiking of the tibial tubercles, does it predict osteoarthritis of the knee?, (A1)

Spinal osteoarthritis and osteoporosis: evidence for a negative association independent of obesity, (A1) 74

Spinal problems, manipulative physiotherapy for in primary care: outcomes of care, 1070

SPITZER WO, HAGGERTY JL, BERKSON L, DAVIS W, PALMER W, TAMBLYN R, LAPRISE R, FAITH JM, ELMORE JG, HORWITZ RI. Continuing occurrence of eosinophilia myalgia syndrome in Canada, 246

Splenic infarction, symptomatic complicating adult Kawasaki disease, (L) 579 Spondylarthritis, seronegative, interleukin-2 in and in psoriatic arthritis but not in rheumatoid arthritis synovium, (A1) 151

Spondylarthropathy active, elevated serum nitrate concentrations, 616

the antiperinuclear factor, 912

predictive factors of severity of spondylarthropathy in North Africa, 1139

profile of HLA-B27-related unclassifiable seronegative in females and its comparison with the profile in males, 137

seronegative, different abdominal scintigraphy pattern from patients with ulcerative colitis and Crohn's disease, 946

Spondylitis-like and symmetric polyarthralgia associated with gemfibrozil therapy, (L) 692

Spouse, patient's, effects of rheumatoid arthritis on, (A2) 10

SRINIVASAN A, AMOS M, WEBLEY M. The effects of joint washout and steroid injection compared with either joint washout or steroid injection alone in rheumatoid knee effusion, 771

SS-A/La antibodies, maternal HLA antigens and antibodies to SS-A/Ro and SS-B/La. Comparison with systemic lupus erythematosus and primary Sjögren's syndrome, 901

SS-A/Ro antibodies, maternal HLA antigens and antibodies to SS-A/Ro and SS-B/La. Comparison with systemic lupus erythematosus and primary Sjögren's syndrome, 901

Stability

of health status measurement in rheumatoid arthritis, 1162

of pyridinium crosslinks to photolysis, (A1) 54

STALLA F See Gattorno M STANEK G See Erlacher L

STANESCU R See Bailey AJ

STANESCU V See Bailey AJ

Stanford Health Assessment Questionnaire: ordinal or interval scale?, (A2) 51

STANKOVIC A See Pejovic M

STANWORTH DR See Close DR Staphylococcal arthritis in human immunodeficiency virus infection, (L) 397

STARR M See Hudson N

STARR MR See Hudson N

STAUD R. Brachial plexitis in a patient with acute parvovirus B19 infection, (L) 480

STEAD H See Wright D

STEEN IN See Bradshaw C

STEERE AC See Rees DHE

STEERS G See Davies EJ

STEINER A See Jones SD; Koh WH

STEINER CW See Koeller M

STENSTAD T, HUSBY G. Therapeutic effect of brefeldin on experimental reactive AA amyloidosis, (A1) 118

STENSTAD T, MAGNUS JH, HUSBY G. Proteoglycans associated with experimental reactive AA amyloidosis, (A1) 118

STEPHENSON AM See McDonagh JE

Steroid injection

effects and joint washout compared with either joint washout or steroid injection alone in rheumatoid knee effusion, 771 lumbar epidural in sciatica: who benefits?, (A1) 40

scintigraphic localization of site for plantar fasciitis, (A1) 42

Steroid myopathy induced by epidural triamcinolone injection, 385 Steroid placement, comparison of the accuracy with clinical outcome in

patients with shoulder symptoms, (A1) 41 Steroid therapy

comparison of intravenous, intramuscular and oral pulse in the management of active rheumatoid arthritis, (A2) 19

long-term, effect of and of systemic lupus erythematosus on bone mass in pre-menopausal women, 742

oral as bridge therapy in rheumatoid arthritis patients starting with parenteral gold. A randomized double-blind placebo-controlled trial, 347

STEVENS CR See Abbot SE; Sahinoglu T; Zhang Z

STEVENS RJ, HUGHES RA. The aetiopathogenesis of giant cell arteritis, 960 STEVENS RJ. Serological diagnosis of Lyme disease, (L) 992

STEWART I See Howarth S

STEWART LJ See Whyte J

STICHTENOTH DO, WOLLENHAUPT J, ANDERSONE D, ZEIDLER H, FRÖLICH JC. Elevated serum nitrate concentrations in active spondylarthropathies, 616

Still's disease

adult onset

destructive lymphadenopathy and T-lymphocyte activation, 978 regional audit, (A2) 22

adult onset in Northern India: comparison with juvenile onset Still's disease, 429

STOLL T, HOFER HO, HUBER E, SEIFERT B, MICHEL BA. A new quantitative and simple muscle strength measurement: interobserver reliability in healthy subjects, (A1) 20

STOLL T See also Gordon C; Sutcliffe N

Stomatitis, management of oral complications of disease modifying drugs in rheumatoid arthritis, (A2) 20

STOREY GO, SCOTT DL. Sexually acquired arthritis in nineteenth century London, (A2) 31

STRANGE R See Hassell A; Ollier WER

STRATTON R See Dolan AL

STREET P See Darlington LG

Stress protein 32kDa expression in arthritic synovia, (A1) 47

STRÖM H. Lymphocyte reactivity to Clostridium perfringens antigens and toxoid in patients with arthritis, (L) 1095

Strontium intestinal absorption and axial bone mineral density, (A1) 75

STUCKI G, STUCKI S, BRÜHLMANN P, MAUS S, MICHEL BA. Comparison of the validity and reliability of self-reported articular indices, 760 STUCKI S See Stucki G Studies of sleep in fibromyalgia; techniques, clinical significance and future

directions, (E) 897 Study of the role of parvovirus B19 in rheumatoid arthritis, 809

STURGESS A See Helsloot J

STURROCK RD See Canvin JMG; Craig JP; Griffiths ID; Hall MG; Karim SN; McInnes IB; Munro R; Neilly JB

SUAREZ-ALMAZOR ME See Avina-Zubieta JA

Subchondral cysts as a favourable prognostic sign in rheumatoid arthritis, (A1) 104

Subcutaneous and periarticular calcification-a variant of limited scleroderma?, (A2) 53

Substance P, FCIII receptor-mediated arrest under flow conditions is

enhanced by, (A1) 52 Subungual splinter haemorrhages, prevalence and clinical significance in patients with an antiphospholipid syndrome, (A1) 16

Successful management of reactive haemophagocytic syndrome in systemiconset juvenile chronic arthritis, (L) 888 SUEOKA N See Kuroki S

SULLI A See Cutolo M

Sulphapyridine, sulphasalazine or 5-aminosalicylic acid-which is the active moiety in rheumatoid arthritis?, (S2) 16

Sulphasalazine

clinical efficacy-a review, (S2) 35

in combination with methotrexate, effects on the production and circulating concentrations of cytokines and their antagonists, 747 does low dose desensitization affect the toxicity in the treatment of

rheumatoid arthritis?, (A1) 97

effect on neutrophil superoxide protection in RA, (A2) 14

history of the use of in rheumatology, (S2) 2

influence on the evolution of gut inflammation and locomotor manifestations in the spondylarthropathies, (A1) 115

long-term usage and side-effect profile in rheumatoid arthritis, (S2) 40 mechanism of action in rheumatoid arthritis, (S2) 7 and methotrexate

antirheumatic agents share an anti-inflammatory mechanism. (S2)

combination in the management of rheumatoid arthritis-view based on personal clinical experience, (S2) 109

combination in rheumatoid arthritis: early clinical impressions, (S2)

penicillamine, audit of patients taking them individually, sequentially and concurrently in inflammatory arthritis, (A2) 17

reduction in cell adhesion molecule levels following treatment in patients with rheumatoid arthritis, (A1) 98

should tests for proteinuria be included in the monitoring schedule?, 790 sulphapyridine or 5-aminosalicylic acid-which is the active moiety in rheumatoid arthritis?, (S2) 16

suppression of proteinuria and prolongation of survival in MRL-lpr/lpr mice. (L) 85

survival on-results from the Norfolk Arthritis Register, (A2) 14 why should rheumatoid arthritis patients recently recruited to studies have milder disease than those enrolled over the previous decade? Does this trend influence outcome?. (A1) 98

Sulphasalazine-induced autoimmune abnormalities in patients with rheumatic disease, 435, (L) 1097

Sulphasalazine-induced lupus in psoriatic arthritis, 383

Sulphoxidation capacity variation in systemic lupus erythematosus patients and healthy controls of different ethnic origins, (A1) 6

SULTANA S See Hameed K

SUN K-H See Hsieh S-C

Suppression of proteinuria and prolongation of survival in MRL-lpr/lpr mice treated with sulphasalazine, (L) 85

Supraspinatus tendon degeneration and basic calcium phosphate salts, (A1)

Surgery

outcome to stabilise the cervical spine in rheumatoid arthritis-pain, neurological status and patient satisfaction, (A1) 92 to the ankle, 854

to the shoulder-shoulder replacement, 653

Surgical intervention, nature in patients with ankylosing spondylitis, (A1) 115 SURIS X See Peris P

Survival

following the onset of scleroderma. Results from a UK clinic population,

on sulphasalazine--results from the Norfolk Arthritis Register, (A2) 14 Sustained-release transdermal glyceryl trinitrate patches as a treatment for primary and secondary Raynaud's phenomenon, 636

SUTCLIFFE N, STOLL T, MALIK J, PYKE S, ISENBERG DA. Do patients with SLE, SLE with Sjögren's syndrome and primary Sjögren's syndrome differ in damage or function?, (A2) 26

SVENDSEN L See Drewes AM SWABY DSA See Yeo LMW

SWANINK CMA See Griep EN

SWANNELL A See Bacon PA

SWINBURN HM See Dolan AL

SYED A See Hameed K

SYKES B See Loughlin J SYKES H See MacDonald AG

Symmetrical compartmental involvement in knee osteoarthritis, (A1) 134

SYMMONS D See Jones M; Jones PBB

SYMMONS DPM. Measuring outcome in rheumatoid arthritis-which measures are suitable for routine clinical use?. (E) 802

SYMMONS DPM, SILLS JA, DAVIS SM. The incidence of juvenile dermatomyositis: results from a nation-wide study, 732

SYMMONS DPM See also Bankhead CR; Brennan P; Harrison BJ; Thomson W; Watts RA

SYMONS J See Rooney M

Symptomatic enlarged iliopsoas bursae in the presence of a normal plain hip radiograph, 365

Symptomatic splenic infarction complicating adult Kawasaki disease, (L) 579 Syndesmophyte formation effect on lumbar spine bone density in patients with ankylosing spondylitis, (A1) 117

Synovial chondromatosis, scintigraphic findings at presentation, (L) 687 Synovial fibroblastic cells, interleukin-6 (IL-6) induces the proliferation in the presence of soluble IL-6 receptor, 321 Synovial fibroblasts, shape changes from RA patients, (A1) 145

Synovial fluid

glycolipid antibodies in rheumatoid arthritis-synovial neuropathy or synovial damage, (A2) 4

sample screening from patients with undifferentiated oligoarthritis for bacterial antigens using PCR, (A1) 116

Synovial lining-associated epitope 67: tissue distribution and identification by Western blotting, (A1) 149

Synovial membrane, IL-1 and IL-1ra production by monocyte/macrophage in rheumatoid arthritis, (A1) 150

Synovial microvascular endothelial cells

in culture express active xanthine oxidase, (A1) 51

exhibit enhanced shear rate dependent leucocyte adhesion in vitro, (A1)

Synovial permeability, determination of apparent in the knee joint of patients suffering from osteoarthritis and rheumatoid arthritis, 520 Synovial targeting, comparable co-expression of VCAM-1* and DAF* on

synovial intimal cells and follicular dendritic cells in rheumatoid arthritis, (A2) 4

Synovial tissue

presence of bacterial flora-derived antigen in macrophages and dendritic cells 1112

volume quantification with gadolinium-DTPA enhanced magnetic resonance, (A1) 125

Synoviocytes, lamellar body secretion: ultrastructural analysis of an unexplored function, 13

Synovitis inflammatory, gut involvement in the pathogenesis, (E) 801

pathological synovial fluids promote apoptosis of peripheral blood and synovial neutrophils: a role in resolution?, (A1) 58

relationship with erosions in rheumatoid arthritis, (A1) 105

remitting seronegative symmetrical with pitting oedema (RS3PE syndrome) association with HLA-A2, (L) 889

small joint in rheumatoid arthritis: should it be assessed separately, 51

arthritic, 32kDa stress protein expression, (A1) 47

macrophage populations in: inter-relationships and correlation with articular damage, (A1) 149 rheumatoid, clonality of immunoglobulin production, (A1) 154

Systemic lupus erythematosus

acquired Brown's syndrome, 1092

adult-and childhood-onset: a comparison of onset, clinical features, serology and outcome, 866

analysis of thrombotic events in antiphospholipid antibody positive lupus patients, (A1) 15

anti-Clq antibodies are associated with hypo-complementaemia, defective opsonisation of gram-positive organisms, and death due to infection. (A1) 10

anti-Sa antibodies, (A1) 9

assessing activity using plasma factor VIII von Willebrand factor antigen, (A1) 8

BCL-2 expression is unaltered in unfractionated peripheral blood mononuclear cells, 316

CD5-positive and CD5-negative plaque-forming cells against poly-Llysine treated sheep erythrocytes, (A1) 6

cerebral venous thrombosis and acquired protein S deficiency: an uncommon cause of headache, 572

characterisation of anti-endothelial cell membrane antibodies, (A1) 11 complement and immune complexes, (A1) 82

damage correlates with disease duration in Afro-Caribbeans and Caucasians but not Asians, (A1) 17

defective spontaneous and bacterial lipopolysaccharide-stimulated production of interleukin-1 receptor antagonist by polymorphonuclear neutrophils, 107

development after chemotherapy and radiotherapy for malignant thymoma, 1175

development of features in patients with primary antiphospholipid syndrome, (A1) 16

patients with SLE, SLE with Sjögren's syndrome and primary Sjögren's syndrome differ in damage or function?, (A2) 26

effect and of long-term steroid therapy on bone mass in pre-menopausal women, 742

high resolution computed tomography of the lung in asymptomatic patients with normal chest radiographs, (A1) 14 homozygosity for glutathione-transferase GSTM1 and GSTT1 null

alleles may confer risk for anti-Ro antibodies, (A1) 5

hypocomplementaemia, C3 nephritic factor and type III mesangiocapillary glomerulonephritis progressing to, (L) 90 increased interleukin-2 receptor expression on peripheral blood lympho-

cytes, (A1) 6 increased levels of antibodies to heat shock proteins with increasing age in MRL/MP-LPR/LPR mice, 610

increased levels of β 2 glycoprotein-1 antigen and β 2 glycoprotein-1 binding antibodies are associated with a history of thromboembolic complications, 1031

magnetic resonance imaging of the brain, 1055

and mannose binding protein in southern Chinese, (A1) 7

mannose-binding protein and complement C4 polymorphism are associated with SLE in a Spanish population, (A1) 78

maternal HLA antigens and antibodies to SS-A/Ro and SS-B/La. Comparison with systemic lupus erythematosus and primary Sjögren's syndrome, 901

measurement of antibodies to reactive oxygen species damaged DNA as a simple marker of disease activity, (A1) 10

menstrual disorders in girls treated with cyclophosphamide, 737 occurrence, nature and distribution of flares in a cohort of patients: a rheumatological view, 257

paediatric, antiphospholipid antibodies in, and in juvenile chronic arthritis and overlap syndromes, 873

patients with both lupus anticoagulant and high-titre anticardiolipin antibodies are at risk for clinical manifestations related to the antiphospholipid syndrome, 873

polymorphism of the heat-shock protein gene HSP70-2, 721

primary nocardial meningitis, 178

primary tuberculous muscle abscess, 1177

quantitative analysis of lymphocyte adhesion molecule deployment, (A1)

relapsing, changing autoantibody profiles with variable clinical manifestations, 915

relative contributions of HLA-DQA and complement C4A loci in determining susceptibility to, 221

soluble FAS is increased in, (A1) 5

thyroid autoimmunity in: the clinical significance of a fluctuating course, 361

urinary retinol binding protein excretion, (A1) 17

variation in sulphoxidation capacity in patients and healthy controls of different ethnic origins, (A1) 6

Systemic rheumatic disease, effectiveness of minimally supervised home aerobic training, 1064 Systemic sclerosis

abnormal anal function, (A1) 26

altered levels of soluble adhesion molecules, 814 autoantibodies to myeloperoxidase, (A2) 24

detection of antiribosomal P protein antibodies, 908

increased incidence of intermittent claudication: a prospective study, (A1) 26

management, (E) 3

neurological involvement, (E) 1007

neurovascular control mechanisms in and in primary Raynaud's phenomenon as assessed by iontophoresis and laser Doppler techniques, (A2) 24

predictive value of von Willebrand factor and capillary microscopy-a prospective study, (A1) 27

and primary Raynaud's phenomenon, capillary measurements by video capillary microscopy, (A1) 25

progressive, phrenicus palsy in, 684 small bowel bacterial overgrowth: detection using direct and indirect methods and treatment outcome, 265

small bowel bacterial overgrowth and omeprazole, (A1) 67

special circulating adhesion molecule levels reflect disease severity, 1048 ultrasound measurement of skin thickness, 535

use of intravenous prostacyclin to treat pulmonary hypertension associated with, (A1) 30

Systemic vasculitis

antigen specificity of ANCA, (L) 394

incidence in the Norwich Health Authority, (A1) 21

increased endothelium derived relaxing factor/nitric oxide production,

intravenous cyclophosphamide in the treatment: a five year follow-up study, (A1) 22

pulse cyclophosphamide therapy, (A1) 21

TELLUS MM. Effective treatment of anti Jo-1 antibody-positive polymyositis with cyclosporin, (L) 1187

T lymphocyte activation and destructive lymphadenopathy in adult-onset Still's disease, 978

T-cell

clonality in synovial fluid from rheumatoid joints and after culture in interleukin-2, 232

immunity, dominant human leukocyte antigen DR4-Dw15 restricted mycobacterial 65 kDa antigen-specific in Chinese patients with rheumatoid arthritis, 498

receptor variable gene segment TCRBV6S1 null allele frequencies in UK juvenile chronic arthritis, (A1) 144

receptor VB family expression in peripheral blood in multicase rheumatoid arthritis families, (A1) 141

response to recombinant topoisomerase-1 is independent of disease and of autoantibody status, (A1) 30

TAAGHOLT SJ See Drewes AM

TAGGART A See Doherty E

TAIT TJ. Wegener's granulomatosis and CREST syndrome, (L) 483

TAKAHASHI M. Receiver operating characteristic analysis of urinary pyridinium cross-links in arthritis, (L) 792

TAKEHARA K See Fujimoto M; Ihn H TAMAKI K See Fujimoto M: Ihn H

TAMBLYN R See Spitzer WO

Targeting endothelial activation in inflammation using 111In-labelled anti-Eselectin monoclonal antibody, (A1) 81

TARGOFF IN See D'Cruz D

Tarsal tunnel syndrome in ankylosing spondylitis, (L) 488

TASIAUX N See Dobbie JW

TAUB NA See D'Cruz D

TAYLOR D See Cheung NT

TAYLOR DJ See Cheung NT

TAYLOR HG See Averns HL

TAYLOR J See Thompson PW TAYLOR M See Kaye SA

TAYLOR P See Peacock DJ

TAYLOR PC See Williams DG

TeR $\Gamma\delta$ lymphocytes, arthropathy, leucopenia and recurrent infection associated with, 978

TCRBV6S1*3 null allele may confer additional RA risk in HLA shared epitope positive individual, (A1) 141

TEALE R See Bacon PA

Tear evaporation rate and osmolality, abnormal in rheumatoid arthritis, (A1) 91

TEBIB JG, LETROUBLON MC, NOEL E, BIENVENU J, BOUVIER M. sIL-2R levels in rheumatoid arthritis: poor correlation with clinical activity is due in part to disease duration, 1037

TEH LS, MANNING J, MOORE T, TULLY MP, O'REILLY D, JAYSON MIV. Sustained-release transdermal glyceryl trinitrate patches as a treatment for primary and secondary Raynaud's phenomenon, 636

TEH LS See also Davies E

TELFER J See McInnes IB

Temporal arteritis, macrophage derived granulocyte macrophage colony stimulating factor may augment HLA-DR expression on inflammatory cells. (A1) 151

TEN WOLDE S, DIJKMANS BAC, VAN ROOD JJ, CLAAS FHJ, DE VRIES RRP, HAZES JMW, VAN RIEL PCLM, VAN GESTEL A, BREEDVELD FC. HLA phenotypes and therapeutic response to parenteral gold in rheumatoid arthritis, (A1) 97

TEN WOLDE S. DUKMANS BAC, VAN ROOD JJ, CLAAS FHJ, DE VRIES RRP. HAZES JMW, VAN RIEL PLCM, VAN GESTEL A, BREEDVELD FC. Human leucocyte antigen phenotypes and gold-induced remissions in patients with rheumatoid arthritis, 343

TEN WOLDE S, DIJKMANS BAC, JANSSEN M, HERMANS J, VANDENBROUCKE JP, LAMERS CBHW. A double-blind placebo-controlled study of ranitidine for the prevention of recurrent peptic ulcer disease in rheumatoid arthritis patients on non-steroidal anti-inflammatory drugs, (A1) 94

Tender points

in graded back pain: more than pain in the back II, (A2) 38 number and extent of reported pain, (A1) 38

Tenidap, a cytokine-modulating anti-rheumatic drug, and diclofenac, a comparative study in rheumatoid arthritis: a 24-week analysis of a 1-year

clinical trial, 554 TENNANT A, FEAR J, HILLMAN M, CHAMBERLAIN MA. The Stanford Health Assessment Questionnaire: ordinal or interval scale?, (A2) 51

TENNANT A, McKenna SP. Conceptualizing and defining outcome, (E) 899 TENNANT A See also Hillman M

Tenosynovitis, fluoroquinolone-induced of the wrist mimicking de Quervain's disease, (L) 583

Tenoxicam, double blind comparison of the endoscopic and clinical effects of tenoxicam and diclofenac in rheumatoid arthritis, (A1) 95 Testosterone, protective role in adjuvant-induced arthritis, 1117

in vitro immunomodulatory effect in rheumatoid arthritis, (A1) 152 combined oral with intravenous clindamycin in rheumatoid arthritis: an open evaluation, (A1) 96

TGF-β1 induces myofibroblastic differentiation in human synovial fibroblasts in vitro: modulation by basic FGF, (A2) 29

Thalassaemia major, anterior knee pain in thalassaemia major treated with L1, a new iron chelating agent, (A1) 119

Therapeutic approaches for early rheumatoid arthritis. How early? How aggressive?, (S2) 87

Therapeutic effect

of an engineered human anti-tumour necrosis factor alpha antibody (CDP571) in rheumatoid arthritis, 334

of brefeldin on experimental reactive AA amyloidosis, (A1) 118

Therapeutic monoclonal antibodies, 707

Therapeutic ultrasound as a treatment of plantar heel pain, (A1) 42

They turned down my abstract again, (E) 303

Thiazide use effects on bone density: the Chingford study, (A1) 129

THOEN JE See Glennas A THOMAE K See Engelhardt G

THOMAS E See Croft PR

falls, (A1) 72

THOMAS JT See Hillarby MC THOMPSON PW, BHASKARAN V, EVANS C, EBRAHIM E. A randomized controlled trial of brisk walking in the prevention of osteoporosis and

THOMPSON PW, KIRWAN JR. Joints count: a review of old and new articular indices of joint inflammation, (E) 1003

THOMPSON PW, TAYLOR J, NEWTH J, FISHER A, JONES C. Identification of an at risk group for osteoporosis using heel ultrasound measurement in general practice, (A1) 37

THOMPSON PW See also Carr AJ; Dunne CA

THOMPSON RN See MacDonald AG

THOMSEN C See Nørregaard J

THOMSON EA See McInnes IB

THOMSON W, CROSS S, OLLIER W, CAMPBELL RDC. The MHC encoded TN-X gene may be important in RA, (A1) 140

THOMSON W, HARRISON BJ, PEPPER L, CHAKRAVARTY K, SYMMONS DPM. HLA associations of inflammatory polyarthritis developing postimmunisation, (A1) 76

THOMSON W See also Donn RP

Hughes syndrome: antiphospholipid antibodies move closer to in 1994, (E) 493

increased levels of β 2 glycoprotein-1 antigen and β 2 glycoprotein-1 binding antibodies are associated with a history of thromboembolic complications in patients with SLE and primary antiphospholipid syndrome, 1031

in patients with anti-cardiolipin antibodies is not associated with activated protein-C resistance due to factor VLeiden mutation, (A1) 15

Thrombotic events, analysis in antiphospholipid antibody positive lupus patients, (A1) 15

Thymic tumour, type 1 cryoglobulinaemia associated with: successful treatment with plasma exchange, 285 Thymoma, malignant, development of systemic lupus erythematosus after

chemotherapy and radiotherapy, 1175 Thyroid autoimmunity in systemic lupus erythematosus: the clinical significance of a fluctuating course, 361

Thyroid function tests, audit in a rheumatic disease clinic—results and impression that thyroxine excerbates RA, (A2) 21

Tiaprofenic acid, indomethacin and placebo in knee osteoarthritis: the long term outcome of therapy, (A2) 44

Tibial tubercles, does spiking predict osteoarthritis of the knee?, (A1) 36

Tibialis posterior tendon rupture, factor in planovalgus rheumatoid foot, 645 Tibiofemoral osteophyte in the anteroposterior radiograph predicts joint space narrowing in the patellofemoral joint, (A1) 136

TICKELL C. Environmental change and disease, 1105

TILLEY BC, HEYSE SP, RUSINOVA M, PILLEMER S, CLEGG DO. Global tests for multiple outcomes in rheumatoid arthritis trials, (A1) 89

TINAHONES FJ. COLLANTES E. C-SORIGUER FJ. GONZÁLEZ-RUIZ A. PINEDA M, AÑON J, SANCHEZ GUIJO P. Increased VLDL levels and diminished renal excretion of uric acid in hyperuricaemic-hypertriglyceridaemic

Tissue inhibitor of metalloproteinase (TIMP)

collagenase, collagenase-TIMP complexes and glycosaminoglycan levels in sequential samples of synovial fluid aspirated from patients with osteoarthritis, (A1) 138

effect of growth factors on production from human tendon fibroblasts in culture, (A1) 138

TN-x gene, MHC encoded may be important in RA, (A1) 140

TNF monoclonal antibodies, isotype variation. Is there a therapeutic difference in rheumatoid arthritis?, (A1) 153

TNF Nco-1 RFLP is not a predictive risk factor in rheumatoid arthritis, (A1)

TOBIAS JH, DALZELL N, CHILD AH. Assessment of bone mineral density in women with Marfan syndrome, 516

TOBIN J See Lim K

TODD C See O'Neill TW TODD P See Dolan AL

TOIVANEN P See Mäki-Ikola O; Roivainen A

TOKUCHIRA M See SATOH M

Tolerance to adjuvant arthritis induced by anti-CD4 monoclonal antibodies does not confer tolerance to type II collagen-induced arthritis, (A1) 65 TOMLINSON A See Craig JP

TONG W See Humphries J

Tonsillectomy and ankylosing spondylitis, (A2) 35

Topoisomerase-1, recombinant, T-cell response is independent of disease and of autoantibody status, (A1) 30

TORGERSON DJ, CAMPBELL MK, REID DM. Life-style, environmental and medical factors influencing peak bone mass in women, 620

TORGERSON DJ See also Francis RM

TÖRNWALL J. VIP in salivary glands in Sjögren's syndrome, (L) 891

Total and free methotrexate pharmacokinetics, with and without piroxicam, in rheumatoid arthritis patients, 421

TOUBI E See Buchanan NMM

TOWBIN H See Seitz M

Toxic epidermal necrolysis, pancytopenia and adult respiratory syndrome, (L) 798

Trabecular organisation measurement of the OA knee by fractal signature analysis of high definition macroradiographs, (A1) 121

Transient regional osteoporosis of the hip, (L) 296

Transporter genes, polypeptide (TAP) polymorphisms and genetic susceptibility to rheumatoid arthritis, 207

TRCB gene complex, mapping the genetic association with hypergammaglobulinaemic primary Sjögren's syndrome, (A1) 77

TREADWELL B See Rajapakse C

TRENNAM C See Naughton D

Triamcinolone, steroid myopathy induced by epidural injection, 385 Trivedi P See Vancheeswaran R

Trochlear and abducent palsies in a patient with Siögren's syndrome. (L) 484 Tropical infections, rheumatoid factor—associated cross reactive idiotypes,

TROUGHTON PR, EL MANZALAWI E, PLATT R, GAIKWAD P, BIRD HA, BASSIOUNI H, SALAHELDIN IB, BASSIOUNI M, WRIGHT V. Interleukin-8 (IL-8) and hypoxic reperfusion injury in rheumatoid arthritis and seronegative polyarthritis, (A1) 50

TROUGHTON PR See also Bradley SM

Tryptase, mast cell in the synovial fluid of patients with arthritis, (A1) 56 Tsal C-Y. Suppression of proteinuria and prolongation of survival in MRLlpr/lpr mice treated with sulphasalazine, (L) 85

TSAI C-Y See also Hsieh S-C TSAI S-T See Hsieh S-C

TSAI Y-Y See Hsieh S-C

Tuberculosis, and human immunodeficiency virus, multiple joint avascular necrosis-a rare but important cause, 387

Tuberculous muscle abscess in a patient with systemic lupus erythematosus,

TUCKER LB, MENON S, SCHALLER JG, ISENBERG DA. Adult-and childhoodonset systemic lupus erythematosus: a comparison of onset, clinical features, serology and outcome, 866

TUCKER R See Fisher M

TUDDENHAM EGD See Inamo Y

TULLY MP See Teh LS

Tumour necrosis factor, therapeutic effects of an engineered human antitumour necrosis factor alpha antibody (CDP571) in rheumatoid arthritis, 334

Tumour necrosis factor-α, neutrophil-derived: a potential mechanism for promotion of inflammation in procainamide-induced lupus, (A1) 3

TUNN E See Francombe J

Two distinct sites on the RNA polymerase-II large subunit are recognized by scleroderma autoantibodies but one site is unique to the phosphorylated (IIO) form, (A1) 31

Type 1 cryoglobulinaemia associated with a thymic tumour: successful treatment with plasma exchange, 285

UENO Y. A case of hepatitis C virus (HCV)-associated arthritis. Quantitative analysis of HCV RNA of the synovial fluid and the serum, (L) 691

UENO Y. Non-Hodgkin's lymphoma and polyarthritis, (L) 293 Ulcers of the small bowel induced by non-steroidal anti-inflammatory drugs in the rat: implications for clinical practice, 727

measurement of skin thickness in systemic sclerosis, 535

scans and indium-111 antimyosin antibody scans in the evaluation of patients with suspected polymyositis, (A2) 27

therapeutic as a treatment of plantar heel pain, (A1) 42

Uncoupling of oxidative phosphorylation

by non-steroidal anti-inflammatory drugs related to their pKA, (A2) 23 is the initiating factor in NSAID-induced gastrointestinal damage, (A1)

UNDERWOOD MR, DAWES P. Inflammatory back pain in primary care, 1074 UPPAL SS, PANDE I, SINGH G, KAILASH S, KAKKER R, KUMAR A, MEHRA NK, GIRI TK, SEKHARAN NG, MALAVIYA AN. Profile of HLA-B27related unclassifiable seronegative spondylarthropathy in females and its comparison with the profile in males, 137

UPPAL SS, PANDE IRA, KUMAR A, KAILASH S, SEKHARAN NG, ADYA CM, MALAVIYA AN. Adult onset Still's disease in Northern India: comparison with juvenile onset Still's disease, 429

Upper motor neuron effects on the cutaneous response to topical capsaicin, (A1) 53

Urate crystals, quantitative study of phagocytosis in the synovial fluid of asymptomatic joints of patients with gout, 724

Urate metabolism, effects of alcoholic beverages on urate metabolism in gout sufferers, 756

URIARTE E See Belzunegui J

Urinary levels of pyridinoline and deoxypyridinoline may be influenced by turnover of non-osseous tissues in patients with joint disease, (L) 1186 Urinary retinol binding protein excretion in SLE, (A1) 17

Urinary tract infection in patients with rheumatoid arthritis. A predictable problem?, (A1) 85

Urokinase and plasminogen activator inhibitor-1 levels in synovial fluids in normal, osteoarthritis and rheumatoid arthritis knees, (A1) 59

of a contrast enhanced MRI in the assessment of therapeutic response to a disease-modifying antirheumatic drug, 956

of focus group interview in the evaluation of patients with ankylosing spondylitis, 150

of a hand-pain diagram as a screening tool for carpal tunnel syndrome, (A1) 68

of intravenous prostacyclin to treat pulmonary hypertension associated with systemic sclerosis, (A1) 30

of three-phase bone scan in the diagnosis of repetitive strain injury, (A2)

Utility of lateral bone density measurements in osteoarthritis of the lumbar spine, (A1) 127

UTTLEY WS See Smith GW

Uveitis and pamidronate, (L) 479

VAGHMARIA A, EVANS D, GOLIK P, BAILEY I, BRADBURY S. Musculoskeletal problems in general practice: a prospective survey of consultations and management, (A2) 50

VAGHMARIA A See also Hav EM

VAISHNAW AK, WALPORT MJ, MORLEY BJ. Regulation of human complement component C4 gene expression, (A1) 156

VALE S See Kay LJ

Validation of a new method of assessing median nerve conduction at the wrist in carpal tunnel syndrome, (A1) 43

VALLS I See Saraux A

Value

of serological testing for Lyme disease in the UK, 132 of von Willebrand factor and angiotensin converting enzyme levels in

vasculitis, (A1) 22

VAN DE PUTTE LBA See Barrera P; van Gestel AM; Haagsma CJ; van Riel **PLCM**

VAN DEN BORNE BEEM, LANDEWÉ RBM, GOEI THE HS, VAN RUTHOVEN AWAM, BREEDVELD FC, DUKMANS BAC. Cyclosporine A in early rheumatoid arthritis patients: a 24 months chloroquine controlled follow up study, (A1) 99

VAN DEN ENDE CHM, HAZES JMW, LE CESSIE S, BREEDVELD FC, DIJKMANS BAC. Discordance between objective and subjective assessment of functional ability of patients with rheumatoid arthritis, 951

VAN DEN ENDE CHM, HAZES JMW, HERMANS J, BREEDVELD FC, DIJKMANS BAC. A randomized controlled trial of four different exercise programs in patients with rheumatoid arthritis, (A1) 93

VAN DER HEIDE A See Geenen R

VAN DER HEIJDE DMFM. Joint erosions and patients with early rheumatoid arthritis, (S2) 74

VAN DER KORST JK See Griep EN

VAN DER LAAR MAFJ See Kaandorp CJE VAN DER LINDEN Sj See Blaauw AAM

VAN DER MEER JWM See Barrera P; Griep EN VAN DER VLEUTEN CPM See Blaauw AAM

VAN DISTEL G See Boonen S

VAN GESTEL A See ten Wolde S

VAN GESTEL AM, LAAN RFJM, HAAGSMA CJ, VAN DE PUTTE LBA, VAN RIEL PLCM. Oral steroids as bridge therapy in rheumatoid arthritis patients starting with parenteral gold. A randomized double-blind placebocontrolled trial, 347

VAN GESTEL AM See also van Riel PLCM

VAN PAASSEN HC See Melief MJ

VAN RIEL PLCM. Bath Ankylosing Spondylitis Functional Index, (L) 794

VAN RIEL PLCM, VAN GESTEL AM, VAN DE PUTTE LBA. Long-term usage and side-effect profile of sulphasalazine in rheumatoid arthritis. (\$2) 40

VAN RIEL PLCM See also Barrera P; Haagsma CJ; ten Wolde S; Van Gestel AM

VAN RIJN HJM See Lems WF VAN RIJTHOVEN AWAM See van den Borne BEEM

VAN ROERMUND PM See van Valburg AA

VAN ROOD JJ See ten Wolde S

VAN SCHAARDENBURG D See Kaandorp CJE; Lycklama à Nijeholt GJ

VAN VALBURG AA, VAN ROERMUND PM, VERBOUT AJ, LAFEBER FPJG, BIJLSMA JWJ. Shortly after anterior cruciate ligament transection dogs develop characteristics similar to human moderate knee osteoarthritis,

VAN VUGT RM See Lems WF

VANCHEESWARAN R, BLACK CM, HASSON N, DAVID J, TRIVEDI P, WOO P. Differences between adult and childhood onset scleroderma, (A1) 29

VANDENBROUCKE JP See ten Wolde S; Vliet Vlieland TPM; Voskuyl AE VANDEVYER C, GEUSENS P, CASSIMAN J-J, RAUS J. Peptide transporter genes (TAP) polymorphisms and genetic susceptibility to rheumatoid arthritis, 207

Variation in sulphoxidation capacity in systemic lupus erythematosus patients and healthy controls of different ethnic origins, (A1) 6

VARSANI H See Rooney M

Vascular risk factors and rheumatoid arthritis, (A2) 7

Vasculitis

altered levels of soluble adhesion molecules, 814

cutaneous leucocytoclastic: a clinical and aetiological study, 942 factors associated with the development in rheumatoid arthritis: results of a case-control study, (A1) 86

fatal vasculitis following treatment with co-trimoxazole, (L) 84 rheumatoid, levels of cICAM-1, cICAM-3 and cE-selectin, (A1) 87

soluble immunoglobulin inhibits FCIII receptor-mediated neutrophil arrest to immune complexes under flow conditions: a possible mechanism of action in treatment, (A1) 24

value of von Willebrand factor and angiotensin converting enzyme levels. (A1) 22

Vasectomy, does it predispose to osteoporosis?, (A2) 42

VASERMAN N See Lioté F

Vasodilatation to acetylcholine, reduced in the finger microcirculation in primary Raynaud's phenomenon, (A1) 25

VEALE D See Maple C

VEALE DJ, BELCH JJF. Increased incidence of intermittent claudication in patients with systemic sclerosis: a prospective study, (A1) 26

VEALE DJ, D'ARRIGO C, CANDAL COUTO J, BELCH JJF. Soluble immuno-globulin inhibits FCI II receptor-mediated neutrophil arrest to immune complexes under flow conditions: a possible mechanism of action in treatment of vasculitis, (A1) 24

VEALE DJ, D'ARRIGO C, CANDAL COUTO J, WOOF J, GREER M, CREE I, BELCH JJF. FC/II receptor-mediated arrest under flow conditions is enhanced by Substance P, (A1) 52

VEALE DJ, Ho M, MORLEY KD. Sulphasalazine-induced lupus in psoriatic arthritis, 383

VEALE DJ, KIRK G, McLAREN M, BELCH JJF. Correlation of scleroderma skin score to change in soluble ICAM-1 levels reflects endothelial activation, (A1) 28

VEALE DJ See also Mulherin D

VECCHIO P, KAVANAGH R, HAZLEMAN BL, KING RH. Shoulder pain in a community-based rheumatology clinic, 440

VENABLES PJW, BOYD MT, WEERASINGHE D, BROOKES SM, JIAN-MEI L, MAINI RN. Antibodies to a human endogenous retrovirus envelope glycoprotein in rheumatoid arthritis, (A1) 154

VENABLES PJW See also Brookes SM; Rigby S

VENKOVSKY J See Macgregor AJ Venous thrombosis, cerebral and acquired protein S deficiency: an uncommon cause of headache in systemic lupus erythematosus, 572 Verbout AJ See van Valburg AA

VERCOULEN JHMM See Griep EN

Vertebral fractures

comparison of the effectiveness and cost of treatment in women, 1167 prevalence in patients with RA, treated with corticosteroids, (A1) 73 risk in DISH, (A1) 127

VERWEIJ CL See Kloppenburg M

VESTERGAARD-POULSEN P See Nørregaard J

VEYS EM See Mielants H VILAR MJP See Cury SE VILARDELL M See Marsal S

Villiaumey, Jaques, obituary, 592

VIP in salivary glands in Sjögren's syndrome, (L) 891

VIRGO S See Helsloot J

Visual loss, irreversible in dermatomyositis, 1179

Vitamin D receptor

early postmenopausal bone loss and polymorphisms of, (A1) 79 gene alleles and bone density in postmenopausal twins, (A1) 82 gene polymorphism in postmenopausal female athletes, (A1) 79

VLDL levels, increased and diminished renal excretion of uric acid in hyperuricaemic-hypertriglyceridaemic patients, 920

VLIET VLIELAND TPM, ZWINDERMAN AH, VANDENBROUCKE JP, BREEDVELD FC, HAZES JMW. In-patient treatment for active rheumatoid arthritis: clinical course and predictors of improvement, 847 Von Willebrand factor

and angiotensin converting enzyme level value in vasculitis, (A1) 22 and angiotensin converting enzyme release kinetics by endothelial cells, (A1) 23

and capillary microscopy, predictive value in systemic sclerosis (scleroderma)—a prospective study, (A1) 27
VOSKUYL AE, BREEDVELD FC, WESTEDT ML, ZWINDERMAN AH, VANDENBROUKE JP, HAZES JMW. Factors associated with the development of vasculitis in rheumatoid arthritis: results of a case-control study, (A1) 86

VOSKUYL AE, MARTIN S, MELCHERS I, ZWINDERMAN AH, WEICHSELBRAUN I, BREEDVELD FC. Levels of circulating intercellular adhesion molecule-1 and -3 but not circulating endothelial leukocyte adhesion molecule are increased in patients with rheumatoid arthritis, 311
VOSKUYL AE, MARTIN S, MELCHERS I, ZWINDERMAN AH, BREEDVELD FC.

Levels of clCAM-1, clCAM-3 and cE-selectin in patients with rheumatoid vasculitis, (A1) 87

VUORI K See Roivainen A VYSE T See Marguerie C

Vyse TJ See Bartók I

WADA T See Shingu M

WADSWORTH F. Gynaecological symptoms in fibromyalgia, (L) 888

WALDMANN H See Luqmani RA

WALKER D See Heslop P

WALKER DJ, CHANDLER C, ESSEX T, HESLOP P, STEAD H, KIFF P. Patient activity as an objective measure of NSAID therapy: an arthrotec and NUMACT study, (A2) 15

WALKER DJ See also Kay LJ; McDonagh JE; Wright D

Walking, randomized controlled trial in the prevention of osteoporosis and falls, (A1) 72

WALL P See Kavanagh R

WALLIS GA See Hillarby MC

WALLS AF See Buckley MG WALPORT MJ See Bartók I; Davies KA; Hogarth MB; Mackworth-Young CG; Marguerie C; Naughton M; Vaishnaw AK

WALSH S See Hardwick LJ; Loughlin J WALTER SD See Bosi Ferraz M

WALTERS C See Buckley MG

WALTERS G See De Lord D

WARD AB See Aung TS

WARD ME See Campbell F

WARD PS, BODEN T, WOODGER R, FOULKES R. Isotype variation of TNF monoclonal antibodies. Is there a therapeutic difference in rheumatoid arthritis?, (A1) 153

WARD R See Buckland-Wright JC

WARING RH See Griffiths B

WASSIF W See Smith EC

WATERHOUSE L, FARR M, JOHNSON AE, BACON PA. Disability in rheumatoid arthritis-it's relation to clinical and radiological courses over 3 years, (A2) 11

WATLING B See Bradshaw C

WATSON A See Winrow VR

WATT I See Jevtic V; Rogers J; Shahid MS

WATTS RA, CARRUTHERS DM, SCOTT DGI. Isolated nail fold rheumatoid vasculitis-a benign condition?, (A1) 68

WATTS RA, JOLIFFE VA, SCOTT DGI. Has polyarteritis nodosa been classified out of existence?, (A2) 25

WATTS RA, SCOTT DGI, CARRUTHERS DM, SYMMONS DPM. Incidence of systemic vasculitis in the Norwich Health Authority, (A1) 21 WATTS RA See also Carruthers DM

Weak association between self-reported dry eyes and dry mouth and objective tests of salivary and lacrimal function: a population study, (A1) 33

WEBLEY M See Srinivasan A

WEDLEY J See Dolan AL

WEEKS D See Hardwick LJ WEERASINGHE D See Venables PJW

Wegener's granulomatosis and CREST syndrome, (L) 483, (L) 484 WEICHSELBRAUN I See Voskuyl AE

WEINBLATT ME. Efficacy of methotrexate in rheumatoid arthritis, (S2) 43

WEISS JB See Fakhoury H

WEISSENSTEINER T See Campbell F

WELSH KI See Denton CP

WESTEDT ML See Voskuyl AE WESTHOVENS R See Boonen S WESTON A See Murphy K

- WESTON J See Athanassiou P
- What quality of life assessments should be used in RA?, (A2) 11
- WHEELER MJ See Hart DJ
- WHELLER SK See Perretti M
- Where is C7 produced?, (A1) 155 WHITEFIELD L, ISENBERG DA, BRAZIER DJ, FORBES J. Acquired Brown's syndrome in systemic lupus erythematosus, 1092
- WHITELOCK HC See Kennedy LG
- WHITING S See Jones M
- Why should rheumatoid arthritis patients recently recruited to sulphasalazine studies have milder disease than those enrolled over the previous decade? Does this trend influence outcome?, (A1) 98
- WHYTE J, EARNSHAW WC, CHAMPOUX JJ, PARKER LH, STEWART LJ, HALL ND, McHugh NJ. A T-cell response to recombinant topoisomerase-1 is independent of disease and of autoantibody status, (A1) 30
- WHYTE J, SORIANO E, EARNSHAW WC, McHugh NJ. Frequency of autoantibodies to a major epitope on the carboxyl terminal fragment of CENP-B in patients with autoimmune disease, 407
- WHYTE J See also Harvey GR
- WILKINSON I See Clunie G
- WILKINSON LS See Edwards JCW
- WILKINSON PC See Al-Mughales J
 WILLIAMS AS, PUNN YL, AMOS N, COOPER AM, WILLIAMS BD. The effect of liposomally conjugated methotrexate upon mediator release from human peripheral blood monocytes, 241
- WILLIAMS BD See Williams AS
- WILLIAMS DG, TAYLOR PC. Clonality of immunoglobulin production in rheumatoid synovium, (A1) 154
- WILLIAMS DG See also Adebajo AO
- WILLIAMS HJ. Overview of combination second-line or disease-modifying antirheumatic drug therapy in rheumatoid arthritis, (S2) 96
- WILLIAMS N See Herrick AL
- WILLIAMS P See James D; Prouse P; Scott DL
- WILLIAMS R See Gaffney K
- WILLIAMS S See Buckland-Wright JC; Smith EC
- WILLIS D, MOORE AR, GOWLAND G, WILLOUGHBY DA. OM-89 modulation of chronic inflammation: relevance to clinical use, 525
- WILLIS D, MOORE AR, GOWLAND G, WILLOUGHBY DA. Polyarthritis in the rat: effects of tolerance and sensitization to the bacterial extract OM-89 with a possible mode of action, 1135 WILLOUGHBY DA See Willis D
- WINFIELD J See James D; Prouse P
- WINKLER TH See Ehrenstein MR
- WINROW VR, MAPP PI, WATSON A, GAFFNEY K, BLAKE DR. 32kDa stress protein expression in arthritic synovia, (A1) 47 WINTERSBERGER W See Erlacher L
- WINYARD PG See Dai L; Jones HW; Nazhat NB; Zhang Z
- WISEMAN M See Cheung NT
- WOLLENHAUPT J See Stichtenoth DO
- WOLLHEIM FA See Scheja A
- WOLMAN R See Spector TD WOLTER D, See McRorie ER
- Wong D See Lau CS

- WONG R See Buckley MG WONG RWS See Lau CS WONG WM, COY SD, HOWELL WM, SMITH JL, CAWLEY MID. Interleukin-2 in psoriatic arthritis and seronegative spondylarthritis but not in rheumatoid arthritis synovium, (A1) 151
- Woo P. British Paediatric Rheumatology Group news, 59, 453, 865 Woo P See also Inamo Y; Murray JG; Rooney M; Vancheeswaran R
- WOOD C See Casey MC
- WOODGER R See Ward PS
- WOODWARD E See Doherty M
- WOODY J See Choy EHS
- Woof J See Veale DJ
- WOOLF AD See Deodhar AA; O'Neill TW
- WORDSWORTH BP See Brown MA; Gendi NST; Hardwick LJ WORDSWORTH P See Loughlin J

- WORKMAN E See Kavanagh R WORRALL JG, LOCKE IC, LEAKER B, BLACK CM, CAMBRIDGE G. Autoantibodies to myeloperoxidase in systemic sclerosis, (A2) 24
- WORRALL JG See also Henderson KJ
- WRIGGLESWORTH J See Rafi S

- WRIGGLESWORTH JM See Mahmud T
- WRIGHT A See Hillman M
- WRIGHT D, BARROW S, FISHER AD, HORSLEY SD, JAYSON MIV. Influence of physical, psychological and behavioural factors on consultations for back pain, 156
- WRIGHT D, CHANDLER C, ESSEX T, HESLOP P, STEAD H, KIFF P, WALKER DJ. Diurnal variation in 24 hour ambulatory recordings of patient activity is different in rheumatoid and osteoarthritis, (A2) 16
- WRIGHT EA See Daltroy LH
- WRIGHT ED. Discitis caused by Corynebacterium pseudodiphtheriticum
- following ear, nose and throat surgery, (L) 585
 WRIGHT M, JOBANPUTRA P, BAVINGTON C, SALTER DM, NUKI G. Signal transduction pathways in human chondrocytes activated by intermittent pressurisation, (A1) 58
- WRIGHT V. Historical overview of non-steroidal anti-inflammatory drugs,
- WRIGHT V. Introduction: decisions in non-steroidal anti-inflammatory prescribing, (S1) 1
- WRIGHT V See also Troughton PR
- Wrist
 - fluoroquinolone-induced tenosynovitis mimicking de Quervain's disease, (L) 583
 - predominance in seronegative rheumatoid arthritis: fact, fiction or natural history, (A2) 9 surgery to joint, 274
- WURZNER R See Naughton M
- WYLIE G, APPELBOOM T, BOLTEN W, BREEDVELD FC, FEELY J, LEEMING MRG, LE LOET X, MANTHORPE R, MARCOLONGO R, SMOLEN J. Comparative study of tenidap, a cytokine-modulating anti-rheumatic drug, and diclofenac in rheumatoid arthritis: a 24-week analysis of a 1year clinical trial, 554
- WYNN-JONES CH See Aung TS
- Xanthine oxidase, expression of active, by synovial microvascular endothelial cells in culture, (A1) 51
- Xerostomia prevalence in a population with primary generalised osteoarthritis, (A1) 132
- YAMADA H See Kuroki S
- YAMAGUCHI M See Kuroki S
- YANNI G See Cauli A: Pitzalis C
- YAP PL See Jobanputra P
- YAQUB R, FAWTHROP F, BAYLISS M, DOHERTY M. Levels of chondroitin sulphate and keratan sulphate epitopes, glycosaminoglycans and hyaluronan in normal knee synovial fluid, (A1) 55
- YASUTAKE C See Shingu M
- YATES DAH. Obituary of AT Richardson, 188
 YEAP SS, KEATING NA, HOSKING DJ. Should close relatives of patients with osteoporosis be investigated?, (A1) 73
- YEAP SS, KEATING NA, KENNEDY TD. Assessment of referral guidelines to a bone densitometry service, (A1) 37
- YEO LMW, SWABY DSA, SITUNAYAKE RD, MURRAY PI. Irreversible visual loss in dermatomyositis, 1179
- YOUINOU P See Berthelot JM; Jorgensen C; SARAUX A
- Young A. Short-term outcomes in recent-onset rheumatoid arthritis, (S2) 79 Young A See also James D; Prouse P
- YOUNG S See Robinson C
- Yu C-L See Hsieh S-C; Lai N-S
- YII H-S See Hsieh S-C
- ZEIDLER H See Stichtenoth DO
- ZEUTHEN EL See Birk MA
- ZHANG Z, FREARS E, NAZHAT NB, BLAKE DR, STEVENS CR, WINYARD PG. Oxidative inactivation of human alpha-1-proteinase inhibitor by simultaneous generation of superoxide and nitric oxide, (A1) 48
- ZHANG Z See also Dai L; Jones HW
- ZINDERMAN AH See Voskuyl AE
- ZULFIQUAR A See Segasothy M
 ZWEDE M, KITAS GD, LUQMANI R, BACON PA. Kinetics of von Willebrand factor and angiotensin converting enzyme release by endothelial cells, (A1) 23
- ZWEDE M See also Kitas GD
- ZWINDERMAN AH See Vliet Vlieland TPM: Voskuvl AE

